Triazabicyclodecene: an Effective Isotope Exchange Catalyst in CDCl₃

Supporting Information

Cyrille Sabot,[†] Kanduluru Ananda Kumar,[†] Cyril Antheaume,[‡] Charles Mioskowski^{†}*

Laboratoire de Synthèse Bio-Organique, UMR 7175/LC1-CNRS, and Service Commun de
RMN, IFR85, Université Louis Pasteur de Strasbourg, 74 route du Rhin, B.P 60024, 67401
Illkirch cedex, France

mioskow@aspirine.u-strasbg.fr

Table of contents

Experimental procedures	S4
General considerations	S4
Representative procedure for deuteration reaction	
(compounds in Tables 2 and 3)	S4-S6
2-Acetoxyacetophenone 1i	S5
<i>trans</i> -2-Dimethylamino-4,5-diphenylimidazoline 2	S5-S6
Characterization data (¹H/¹³C)	S7
2-Acetoxyacetophenone 1i	S7

S-Methyltetramethylthiuronium iodide 2'	S8
<i>trans</i> -2-Dimethylamino-4,5-diphenylimidazoline 2	S9
Spectral data (¹H/²H)	S10
TMG	S10
4'-Methoxyacetophenone- <i>d</i> ₃ 1a-d₃	S11
Acetophenone- <i>d</i> ₃ 1b-d₃	S12-S15
4'-Nitroacetophenone- <i>d</i> ₃ 1c-d₃	S16
2-Methoxyacetophenone- <i>d</i> ₂ 1d-d₂	S17
2-Methylacetophenone- <i>d</i> ₂ 1e-d₂	S18
2-Chloroacetophenone- <i>d</i> ₂ 1f-d₂	S19
2,2-Dimethylacetophenone- <i>d</i> ₁ 1g-d₁	S20
1-Phenylpropane-1,2-dione- <i>d</i> ₃ : 1h-d₃	S21
2-Acetoxyacetophenone- <i>d</i> ₂ 1i-d₂	S22
Naphthylmethylketone- <i>d</i> ₃ 1j-d₃	S23
2-Acetylpyridine- <i>d</i> ₃ 1k-d₃	S24
2-Acetylthiophene- <i>d</i> ₃ 1l-d₃	S25
Benzylacetone- <i>d</i> ₅ 1m-d₅	S26-S28
Cyclodecanone- <i>d</i> ₃ 1n-d₄	S29-S31
1-Phenylacetylene- <i>d</i> ₁ 3a-d₁	S32
4-Phenylbut-1-yne- <i>d</i> ₁ 3b-d₁	S33
1-Phenyl-2-methylacetylene- <i>d</i> ₃ 3c	S34
2-Ethynyl-1-methylimidazole 3d-d₁	S35-S36
Methyl-1-phenylacetate 4-d₂	S37
3-Nitrofluorene 5-d₂	S38
Indene 6-d₃	S39

^1H NMR study	S40
References	S41

Experimental procedures

General considerations

All substrates were commercially available and used without further purification except 2-acetoxyacetophenone **1i** synthesized from commercially available 2-hydroxyacetophenone. The *trans*-2-dimethylamino-4,5-diphenylimidazoline **2** was obtained in two steps from commercially available tetramethylthiourea. Deuterated chloroform used for deuterium incorporation reactions, contained D, 99.80% and H₂O<0.01%. ¹H and ¹H-{²H} NMR spectral data were recorded at 600 or 300 MHz, on spectrometers using a software pulse program library. CDCl₃ was used as lock and CHCl₃ as internal standard (7.27 ppm). ²H NMR spectral data were recorded at 46 MHz and chemical shifts are reported relative to CDCl₃ (7.27 ppm). HRMS analysis was performed in positive electrospray mode.

Representative procedure for deuteration reaction (compounds in Tables 2 and 3):

For compound **1b**:

To a solution of 10 mol % of TBD (10 mg, 0.072 mmol) in 3 mL of CDCl₃, was added **1b** (84 μL, 0.72 mmol). The reaction mixture was stirred at room temperature for 12 hours and quenched with 1N HCl (1 mL). The organic layer was washed with water (2 x 2 mL), brine (1 mL), dried over anhydrous Na₂SO₄ and filtered. Filtrate was concentrated to afford **1b-d₃**. The total incorporation yield was determined by ¹H NMR spectroscopy relative to the intensity of a non exchangeable proton in the molecule, or by GC-MS. For example, the global deuterium incorporation of compound **1b-d₃** determined by GC-MS, was calculated as follow: $1/3 \times 5.3\% + 2/3 \times 28.5\% + 3/3 \times 65.6\% = 86.4\%$ of total incorporation. Moreover,

the site of deuterium incorporation in the substrate was determined by ^2H NMR spectroscopy relative to CHCl_3 (7.27 ppm).

2-Acetoxyacetophenone **1i**

1i

To a solution of 2-hydroxyacetophenone (1 g , 9.3 mmol) dissolved in pyridine (40 mL), dried over 3Å molecular sieves beforehand, was added acetyl chloride (667 μL , 9.3 mmol) by dropwise at 0 °C. The reaction mixture was stirred at room temperature for 8 hours. The solution was filtered, and filtrate concentrated. The residue was dissolved with 1N HCl and the organic layer was extracted twice with diethylether. The combined organic layers were washed with brine and dried over anhydrous sodium sulfate, filtered and concentrated to afford a brown residue. The crude product was purified by chromatography on a silica gel column eluting with 20% ethyl acetate in cyclohexane to give 1.320 g (80% yield) of **1i** as a yellow oil. Analytical data were in accordance with the previous literature report.¹ R_f = 0.44 (UV, Vanilline) in 30% EtOAc/ Cyclohexane. ^1H NMR (CDCl_3 , 300 MHz): δ 2.24 (s, 3H), 5.35 (s, 2H), 7.45-7.55 (t, J =7.5 Hz, 2H, Har), 7.60-7.70 (t, J =7.5 Hz, 1H, Har), 7.90-8.00 (t, J =7.5 Hz, 2H, Har). ^{13}C NMR (CDCl_3 , 75 MHz): δ 20.8, 66.3, 128.0, 129.1, 134.1, 134.4, 170.6, 192.4. IR (KRS-5): 2936, 1751, 1705, 1598, 1450, 1421, 1375, 1220. CIMS 179 $[\text{M}+\text{H}^+]$, 196 $[\text{M}+\text{NH}_4^+]$.

trans-2-Dimethylamino-4,5-diphenylimidazoline **2**

Step 1: S-Methyltetramethylthiuronium iodide **2'**

To a solution of tetramethylthiourea (1 g, 7.57 mmol) in 8 mL of dried methanol at 0 °C under argon, was added methyl iodide (870 μ L, 14.09 mmol) by dropwise. After 16 h stirring at room temperature, the solvent was removed under reduced pressure and 2,065 g (99% yield) of **2'** as a yellow solid was obtained (hygroscopic).² RMN ^1H (300 MHz ; CDCl_3) δ (ppm): 2.62 (s, 3H), 3.35 (s, 12H). RMN ^{13}C (75 MHz ; CDCl_3) δ (ppm): 18.6, 45.3, 176.9.

Step 2: *trans*-2-Dimethylamino-4,5-diphenylimidazoline **2**

To a solution of *trans*-1,2-diphenylethylenediamine (1 g, 4.71 mmol) in 14 mL of dried acetonitrile, at 0 °C, under argon, was added S-methyltetramethylthiuronium iodide **2'** (1.808 g, 6.59 mmol). The solution was stirred at room temperature for 2 h and further 2 h at reflux. After cooling, the solution was diluted with 50 mL of dichloromethane and treated by a solution 5% of sodium carbonate (2 x 50 mL). The organic layer was dried over potassium carbonate, filtered and filtrate concentrated. After recrystallization (dichloromethane and a minimum of *n*-hexane), 953 mg (76% yield) of a white solid **2** was obtained. mp 202-203 °C RMN ^1H (300 MHz ; CDCl_3) δ (ppm): 3.02 (s, 6H), 4.69 (s, 2H), 4.78 (bs, 1H), 7.15-7.48 (m, 10H). RMN ^{13}C (50 MHz ; CDCl_3) δ (ppm): 38.8, 74.0, 126.9, 127.7, 128.8, 143.8, 162.3. IR (KRS-5) ν (cm^{-1}): 3028, 2866, 2827, 1667, 1609, 1485, 1451. MS (Electrospray) 266.1612 m/z (266.1657 calcd for $\text{M}+\text{H}$).

Characterization data ($^1\text{H}/^{13}\text{C}$)

Spectral data ($^1\text{H}/^2\text{H}$)

GC

MS

GC

MS

GC

1m

MS

GC

1m-d₅

MS

GC

MS

GC

MS

Reaction of **2d** with TBD

Reaction **2d** with PSTBD

Mechanistic study

References

¹ Itsuno, I.; *J. Chem. Soc. Perkin Trans. I* **1985**, 2039.

² Fujisaki, S.; Fujiwara, I.; Norisue, Y.; Kajigaeshi, S. *Bull. Chem. Soc. Jpn.* **1985**, 58, 2429.