Supporting Information

Triethylborane-Induced Intermolecular Radical Addition to Ketimines

Hideto Miyabe, Yousuke Yamaoka, and Yoshiji Takemoto*

Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

e-mail: takemoto@pharm.kyoto-u.ac.jp

Table of contents

Experimental procedure and characterization data for **3-13b**: S1-7.

¹H and ¹³C NMR spectrum of all obtained compounds: S8-S53.

Methyl 2-[(E)-2-Benzoylhydrazono]propanoate (3). According to literature, ketimine **3** was prepared as a colorless crystal.¹ As a colorless crystal. mp 195-197 °C (AcOEt/hexane). IR (CHCl₃) 1706 cm⁻¹. ¹H NMR (CDCl₃) δ 9.07 (1H, s), 7.90 (2H, br m), 7.58 (1H, t, *J*=7.6 Hz), 7.48 (2H, t, *J*=7.6 Hz), 3.87 (3H, s), 2.21 (3H, s). MS (EI⁺): *m/z* 221 (M+H⁺, 92), 221 (100). Anal. Calcd for C₁₁H₁₂N₂O₃: C, 59.99; H, 5.49; N, 12.72. Found: C, 59.83; H, 5.41; N, 12.72.

Procedure for radical addition to ketimine 3. To a solution of ketimine **3** (100 mg, 0.45 mmol) in CH₂Cl₂ (4.5 mL) were added R²I (13.6 mmol) and Et₃B (1.0 M in hexane, 1.1 mL, 1.1 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 15 min, the reaction mixture was diluted with saturated NaHCO₃ and then extracted with AcOEt. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=2:1)

afforded **4a-d**. The ethyl radical addition reaction was performed in the absence of R²I.

Methyl 2-(2-Benzoylhydrazinyl)-2-methylbutanoate (4a). A colorless oil. IR (CHCl₃) 1728, 1677 cm⁻¹. ¹H NMR (CDCl₃) δ 7.92 (1H, br s), 7.74 (2H, d, *J*=7.0 Hz), 7.51 (1H, t, *J*=7.0 Hz), 7.43 (2H, t, *J*=7.0 Hz), 5.04 (1H, br s), 3.76 (3H, s), 1.81 (2H, q, *J*=7.6 Hz), 1.38 (3H, s), 0.96 (3H, t, *J*=7.6 Hz). ¹³C NMR (CDCl₃) δ 175.8, 166.8, 132.9, 131.7, 128.7, 126.9, 65.7, 52.2, 29.9, 20.7, 8.2. MS (EI⁺): *m/z* 250 (M⁺, 6). HRMS (EI⁺): calcd for C₁₃H₁₈N₂O₃ (M⁺): 250.1317, Found: 250.1323.

Methyl 2-(2-Benzoylhydrazinyl)-2,3-dimethylbutanoate (4b). A colorless crystal. mp 85-87 °C (AcOEt/hexane). IR (CHCl₃) 1727, 1657 cm⁻¹. ¹H NMR (CDCl₃) δ 7.86 (1H, br s), 7.73 (2H, d, *J*=7.3 Hz), 7.51 (1H, t, *J*=7.3 Hz), 7.43 (2H, t, *J*=7.3 Hz), 5.13 (1H, br s), 3.75 (3H, s), 2.15 (1H, m), 1.33 (3H, s), 1.00 (6H, br d, *J*=7.0 Hz). ¹³C NMR (CDCl₃) δ 175.9, 166.6, 133.0, 131.7, 128.7, 126.8, 68.4, 52.1, 33.8, 17.5, 17.2, 16.9. MS (EI⁺): *m/z* 264 (M⁺, 3), 105 (100). Anal. Calcd for C₁₄H₂₀N₂O₃: C, 63.62; H, 7.63; N, 10.60. Found: C, 63.50; H, 7.48; N, 10.60.

Methyl 2-(2-Benzoylhydrazinyl)-2-cyclohexylpropanoate (4c). A colorless crystal. mp 67-69 °C (AcOEt/hexane). IR (CHCl₃) 1726, 1657 cm⁻¹. ¹H NMR (CDCl₃) δ 7.84 (1H, br s), 7.73 (2H, d, *J*=7.6 Hz), 7.50 (1H, t, *J*=7.6 Hz), 7.43 (2H, t, *J*=7.6 Hz), 5.12 (1H, br s), 3.75 (3H, s), 1.92-1.65 (6H, m), 1.34 (3H, s), 1.30-1.10 (5H, m). ¹³C NMR (CDCl₃) δ 175.9, 166.6, 133.0, 131.7, 128.7, 126.8, 68.5, 52.1, 44.3, 27.5, 27.1, 26.6, 26.5, 26.2, 18.0. MS (EI⁺): *m/z* 304 (M⁺, 2), 105 (100). HRMS (EI⁺): calcd for C₁₇H₂₄N₂O₃ (M⁺): 304.1787, Found: 304.1783.

Methyl 2-(2-Benzoylhydrazinyl)-2,3,3-trimethylbutanoate (4d). A colorless crystal. mp 91-94 °C (AcOEt/hexane). IR (CHCl₃) 1724, 1657 cm⁻¹. ¹H NMR (CDCl₃) δ 7.71 (2H, d, *J*=7.3 Hz), 7.57 (1H, br s), 7.51 (1H, t, *J*=7.3 Hz), 7.43 (2H, t, *J*=7.3 Hz), 5.22 (1H, br s), 3.77 (3H, s), 1.36 (3H, s), 1.09 (9H, s). ¹³C NMR (CDCl₃) δ 175.2, 166.9, 133.1, 131.8, 128.8, 126.9, 71.6, 51.9, 36.6, 25.9, 16.4. MS (EI⁺): *m/z* 278 (M⁺, 0.5), 105 (100). Anal. Calcd for C₁₅H₂₂N₂O₃: C, 64.73; H, 7.97; N, 10.06. Found: C, 64.84; H, 7.90; N, 10.01.

3-(Benzyloxy)imino-1-methylindolin-2-one (5). To a solution of *N*-methylisatin (1.0 g, 6.2 mmol) in pyridine (10 mL) was added BnONH₂·HCl (0.99 g, 6.2 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 2 h, the reaction mixture was diluted with Et₂O and then washed with water. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by recrystallization (hexane-AcOEt) afforded **5** (1.35 g 82%) as a yellow crystal. mp 117-119 °C (AcOEt/hexane). IR (CHCl₃) 1722, 1610 cm⁻¹. ¹H NMR (CDCl₃) δ 7.54 (1H, d, *J*=7.3 Hz), 7.46 (2H, d, *J*=7.0 Hz), 7.39-7.28 (4H, m), 7.03 (1H, t, *J*=7.3 Hz), 6.78 (1H, d, *J*=7.9 Hz), 5.46 (2H, s), 3.21 (3H, s). ¹³C NMR (CDCl₃) δ 163.6, 144.5, 144.0, 136.2, 132.5, 128.6, 128.4 (2C), 127.9, 123.0, 115.7, 108.3, 79.3, 26.0. MS (EI⁺): *m/z* 266 (M⁺, 11), 91 (100). Anal. Calcd for C₁₆H₁₄N₂O₂: C, 72.16; H, 5.30; N, 10.52. Found: C, 72.40; H, 5.35;

N, 10.52.

3-(2,2-Diphenylhydrazono)-1-methylindolin-2-one (6). According to literature, ketimine **6** was prepared as a yellow crystal.² mp 175-178 °C (AcOEt/hexane). IR (CHCl₃) 1706, 1606 cm⁻¹. ¹H NMR (CDCl₃) δ 7.40-7.20 (10H, m), 7.13 (1H, t, *J*=7.6 Hz), 6.74 (1H, d, *J*=7.6 Hz), 6.49 (1H, t, *J*=7.6 Hz), 5.51 (1H, d, *J*=7.6 Hz), 3.28 (3H, s). ¹³C NMR (CDCl₃) δ 165.9, 146.0, 144.2, 133.1, 129.8, 129.3, 125.9, 125.4, 123.4, 121.4, 115.9, 107.8, 26.0. MS (EI⁺): *m/z* 327 (M⁺, 19), 168 (100). Anal. Calcd for C₂₁H₁₇N₃O: C, 77.04; H, 5.23; N, 12.84. Found: C, 77.19; H, 5.35; N, 12.74.

3-(2-Benzoylhydrazono)-1-methylindolin-2-one (7). According to literature, ketimine **7** was prepared as a yellow crystal.³ mp 184-186 °C (AcOEt/hexane). IR (CHCl₃) 1731, 1685 cm⁻¹. ¹H NMR (CDCl₃) δ 8.02 (2H, d, *J*=7.6 Hz), 7.88 (1H, br s), 7.61 (1H, t, *J*=7.6 Hz), 7.53 (2H, t, *J*=7.6 Hz), 7.42 (1H, t, *J*=7.6 Hz), 7.18 (1H, t, *J*=7.6 Hz), 6.91 (1H, d, *J*=7.6 Hz), 3.32 (3H, s). One peak due to NH was missing. ¹³C NMR (CDCl₃) δ 163.9, 161.7, 143.3, 137.3, 132.7, 132.0, 131.5, 128.9, 127.7, 123.5, 121.8, 119.5, 108.9, 25.6. MS (EI⁺): *m/z* 279 (M⁺, 4), 105 (100). Anal. Calcd for C₁₆H₁₃N₃O₂: C, 68.81; H, 4.69; N, 15.05. Found: C, 68.69; H, 4.76; N, 15.06.

Procedure for ethyl radical addition to ketimines 5 and 6. To a solution of ketimine **5** or **6** (0.2 mmol) in CH₂Cl₂ (2 mL) was added Et₃B (1.0 M in hexane, 0.5 mL, 0.5 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 1 h, Et₃B (1.0 M in hexane, 0.5 mL, 0.5 mmol) was added to the reaction mixture thrice. After being stirred at the same temperature for 11 h, the reaction mixture was diluted with saturated NaHCO₃ and then extracted with AcOEt. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=4:1, 2-fold development) afforded **8Aa-9Ba**.

Procedure for radical addition to ketimine 7. To a solution of ketimine **7** (55.8 mg, 0.2 mmol) in CH₂Cl₂ (2 mL) were added R²I (6.0 mmol) and Et₃B (1.0 M in hexane, 0.5 mL, 0.5 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 1 h, Et₃B (1.0 M in hexane, 0.5 mL, 0.5 mmol) was added to the reaction mixture. After being stirred at the same temperature for 2 h, the reaction mixture was diluted with saturated NaHCO₃ and then extracted with AcOEt. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=4:1, 2-fold development) afforded **10Aa-10Ad**. The ethyl radical addition reaction was performed in the absence of R²I.

3-(Benzyloxyamino)-3-ethyl-1-methylindolin-2-one (8Aa). A yellow oil. IR (CHCl₃) 1714, 1615 cm⁻¹. ¹H NMR (CDCl₃) δ 7.39-7.30 (2H, m), 7.28-7.19 (3H, m), 7.11 (1H, t, *J*=7.6 Hz), 6.99-6.93 (2H, m), 6.87 (1H, d, *J*=7.6 Hz), 6.12 (1H, br s), 4.41 (1H, d, *J*=11.6 Hz), 4.35 (1H, d, *J*=11.6 Hz), 3.21 (3H, s), 1.78 (1H, m), 1.69 (1H, m), 0.63 (3H, t, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 178.1, 144.8, 137.4, 129.0, 128.8, 128.3, 128.1, 127.6, 123.8,

122.6, 108.0, 77.1, 70.2, 26.8, 26.1, 7.2. MS (EI⁺): m/z 296 (M⁺, 5), 174 (100). HRMS (EI⁺): calcd for C₁₈H₂₀N₂O₂ (M⁺): 296.1525, Found: 296.1529.

3-(N-(Benzyloxy)-N-ethylamino)-3-ethyl-1-methylindolin-2-one (8Ba). A yellow oil. IR (CHCl₃) 1711, 1612 cm⁻¹. ¹H NMR (CDCl₃) δ 7.43 (1H, d, *J*=7.6 Hz), 7.32-7.20 (4H, m), 7.11 (2H, d, *J*=6.1 Hz), 7.06 (1H, t, *J*=7.6 Hz), 6.80 (1H, d, *J*=7.6 Hz), 4.77 (1H, d, *J*=10.4 Hz), 4.64 (1H, d, *J*=10.4 Hz), 3.16 (3H, s), 2.93 (1H, m), 2.79 (1H, m), 2.23 (1H, m), 2.16 (1H, m), 1.22 (3H, t, *J*=7.0 Hz), 0.59 (3H, t, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 175.8, 144.2, 136.9, 128.7, 128.2 (2C), 127.7, 125.1, 122.3, 107.7, 78.6, 74.9, 47.6, 27.9, 25.8, 13.4, 7.8. One carbon peak was missing due to overlapping. MS (EI⁺): m/z 324 (M⁺, 2), 265 (100). HRMS (EI⁺): calcd for C₂₀H₂₄N₂O₂ (M⁺): 324.1838, Found: 324.1835.

3-(2,2-Diphenylhydrazinyl)-3-ethyl-1-methylindolin-2-one (9Aa). A colorless crystal. mp 162-164 °C (AcOEt/hexane). IR (CHCl₃) 1712 cm⁻¹. ¹H NMR (CDCl₃) δ 7.35 (1H, d, *J*=7.5 Hz), 7.25 (1H, m), 7.17 (4H, br t, *J*=8.2 Hz), 7.00-6.90 (3H, m), 6.80 (4H, br d, *J*=8.2 Hz), 6.67 (1H, d, *J*=7.5 Hz), 4.88 (1H, s), 2.72 (3H, s), 1.99 (1H, m), 1.76 (1H, m), 0.76 (3H, t, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 178.6, 147.9, 144.2, 128.8, 128.7, 127.9, 124.5, 122.5, 122.4, 120.6, 107.7, 68.6, 30.2, 25.6, 7.5. MS (EI⁺): m/z 357 (M⁺, 11), 183 (100). Anal. Calcd for C₂₃H₂₃N₃O+1/5H₂O: C, 76.51; H, 6.53; N, 11.64. Found: C, 76.70; H, 6.58; N, 11.60.

3-(1-Ethyl-2,2-diphenylhydrazinyl)-3-ethyl-1-methylindolin-2-one (9Ba). An amorphous substance. IR (CHCl₃) 1708, 1612 cm⁻¹. ¹H NMR (CDCl₃) δ 7.22-7.04 (9H, m), 6.89-6.83 (3H, m), 6.77 (1H, br t, *J*=7.6 Hz), 6.63 (1H, br d, *J*=7.6 Hz), 3.40 (1H, m), 3.32 (1H, m), 2.86 (3H, s), 2.16 (1H, m), 2.06 (1H, m), 1.09 (3H, t, *J*=7.3 Hz), 0.72 (3H, t, *J*=7.3 Hz). MS (FAB⁺): m/z 385 (M⁺, 5), 211 (100). HRMS (FAB⁺): calcd for C₂₅H₂₇N₃O (M⁺): 385.2154, Found: 385.2156.

3-(2-Benzoylhydrazinyl)-3-ethyl-1-methylindolin-2-one (10Aa). A colorless crystal. mp 172-174 °C (AcOEt/hexane). IR (CHCl₃) 1710, 1614 cm⁻¹. ¹H NMR (CDCl₃) δ 7.68 (1H, br s), 7.60 (2H, br d, *J*=7.5 Hz), 7.47 (2H, m), 7.41-7.32 (3H, m), 7.12 (1H, br t, *J*=7.5 Hz), 6.85 (1H, br d, *J*=7.5 Hz), 5.51 (1H, br s), 3.23 (3H, s), 2.19-1.95 (2H, m), 0.69 (3H, t, *J*=7.6 Hz). ¹³C NMR (CDCl₃) δ 177.3, 166.8, 144.0, 132.6, 131.5, 129.3, 128.4, 127.9, 127.0, 124.3, 122.8, 107.9, 69.5, 27.7, 26.0, 7.58. MS (EI⁺): m/z 309 (M⁺, 4), 174 (100). Anal. Calcd for C₁₈H₁₉N₃O₂: C, 69.88; H, 6.19; N, 13.58. Found: C, 69.62; H, 6.31; N, 13.35.

3-(2-Benzoyl-1-ethylhydrazinyl)-3-ethyl-1-methylindolin-2-one (10Ba). An amorphous substance. IR (CHCl₃) 1691, 1614 cm⁻¹. ¹H NMR (CDCl₃) δ 8.52 (1H, br s), 7.92 (2H, d, *J*=7.6 Hz), 7.59-7.44 (4H, m), 7.36 (1H, t, *J*=7.6 Hz), 7.18 (1H, t, *J*=7.6 Hz), 6.88 (1H, d, *J*=7.6 Hz), 3.23 (3H, s), 2.61 (1H, m), 2.39 (1H, m), 2.21 (1H, m), 2.11 (1H, m), 1.02 (3H, t, *J*=7.0 Hz), 0.64 (3H, d, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 179.1, 166.7, 143.7, 133.9, 131.7, 129.3, 128.8, 128.7, 127.3, 124.9, 123.7, 108.5, 72.5, 48.1, 28.6, 25.7, 12.5, 7.6. MS (EI⁺):

m/z 337 (M^+ , 2), 105 (100). HRMS (EI^+): calcd for $C_{20}H_{23}N_3O_2$ (M^+): 337.1790, Found: 337.1795.

3-(2-Benzoylhydrazinyl)-3-isopropyl-1-methylindolin-2-one (10Ab). A colorless crystal. mp 176-178 °C (AcOEt/hexane). IR ($CHCl_3$) 1709, 1613 cm^{-1} . 1H NMR ($CDCl_3$) δ 7.99 (1H, br s), 7.58 (2H, d, $J=7.3$ Hz), 7.49-7.41 (2H, m), 7.38-7.28 (3H, m), 7.08 (1H, br t, $J=7.6$ Hz), 6.84 (1H, br d, $J=7.6$ Hz), 5.56 (1H, br d, $J=5.8$ Hz), 3.21 (3H, s), 2.40 (1H, m), 1.16 (3H, d, $J=6.7$ Hz), 0.67 (3H, d, $J=6.7$ Hz). ^{13}C NMR ($CDCl_3$) δ 178.0, 166.4, 144.4, 132.8, 131.6, 129.4, 128.5, 126.9, 126.5, 125.5, 122.6, 108.0, 71.6, 32.5, 26.0, 16.6, 16.3. MS (EI^+): m/z 323 (M^+ , 4), 280 (100). Anal. Calcd for $C_{19}H_{21}N_3O_2$: C, 70.57; H, 6.55; N, 12.99. Found: C, 70.30; H, 6.46; N, 12.82.

3-(2-Benzoyl-1-isopropylhydrazinyl)-3-isopropyl-1-methylindolin-2-one (10Bb). An amorphous substance. IR ($CHCl_3$) 1689, 1611 cm^{-1} . 1H NMR ($CDCl_3$) δ 8.94 (1H, br s), 7.98 (2H, d, $J=7.0$ Hz), 7.55-7.43 (4H, m), 7.39 (1H, t, $J=7.6$ Hz), 7.16 (1H, t, $J=7.6$ Hz), 6.91 (1H, d, $J=7.6$ Hz), 3.22 (3H, s), 3.10 (1H, m), 2.50 (1H, m), 1.33 (3H, d, $J=6.7$ Hz), 1.08 (3H, d, $J=6.7$ Hz), 0.75 (3H, d, $J=6.7$ Hz), 0.41 (3H, d, $J=6.7$ Hz). ^{13}C NMR ($CDCl_3$) δ 181.5, 166.5, 143.7, 134.1, 131.5, 129.2, 128.7, 127.3, 126.4, 126.3, 122.9, 108.5, 73.3, 53.1, 33.0, 25.6, 22.7, 16.5, 16.4, 15.7. MS (FAB^+): m/z 365 ($M+H^+$, 57), 188 (100). HRMS (FAB^+): calcd for $C_{22}H_{28}N_3O_2$ ($M+H^+$): 366.2182, Found: 366.2177.

3-(2-Benzoylhydrazinyl)-3-tert-butyl-1-methylindolin-2-one (10Ac). A colorless crystal. mp 196-198 °C (AcOEt/hexane). IR ($CHCl_3$) 1710, 1612 cm^{-1} . 1H NMR ($CDCl_3$) δ 7.52 (1H, br d, $J=7.6$ Hz), 7.43-7.33 (3H, m), 7.32-7.23 (3H, m), 7.11 (1H, br s), 7.04 (1H, br t, $J=7.6$ Hz), 6.79 (1H, d, $J=7.6$ Hz), 5.92 (1H, br d, $J=6.4$ Hz), 3.18 (3H, s), 1.11 (9H, s). ^{13}C NMR ($CDCl_3$) δ 177.7, 166.7, 144.9, 132.9, 131.7, 129.2, 128.6, 127.5, 126.8, 126.5, 122.0, 107.7, 74.8, 37.2, 26.1, 24.8. MS (FAB^+): m/z 338 ($M+H^+$, 73), 202 (100). Anal. Calcd for $C_{20}H_{23}N_3O_2$: C, 71.19; H, 6.87; N, 12.45. Found: C, 70.94; H, 6.83; N, 12.35.

Procedure for radical addition to ketimine 3 in aqueous media. To a solution of ketimine **3** (100 mg, 0.45 mmol) in H_2O -MeOH (1:4, v/v, 4.5 mL) were added R^2I (13.6 mmol) and Et_3B (1.0 M in MeOH, 1.1 mL, 1.1 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 30 min, the reaction mixture was concentrated at reduced pressure. After the resulting residue was dissolved in AcOEt, the organic phase was washed with saturated $NaHCO_3$. The organic phase was dried over $MgSO_4$ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=2:1) afforded **4a-d**. The ethyl radical addition reaction was performed in the absence of R^2I .

Procedure for radical addition to ketimine 7 in aqueous media. To a solution of ketimine **7** (55.8 mg, 0.2 mmol) in H_2O -MeOH (1:4, v/v, 4 mL) were added R^2I (6.0 mmol) and Et_3B (1.0 M in MeOH, 0.5 mL, 0.5 mmol) under nitrogen atmosphere at 20 °C. After being stirred at the same temperature for 1 h, Et_3B (1.0 M in hexane, 0.5 mL, 0.5 mmol)

added to the reaction mixture. After being stirred at the same temperature for 2 h, the reaction mixture was diluted with saturated NaHCO₃ and then extracted with AcOEt. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=4:1, 2-fold development) afforded 10Aa-10Bd. The ethyl radical addition reaction was performed in the absence of R2I.

(R)-5,6-Dihydro-3-methyl-5-phenyl-1,4-oxazin-2-one (11). According to literature, ketimine **11** was prepared as a colorless crystal.⁴ mp 67-70 °C (AcOEt/hexane). $[\alpha]_D^{25}$ -223.2 (c 1.08, CHCl₃). IR (CHCl₃) 1740 cm⁻¹. ¹H NMR (CDCl₃) δ 7.45-7.30 (5H, m), 4.84 (1H, m), 4.56 (1H, dd, *J*=11.6, 4.5 Hz), 4.26 (1H, br t, *J*=11.6 Hz), 2.41 (3H, s). ¹³C NMR (CDCl₃) δ 160.3, 155.6, 136.8, 129.0, 128.3, 127.1, 71.5, 59.7, 21.7. MS (EI⁺): *m/z* 189 (M⁺, 10), 104 (100). Anal. Calcd for C₁₁H₁₁NO₂: C, 69.83; H, 5.86; N, 7.40. Found: C, 69.97; H, 5.92; N, 7.36.

Procedure for radical addition to ketimine 11. To a solution of ketimine **11** (50 mg, 0.265 mmol) in CH₂Cl₂ (1 mL) were added BF₃·OEt₂ (0.034 mL, 0.265 mmol), *i*-PrI (7.95 mmol), and Et₃B (1.0 M in hexane, 0.66 mL, 0.66 mmol) under oxygen atmosphere at reflux. After being stirred at the same temperature for 5 min, Et₃B (1.0 M in hexane, 0.66 mL, 0.66 mmol) added to the reaction mixture. After being stirred at the same temperature for 5 min, Et₃B (1.0 M in hexane, 0.66 mL, 0.66 mmol) added to the reaction mixture. After being stirred for 5 min, the reaction mixture was diluted with saturated NaHCO₃ and then extracted with AcOEt. The organic phase was dried over MgSO₄ and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=4:1, 2-fold development) afforded **12** and **13**. The ethyl radical addition reaction was performed in the absence of *i*-PrI. CAUTION: The addition of triethylborane under oxygen may induce fire and explosion.

(3*R*,5*R*)-3-Ethyl-3-methyl-5-phenylmorpholin-2-one (**12a**). A colorless crystal. mp 67-69 °C (AcOEt/hexane). $[\alpha]_D^{25}$ -19.7 (c 1.18, CHCl₃). IR (CHCl₃) 1735 cm⁻¹. ¹H NMR (CDCl₃) δ 7.45-7.30 (5H, m), 4.38-4.23 (3H, m), 2.08 (1H, m), 1.81 (1H, m), 1.44 (3H, s), 0.96 (3H, t, *J*=7.6 Hz). ¹³C NMR (CDCl₃) δ 173.6, 138.1, 128.9, 128.6, 127.1, 75.2, 59.7, 52.9, 30.9, 26.1, 7.3. MS (EI⁺): *m/z* 219 (M⁺, 5), 85 (100). Anal. Calcd for C₁₃H₁₇NO₂: C, 71.21; H, 7.81; N, 6.39. Found: C, 71.11; H, 7.68; N, 6.34.

(3*S*,5*R*)-3-Ethyl-3-methyl-5-phenylmorpholin-2-one (diastereomer of **12a**). A colorless crystal. mp 106-109 °C (AcOEt/hexane). $[\alpha]_D^{25}$ -28.3 (c 0.6, CHCl₃). IR (CHCl₃) 1731 cm⁻¹. ¹H NMR (CDCl₃) δ 7.44 (2H, d, *J*=7.6 Hz), 7.40-7.27 (3H, m), 4.38 (1H, dd, *J*=10.7, 3.1 Hz), 4.30 (1H, dd, *J*=10.7, 3.1 Hz), 4.22 (1H, br t, *J*=10.7 Hz), 2.06 (1H, m), 1.56 (1H, m), 1.53 (3H, s), 1.08 (3H, t, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 173.4, 138.2, 128.9, 128.6, 127.2, 75.4, 61.3, 53.2, 35.4, 26.9, 8.5. MS (FAB⁺): *m/z* 220 (M⁺+H, 95), 154 (100). Anal. Calcd for C₁₃H₁₇NO₂: C, 71.21; H, 7.81; N, 6.39. Found: C, 70.91; H, 7.73; N, 6.20.

(3*R*,5*R*)-3-Isopropyl-3-methyl-5-phenylmorpholin-2-one (**12b**). A colorless crystal. mp

71-73 °C (AcOEt/hexane). $[\alpha]_D^{27} +15.6$ (c 0.98, CHCl₃). IR (CHCl₃) 1733 cm⁻¹. ¹H NMR (CDCl₃) δ 7.44-7.31 (5H, m), 4.39 (2H, m), 4.28 (1H, br t, *J*=11.0 Hz), 2.44 (1H, m), 1.38 (3H, s), 1.01 (3H, d, *J*=7.0 Hz), 0.99 (3H, d, *J*=6.7 Hz). ¹³C NMR (CDCl₃) δ 173.5, 138.5, 128.9, 128.5, 127.1, 74.9, 62.1, 53.7, 34.4, 25.1, 17.7, 16.2. MS (FAB⁺): *m/z* 234 (M⁺+H, 100). Anal. Calcd for C₁₄H₁₉NO₂: C, 72.07; H, 8.21; N, 6.00. Found: C, 72.28; H, 8.26; N, 5.99.

(3*S*,5*R*)-3-Isopropyl-3-methyl-5-phenylmorpholin-2-one (diastereomer of 12b). A colorless crystal. mp 108-110 °C (AcOEt/hexane). $[\alpha]_D^{27} -46.8$ (c 1.2, CHCl₃). IR (CHCl₃) 1711 cm⁻¹. ¹H NMR (CDCl₃) δ 7.45 (2H, d, *J*=7.6 Hz), 7.35-7.26 (3H, m), 4.33 (1H, dd, *J*=10.5, 3.1 Hz), 4.25 (1H, dd, *J*=10.5, 3.1 Hz), 4.19 (1H, br t, *J*=10.5 Hz), 2.16 (1H, m), 1.51 (3H, s), 1.11 (3H, d, *J*=6.4 Hz), 0.97 (3H, d, *J*=6.7 Hz). ¹³C NMR (CDCl₃) δ 173.9, 138.4, 128.8, 128.6, 127.3, 74.9, 63.4, 53.2, 37.6, 25.2, 18.0, 15.4. MS (FAB⁺): *m/z* 234 (M⁺+H, 100). Anal. Calcd for C₁₄H₁₉NO₂: C, 72.07; H, 8.21; N, 6.00. Found: C, 72.04; H, 8.00; N, 6.03.

3-Ethyl-3-methyl-5-phenyl-3*H*-1,4-oxazin-2(6*H*)-one (13a). A colorless oil. IR (CHCl₃) 1742 cm⁻¹. ¹H NMR (CDCl₃) δ 7.75 (2H, d, *J*=7.0 Hz), 7.53-7.40 (3H, m), 5.42 (2H, s), 2.14 (1H, m), 1.89 (1H, m), 1.57 (3H, s), 0.92 (3H, t, *J*=7.3 Hz). ¹³C NMR (CDCl₃) δ 171.4, 159.4, 134.7, 131.3, 128.9, 126.0, 68.2, 61.3, 34.2, 26.0, 9.0. HRMS (EI⁺): *m/z* 217 (M⁺, 8), 104 (100). HRMS (EI⁺): calcd for C₁₃H₁₅NO₂ (M⁺): 217.1103, Found: 217.1097.

(*R*)-3-Isopropyl-3-methyl-5-phenyl-3*H*-1,4-oxazin-2(6*H*)-one (13b). A colorless oil. IR (CHCl₃) 1742 cm⁻¹. ¹H NMR (CDCl₃) δ 7.76 (2H, d, *J*=7.3 Hz), 7.53-7.42 (3H, m), 5.42 (1H, d, *J*=16.5 Hz), 5.37 (1H, d, *J*=16.5 Hz), 2.35 (1H, m), 1.54 (3H, s), 1.10 (3H, d, *J*=6.7 Hz), 0.93 (3H, d, *J*=6.7 Hz). ¹³C NMR (CDCl₃) δ 171.5, 158.6, 134.8, 131.3, 128.9, 126.0, 68.2, 63.2, 37.3, 24.0, 18.2, 16.1. MS (FAB⁺): *m/z* 232 (M⁺+H, 100). HRMS (FAB⁺): calcd for C₁₄H₁₈NO₂ (M⁺+H): 232.1338, Found: 232.1342.

References

- 1) Palla, G.; Predieri, G.; Domiano, P.; Vignali, C.; Turner, W. *Tetrahedron* **1986**, *42*, 3649.
- 2) Coda, A. C.; Desimoni, G.; Invernizzi, G.; Quadrelli, P.; Righetti, P. P.; Tacconi, G. *Tetrahedron* **1987**, *43*, 2843.
- 3) Somogyi, L. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 873.
- 4) Harwood, L. M.; Vines, K. J.; Drew, M. G. B. *Synlett* **1996**, 1051.

4a

4a

7