Umpolung of Michael Acceptors Catalyzed by *N*-Heterocyclic Carbenes

Christian Fischer, Sean W. Smith, David A. Powell, and Gregory C. Fu*

Department of Chemistry, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139

SUPPORTING INFORMATION

I. General

Unless otherwise noted, all reactions were carried out in oven-dried glassware under an atmosphere of argon. THF and CH₂Cl₂ were purified by passage through an alumina column as described by Grubbs.¹ All chemicals were used without further purification, unless otherwise noted. Melting points were measured on a Hoover melting point apparatus and are uncorrected.

The carbene-catalyzed β -alkylations are moisture-sensitive. Therefore, before use, the K₃PO₄ must be dried in a vacuum oven at 150 °C for 24 hours (coarse K₃PO₄, e.g., from Strem or Alfa-Aesar, must be ground into a fine powder prior to drying). The dry K₃PO₄ is stored in a glove box.

II. Preparation of Substrates and Catalysts

These procedures have not been optimized.

Preparation of cyclization substrates (representative procedure). A 100-mL flask was charged with pyridinium chlorochromate (2.6 g, 12 mmol), evacuated, and back-filled with argon. To the orange solid was added CH₂Cl₂ (50 mL) and the bromoalcohol (10 mmol). The resulting dark-brown reaction mixture was stirred for 3 hours at r.t., and then Et₂O (50 mL) was added, leading to the precipitation of a brown solid. The mixture was allowed to stand at r.t. for 2 h, and then it was filtered through a plug of silica gel (washed with Et₂O). The solvent was removed, and the residue was transferred to a 100-mL flask under argon. THF (50 mL) and then the olefination reagent² (14 mmol; added in one portion) were added, and the resulting mixture was allowed to stir at r.t. overnight. The reaction mixture was then adsorbed onto silica gel and purified by column chromatography.

[CAS# 111710-77-1]. Prepared from 5-bromopentan-1-ol (TCI) and (carbethoxymethylene)triphenylphosphorane (Alfa-Aesar) by the representative procedure.

^1H NMR (CDCl_3 , 500 MHz) δ 6.98-6.91 (m, 1H), 5.86-5.82 (m, 1H), 4.22-4.17 (m, 2H), 3.44-3.40 (m, 2H), 2.27-2.22 (m, 2H), 1.92-1.86 (m, 2H), 1.67-1.60 (m, 2H), 1.31-1.28 (m, 3H).

[CAS# 96251-91-1]. Prepared by a literature procedure.³ Tetrahydropyran-2-ol (2.4 g, 24 mmol),⁴ (carbethoxymethylene)triphenylphosphorane (11.2 g, 32.1 mmol; Alfa-Aesar), and CH_2Cl_2 (60 mL) were added to a dry round-bottom flask equipped with a magnetic stir bar under an argon atmosphere. The light-yellow solution was stirred at r.t. for 48 hours, and then it was concentrated. The resulting solid was taken up in 7:3 hexanes: Et_2O (60 mL), and the suspension was stirred at r.t. for 30 minutes. Then, it was filtered through filter paper on a Hirsch funnel (washed with hexanes). The clear filtrate was concentrated to an oil and purified by column chromatography (70:30 hexanes: EtOAc), which furnished 3.33 g (82%) of the desired compound as a light-yellow oil. Analysis by ^1H NMR revealed that the product was an 8:1 mixture of E:Z stereoisomers.

^1H NMR (CDCl_3 , 500 MHz; major isomer) δ 6.93 (dt, J = 15.5, 7.0 Hz, 1H), 5.80 (dt, J = 15.5, 1.5 Hz, 1H), 4.15 (q, J = 7.0 Hz, 2H), 3.62 (t, J = 6.0 Hz, 2H), 2.25-2.18 (m, 2H), 1.93 (br s, 1H), 1.60-1.50 (m, 4H), 1.26 (t, J = 7.0 Hz, 3H).

[CAS# 173043-75-9]. A solution of ethyl 7-hydroxyhept-2-enoate (3.00 g, 17.4 mmol) and pyridine (2.80 mL, 34.8 mmol) in dry CH_2Cl_2 (30 mL) was cooled to 0 °C in an ice bath, and tosyl chloride (5.0 g, 26 mmol) was added over a 10-minute period. The suspension was stirred at 0 °C for 1 hour, warmed to r.t., and stirred at r.t. for 16 hours. The reaction mixture was then poured into a 250-mL separatory funnel that contained

aqueous HCl (1.0 M; 150 mL) and extracted with Et₂O (3 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (90:10 hexanes:EtOAc) furnished 3.58 g (63%) of the desired product as a clear oil.

¹H NMR (CDCl₃, 500 MHz) δ 7.80 (d, *J* = 8.2 Hz, 2H), 7.37 (d, *J* = 8.2 Hz, 2H), 6.91-6.85 (m, 1H), 5.79-5.70 (m, 1H), 4.22-4.17 (m, 2H), 4.06-4.03 (m, 2H), 2.47 (s, 3H), 2.19-2.06 (m, 2H), 1.71-1.66 (m, 2H), 1.53-1.47 (m, 2H), 1.32-1.28 (m, 3H).

[CAS# 107408-35-5]. Prepared from 5-chloropentan-1-ol (Lancaster) and (carbethoxymethylene)triphenylphosphorane (Alfa-Aesar) by the representative procedure.

¹H NMR (CDCl₃, 500 MHz) δ 6.96-6.90 (m, 1H), 5.85-5.81 (m, 1H), 4.20-4.16 (m, 2H), 3.55-3.52 (m, 2H), 2.23 (q, *J* = 7.3 Hz, 2H), 1.82-1.76 (m, 2H), 1.65-1.59 (m, 2H), 1.30-1.26 (m, 3H).

Prepared from 5-bromopentan-1-ol (TCI) and allyl *P,P*-diethylphosphonoacetate (Aldrich) by the representative procedure.

¹H NMR (CDCl₃, 500 MHz) δ 7.00-6.94 (m, 1H), 6.12-6.09 (m, 1H), 5.98-5.90 (m, 1H), 5.88-5.84 (m, 1H), 5.25-5.23 (m, 1H), 4.64-4.61 (m, 2H), 3.42-3.39 (m, 2H), 2.27-2.22 (m, 2H), 1.91-1.85 (m, 2H), 1.66-1.60 (m, 2H).

¹³C NMR (CDCl₃, 125 MHz) δ 166.1, 148.7, 132.2, 121.5, 118.1, 64.9, 33.2, 32.0, 31.2, 26.4.

IR (film) 2940, 2864, 1718, 1654, 1438, 1362, 1252, 1175, 1134, 1026, 988, 932 cm⁻¹.

HRMS (EI) calcd for C₁₀H₁₅BrO₂ (M⁺) 246.0250, found 246.0230.

Prepared from 5-bromopentan-1-ol (TCI) and *N*-methoxy-*N*-methyl-2-

(triphenylphosphoranylidene)acetamide (Fluka) by the representative procedure.

^1H NMR (CDCl_3 , 500 MHz) δ 6.98-6.94 (m, 1H), 6.43 (d, J = 6.6 Hz, 1H), 3.71 (s, 3H), 3.44-3.41 (m, 2H), 3.25 (s, 3H), 2.29 (q, J = 7.0 Hz, 2H), 1.93-1.88 (m, 2H), 1.66-1.63 (m, 2H).

^{13}C NMR (CDCl_3 , 125 MHz) δ 167.0, 146.9, 119.4, 61.9, 33.6, 32.5, 32.3, 31.7, 26.9.

IR (film) 3054, 2986, 2969, 2940, 2863, 2360, 2306, 1662, 1629, 1460, 1421, 1384, 1265, 1180, 996 cm^{-1} .

HRMS (EI) calcd for $\text{C}_9\text{H}_{16}\text{BrNO}_2$ (M^+) 249.0359, found 249.0491.

[CAS# 99765-31-8]. Prepared from 5-bromopentane-1-ol (TCI) and (cyanomethyl)triphenylphosphonium chloride (Alfa-Aesar) by the representative procedure.⁵

^1H NMR (CDCl_3 , 300 MHz) δ 6.75-6.69 (m, 1H major), 6.53-6.44 (m, 1H minor), 5.36-5.30 (m, 1H major+minor), 3.42 (q, J = 6.3 Hz, 2H major+minor), 2.47 (q, J = 6.3 Hz, 2H minor), 2.26 (q, 7.2 Hz, 2H major), 1.91-1.86 (m, 2H major+minor), 1.68-1.60 (m, 2H major+minor).

^{13}C NMR (CDCl_3 , 125 MHz) δ 155.6, 154.8, 118.0, 101.0, 100.9, 33.8, 33.6, 33.0, 32.5, 32.5, 31.6, 27.3, 26.8.

Prepared from the monotosylate of 1,5-dihydroxypentane⁶ and (cyanomethyl)triphenylphosphonium chloride (Alfa-Aesar) by the representative procedure.⁷

^1H NMR (CDCl_3 , 500 MHz) δ 7.79 (d, J = 8.2 Hz, 2H major+minor), 7.37 (d, J = 8.2 Hz, 2H major+minor), 6.67-6.61 (m, 1H major), 6.46-6.40 (m, 1H minor), 5.33-5.28 (m, 1H major+minor), 4.06-4.03 (m, 2H major+minor), 2.47 (s, 3H major+minor), 2.40 (q, J = 8.2 Hz, 2H minor), 2.23-2.18 (m, 2H major), 1.71-1.65 (m, 2H major+minor), 1.56-1.49 (m, 2H major+minor).

^{13}C NMR (CDCl_3 , 125 MHz) δ 155.4, 154.6, 145.6, 133.6, 130.6, 130.5, 128.6, 128.5, 117.9, 101.1, 101.0, 70.5, 70.4, 33.2, 33.1, 31.7, 28.8, 24.8, 24.2, 22.3.

IR (film) 3055, 2987, 2305, 1722, 1712, 1598, 1422, 1358, 1265, 1177 cm^{-1} .

LRMS (ESI) calcd for $\text{C}_{14}\text{H}_{17}\text{NNaO}_3\text{S}$ ($\text{M}+\text{Na}^+$) 302.1, found 302.1.

[CAS# 439665-65-3].⁸ ¹H NMR (CDCl₃, 500 MHz) δ 6.96-6.92 (m, 1H), 6.12-6.09 (m, 1H), 4.23-4.19 (m, 4H), 3.80 (t, J = 6.1 Hz, 2H), 3.49 (t, J = 6.1 Hz, 2H), 1.31-1.27 (m, 3H).

Prepared from 6-bromohexan-1-ol (TCI) and (carbethoxymethylene)triphenylphosphorane (Alfa-Aesar) by the representative procedure.

¹H NMR (CDCl₃, 500 MHz) δ 6.94 (dt, J = 15.6, 7.0 Hz, 1H), 5.82 (ddd, J = 15.6, 1.5, 0.9 Hz, 1H), 4.20-4.16 (m, 2H), 3.42-3.39 (m, 2H), 2.24-2.20 (m, 2H), 1.90-1.84 (m, 2H), 1.51-1.45 (m, 4H), 1.30-1.27 (m, 3H).

¹³C NMR (CDCl₃, 125 MHz) δ 166.6, 148.7, 121.6, 60.2, 33.6, 32.5, 31.9, 27.6, 27.1, 14.2.

IR (film) 2981, 2936, 2860, 1717, 1655, 1463, 1445, 1367, 1308, 1268, 1241, 1221, 1187, 1134, 1096, 1043, 981 cm⁻¹.

HRMS (EI) calcd for C₁₀H₁₇BrO₂ (M⁺) 248.0406, found 248.0411.

[CAS# 71032-10-5]. Prepared from 4-bromobutan-1-ol (TCI) and (carbethoxymethylene)triphenylphosphorane (Alfa-Aesar) by the representative procedure.

¹H NMR (CDCl₃, 300 MHz) δ 6.98-6.91 (m, 1H), 5.91-5.85 (m, 1H), 4.20 (q, J = 7.1 Hz, 2H), 3.42 (t, J = 6.3 Hz, 2H), 2.40-2.36 (m, 2H), 2.06-2.00 (m, 2H), 1.29 (t, J = 7.1 Hz, 3H).

Catalyst 1. A 500-mL flask was charged with *N*-(4-methoxyphenyl)benzamide (24.9 g, 109 mmol; TCI) and PCl₅ (22.9 g, 0.110 mol), evacuated, and back-filled with argon. The flask was fitted with a nitrogen-filled balloon, and the mixture was heated to 80 °C; gas evolution was observed. The resulting yellow-green solution was stirred for 90 minutes at 80 °C. The reaction mixture was then cooled to r.t., and the volatiles were removed under reduced pressure, leaving a green solid that was used in the next step without purification (¹H NMR analysis indicated complete conversion to the imidoyl

chloride).

To the green solid was added THF (150 mL) and NEt_3 (42 mL, 0.30 mol). To this suspension was added 1-(4-methoxyphenyl)hydrazine hydrochloride (19.3 g, 0.110 mol; TCI) as a slurry in THF (100 mL) via a solid addition funnel over 30 minutes. The resulting yellow suspension was stirred at r.t. for 18 h, and then the volatiles were removed on a rotary evaporator. Acetic acid (2% aqueous solution; 200 mL) was added, and the reaction mixture was stirred vigorously for 15 minutes. The mixture was extracted with CH_2Cl_2 (5 x 100 mL), and the combined extracts were washed with brine, dried over Na_2SO_4 , and evaporated to a red oil/foam. This material (**A**) was used in the next step without further purification.

Acetic anhydride (150 mL) and formic acid (75 mL) were mixed in a 500-mL two-necked flask under argon. The solution was heated at 60 °C for 20 minutes, then cooled to r.t. and transferred to **A** via cannula under argon. The red solution was stirred for 36 hours at r.t. The volatiles were then removed on a rotary evaporator, followed by high vacuum. To the crude material was added perchloric acid (35% aqueous solution; 150 mL) and CH_2Cl_2 (150 mL). The mixture was stirred vigorously for 30 minutes, and then the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (3 x 100 mL), and the combined organic layers were washed with brine, dried over Na_2SO_4 , and concentrated to a red oil, which was dried under high vacuum for 2 days. Purification of the triazolium salt was achieved by slow crystallization from acetone/ Et_2O at -20 °C.⁹ This salt was further purified by recrystallization¹⁰ from CH_2Cl_2 to give the desired compound (4.81 g; 10% from *N*-(4-methoxyphenyl)benzamide) as a colorless fluffy solid.

mp: 220 °C.

^1H NMR (DMSO-d_6 , 500 MHz) δ 11.16 (s, 1H), 8.00 (d, J = 8.8 Hz, 2H), 7.64-7.54 (m, 7H), 7.31 (d, J = 8.9 Hz, 2H), 7.20 (d, J = 8.8 Hz, 2H), 3.89 (s, 3H), 3.84 (s, 3H).

^{13}C NMR (DMSO-d_6 , 125 MHz) δ 162.2, 161.9, 154.3, 133.4, 130.4, 130.4, 129.2, 129.1, 125.7, 123.7, 123.5, 116.5, 116.3, 57.0, 56.9.

IR (film) 3062, 2840, 1608, 1564, 1508, 1488, 1451, 1307, 1261, 1175, 1025, 834, 733, 694, 623 cm^{-1} .

HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{20}\text{N}_3\text{O}_2$ (M^+) 358.1550, found 358.1557; HRMS (ESI) calcd for ClO_4 (M^-) 98.9480, found 98.9483.

Catalyst 2. A 500-mL two-necked flask fitted with a reflux condenser was charged with *p*-methoxybenzoyl chloride (13.8 mL, 0.100 mol; TCI), evacuated, and back-filled with argon. Toluene (120 mL) and *p*-anisidine (12.3 g, 0.100 mol; Alfa-Aesar) were added. The amine slowly dissolved, and a precipitate formed. The reaction mixture was heated to 120 °C overnight. The suspension was then cooled to r.t. and filtered through a glass frit. The solid was washed with hexanes and dried under high vacuum

overnight. The amide (25.5 g) was obtained as a colorless, slightly violet solid that was used without further purification.

A 100-mL flask was charged with 4-methoxy-*N*-(4-methoxyphenyl)benzamide (5.15 g, 20.0 mmol) and PCl_5 (4.16 g, 20.0 mol), evacuated, and back-filled with argon. The flask was fitted with a nitrogen-filled balloon, and the mixture was heated to 80 °C; gas evolution was observed. The resulting yellow-green solution was stirred for 2 hours at 80 °C. The reaction mixture was then cooled to r.t., and the volatiles were removed under reduced pressure, leading to a green solid that was used in the next step without further purification (^1H NMR analysis indicated complete conversion to the imidoyl chloride).

To the green solid was added THF (60 mL), 1-(4-methoxyphenyl)hydrazine hydrochloride (3.50 g, 20.0 mol; TCI), and NEt_3 (6.3 mL, 45 mmol). [CAUTION: While we have not observed an exotherm when conducting the reaction on this scale, on larger scale the hydrazine addition has led to significant warming of the reaction mixture. For large-scale reactions, we advise that the hydrazine be added as a slurry in THF (see preparation of 1).] The yellow suspension was stirred at r.t. for 20 hours, and then the volatiles were removed on a rotary evaporator. Acetic acid (2% aqueous solution; 50 mL) was added, and the mixture was vigorously stirred for 15 minutes. The mixture was then extracted with CH_2Cl_2 (3 x 100 mL), and the combined organic extracts were washed with brine, dried over Na_2SO_4 , and concentrated, furnishing a red oil/foam. This material was used in the next step without further purification.

Acetic anhydride (50 mL) and formic acid (25 mL) were mixed in a 250-mL two-necked flask under argon. The mixture was heated to 60 °C for 20 minutes, and then it was cooled to r.t. and transferred to the amidine via cannula under argon. The red solution was stirred for 20 hours at r.t. The volatiles were then removed on a rotary evaporator and under high vacuum. To the residue was added perchloric acid (35% aqueous solution; 50 mL) and CH_2Cl_2 (50 mL). The mixture was stirred vigorously for 30 minutes, and then the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (3 x 50 mL), and the combined organic layers washed with brine, dried over Na_2SO_4 , and evaporated, yielding a red oil. The triazolium salt was crystallized from CH_2Cl_2 /hexanes at -20 °C. This impure triazolium salt was purified by recrystallization¹⁰ from CH_2Cl_2 , which furnished the target compound (1.07 g, 11% from *p*-methoxybenzoyl chloride) as a fluffy beige solid.

mp: 112 °C.

^1H NMR ($\text{DMSO}-d_6$, 500 MHz) δ 11.08 (s, 1H), 7.98 (d, J = 9.2 Hz, 2H), 7.61 (d, J = 8.9 Hz, 2H), 7.46 (d, J = 8.8 Hz, 2H), 7.29 (d, J = 9.2 Hz, 2H), 7.21 (d, J = 8.9 Hz, 2H), 7.08 (d, J = 8.8 Hz, 2H), 3.88 (s, 3H), 3.85 (s, 3H), 3.81 (s, 3H).

^{13}C NMR ($\text{DMSO}-d_6$, 125 MHz) δ 162.0, 161.0, 160.7, 153.1, 130.8, 128.1, 128.0, 124.7, 122.3, 115.3, 115.2, 114.7, 114.4, 55.8, 55.8, 55.6.

IR (film) 3073, 2841, 1611, 1508, 1498, 1307, 1259, 1176, 1027, 835, 623 cm^{-1} .

HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{22}\text{N}_3\text{O}_3$ (M^+) 388.1656, found 388.1653; HRMS (ESI) calcd for ClO_4 (M^-) 98.9480, found 98.9484.

III. Carbene-Catalyzed β -Alkylations

General Procedure for Table 2. In a glove box, a 20-mL vial was charged with catalyst **1** (46 mg, 0.10 mmol), dry K_3PO_4 (0.53 g, 2.5 mmol; run 1: Strem; run 2: Lancaster), glyme (8 mL; Fluka, anhydrous), and the substrate (1.00 mmol). The vial was closed with a teflon cap, wrapped at the cap with electrical tape, removed from the glove box, and submerged in a pre-heated oil bath at 80 °C. After the indicated time, the vial was removed from the oil bath, and the reaction mixture was directly purified by column chromatography (pentane/ Et_2O).

For entries 4, 6, 7, and 8 of Table 2, a small amount of isomerization to the β,γ -unsaturated product was observed if a long reaction time was employed.

Table 2, entry 1 [CAS# 1903-22-6]. The compound was prepared according to the General Procedure. Reaction time: 8 hours. After chromatography on silica gel (pentane: Et_2O 20:1), the desired compound was isolated as a colorless oil: run 1: 146 mg (95%); run 2: 143 mg (93%).

1H NMR ($CDCl_3$, 500 MHz) δ 5.80-5.79 (m, 1H), 4.17-4.12 (m, 2H), 2.78-2.75 (m, 2H), 2.45-2.42 (m, 2H), 1.77-1.72 (m, 2H), 1.69-1.63 (m, 2H), 1.29-1.26 (m, 3H).

Table 2, entry 2 [CAS# 1903-22-6]. The compound was prepared according to the General Procedure. Reaction time: 16 hours. After chromatography on silica gel (pentane: Et_2O 20:1), the desired compound was isolated as a colorless oil: run 1: 144 mg (94%); run 2: 143 mg (93%).

1H NMR ($CDCl_3$, 500 MHz) δ 5.80-5.79 (m, 1H), 4.17-4.12 (m, 2H), 2.78-2.75 (m, 2H), 2.45-2.42 (m, 2H), 1.77-1.72 (m, 2H), 1.69-1.63 (m, 2H), 1.29-1.26 (m, 3H).

Table 2, entry 3 [CAS# 1903-22-6]. The compound was prepared according to the General Procedure. Reaction time: 8 hours. After chromatography on silica gel (pentane: Et_2O 20:1), the desired compound was isolated as a colorless oil: run 1: 136 mg (88%); run 2: 139 mg (90%).

^1H NMR (CDCl_3 , 500 MHz) δ 5.80-5.79 (m, 1H), 4.17-4.12 (m, 2H), 2.78-2.75 (m, 2H), 2.45-2.42 (m, 2H), 1.77-1.72 (m, 2H), 1.69-1.63 (m, 2H), 1.29-1.26 (m, 3H).

Table 2, entry 4. The compound was prepared according to the General Procedure. Reaction time: 8 hours. After chromatography on silica gel (pentane: Et_2O 20:1), the desired compound was isolated as a colorless oil: run 1: 148 mg (89%); run 2: 144 mg (87%).

^1H NMR (CDCl_3 , 500 MHz) δ 5.97-5.90 (m, 1H), 5.83 (d, J = 1.8 Hz, 1H), 5.32 (dd, J = 18.6, 17.1 Hz, 1H), 5.21 (d, J = 10.4 Hz, 1H), 4.60-4.59 (m, 2H), 2.79-2.76 (m, 2H), 2.54-2.43 (m, 2H), 1.77-1.61 (m, 4H).

^{13}C NMR (CDCl_3 , 75 MHz) δ 169.9, 166.5, 132.7, 117.7, 111.3, 64.3, 36.0, 32.7, 26.4, 25.5.

IR (film) 2960, 1872, 1717, 1653, 1455, 1419, 1371, 1352, 1303, 1272, 1231, 1194, 1153, 1119, 1027, 993, 929, 857 cm^{-1} .

HRMS (EI) calcd for $\text{C}_{10}\text{H}_{13}\text{O}_2$ ($\text{M}-\text{H}^+$) 165.0916, found 165.0909.

Table 2, entry 5. The compound was prepared according to the General Procedure. Reaction time: 16 hours. After chromatography on silica gel (pentane: Et_2O 1:1), the desired compound was isolated as a colorless oil: run 1: 115 mg (68%); run 2: 114 mg (67%).

^1H NMR (CDCl_3 , 500 MHz) δ 6.33-6.31 (m, 1H), 3.70 (s, 3H), 3.20 (s, 3H), 2.84-2.81 (m, 2H), 2.49-2.46 (m, 2H), 1.76-1.73 (m, 2H), 1.67-1.64 (m, 2H).

^{13}C NMR (CDCl_3 , 125 MHz) δ 168.9, 167.9, 109.6, 62.1, 37.0, 33.1, 33.0, 27.3, 26.1.

IR (film) 3583, 3503, 3257, 3063, 2958, 2870, 2823, 1659, 1631, 1462, 1407, 1381, 1324, 1178, 1091 cm^{-1} .

HRMS (EI) calcd for $\text{C}_9\text{H}_{15}\text{NO}_2$ (M^+) 169.1097, found 169.1100.

Table 2, entry 6 [CAS# 5732-88-7]. The compound was prepared according to the

General Procedure. Reaction time: 8 hours. Upon cooling, the reaction mixture was diluted with CH₂Cl₂ (30 mL), transferred to a 150-mL separatory funnel, and washed with distilled water (3 x 20 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄, passed through a glass frit, and concentrated via a rotary evaporator. The resulting dark-red oil was purified via kugelrohr distillation at 120 °C (10 torr; if the product was purified by flash chromatography, ~20% of the β,γ-unsaturated isomer was observed). The desired compound (20:1 mixture of the α,β:β,γ isomers) was isolated as a colorless oil: run 1: 78 mg (72%); run 2: 75 mg (70%).

¹H NMR (CDCl₃, 300 MHz) δ 5.25-5.23 (m, 1H), 2.62-2.58 (m, 2H), 2.48-2.43 (m, 2H), 1.82-1.77 (m, 4H).

Table 2, entry 7 [CAS# 5732-88-7]. The compound was prepared according to the General Procedure. Reaction time: 8 hours. Upon cooling, the reaction mixture was diluted with CH₂Cl₂ (30 mL), transferred to a 150-mL separatory funnel, and washed with distilled water (3 x 20 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄, passed through a glass frit, and concentrated via a rotary evaporator. The resulting dark-red oil was purified via kugelrohr distillation at 120 °C (10 torr; if the product was purified by flash chromatography, ~20% of the β,γ-unsaturated isomer was observed). The desired compound (20:1 mixture of the α,β:β,γ isomers) was isolated as a colorless oil: run 1: 92 mg (86%); run 2: 88 mg (82%).

¹H NMR (CDCl₃, 300 MHz) δ 5.25-5.23 (m, 1H), 2.62-2.58 (m, 2H), 2.48-2.43 (m, 2H), 1.82-1.77 (m, 4H).

Table 2, entry 8. The compound was prepared according to the General Procedure. Reaction time: 6 hours. After chromatography on silica gel (pentane:Et₂O 6:1 → 3:1), the desired compound (2:1 mixture of the illustrated olefin isomers, along with ~5% of an unidentified isomer) was isolated as a colorless oil: run 1: 117 mg (75%); run 2: 124 mg (79%).

¹H NMR (CDCl₃, 500 MHz) δ 5.86-5.85 (m, 1H major), 5.80-5.79 (m, 1H minor), 4.72 (m, 2H major), 4.39 (m, 2H minor), 4.20-4.14 (m, 2H major+minor), 3.98 (d, *J* = 7.0 Hz, 2H minor), 3.90-3.87 (m, 2H major), 3.05-3.02 (m, 2H minor), 2.74-2.71 (m, 2H major), 1.31-1.24 (m, 3H major+minor).

¹³C NMR (CDCl₃, 125 MHz) δ 166.3, 166.1, 162.7, 161.7, 111.4, 110.2, 72.0, 71.5, 68.9, 67.0, 60.0, 59.9, 34.4, 32.6, 17.9, 14.3.

IR (film) 2981, 2860, 1710, 1667, 1373, 1351, 1215, 1124, 1076, 1035, 921, 851 cm^{-1} .
HRMS (EI) calcd for $\text{C}_8\text{H}_{12}\text{O}_3$ (M^+) 156.0781, found 156.0784.

Table 2, entry 9 [CAS# 1552-91-7]. The compound was prepared according to the General Procedure. Reaction time: 16 hours. After chromatography on silica gel (pentane:Et₂O 20:1), the desired compound was isolated as a colorless oil: run 1: 135 mg (80%); run 2: 136 mg (81%).

¹H NMR (CDCl_3 , 500 MHz) δ 5.60 (s, 1H), 4.16-4.11 (m, 2H), 2.83-2.81 (m, 2H), 2.20-2.18 (m, 2H), 1.67-1.57 (m, 6H), 1.29-1.26 (m, 3H).

Table 2, entry 10 [CAS# 27741-65-7]. The compound was prepared according to the General Procedure, except that 20% of catalyst **1** was used, and dioxane was employed as the solvent (a slightly lower yield was obtained when glyme was used). Reaction time: 33 hours. After chromatography on silica gel (pentane:Et₂O 20:1), the desired compound was isolated as a colorless oil: run 1: 66 mg (47%); run 2: 70 mg (49%).

¹H NMR (CDCl_3 , 300 MHz) δ 5.59-5.57 (m, 1H), 4.14 (q, J = 7.2 Hz, 2H), 3.15-3.12 (m, 2H), 2.85-2.83 (m, 2H), 2.09 (quintet, J = 7.9 Hz, 2H), 1.27 (t, J = 7.2 Hz, 3H).

References

- (1) Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518-1520.
- (2) Generally, a pre-formed ylide was used. If the olefination agent was generated in situ, then the aldehyde was added to the olefination agent at $-78\text{ }^{\circ}\text{C}$, and the mixture was then allowed to warm to r.t.
- (3) Pandey, G.; Hajra, S.; Ghorai, M. K.; Kumar, K. R. *J. Org. Chem.* **1997**, *62*, 5966-5973.
- (4) Prepared according to: Schniepp, L. E.; Geller, H. H. *J. Am. Chem. Soc.* **1946**, *68*, 1646-1648.
- (5) The desired product was isolated and used as a 2:1 E:Z mixture.
- (6) Prepared according to: Bouzide, A.; Sauv, G. *Org. Lett.* **2002**, *4*, 2329-2332.
- (7) The desired product was isolated and used as a 1.7:1 E:Z mixture.
- (8) Prepared according to: Lautens, M.; Paquin, J.-F.; Piguel, S. *J. Org. Chem.* **2002**, *67*, 3972-3974.
- (9) The initial crystallization, which requires dilute conditions, can be slow (may take up to 5 days).
- (10) CH_2Cl_2 was added at reflux until almost all of the solid had dissolved. After cooling to r.t., the mixture was cooled to $-20\text{ }^{\circ}\text{C}$ for 12 h. The crystallized solid was collected on a glass frit and washed with a small amount of cold CH_2Cl_2 and Et_2O .

STANDARD PROTON PARAMETERS

expt1 s2pu1

SAMPLE		DEC. & VI	
date	Aug 19 2005	dftq	125.674
solvent	CDCl3	dn	C13
file	/data/export/~	dpwr	34
home/gfu/tsm/bu11w	dof		1498.1
inle/tsm-02-042-bv	dm		nnn
romester.fid	dmm		w
ACQUISITION	dmf		10000
sfrq	499.749	dseq	
tn	H1	dres	1.0
at	3.277	homo	
np	65536	PROCESSING	
sw	9998.8	wtfile	
fb	not used	proc	ft
bs	16	fn	65536
tpwr	56	math	f
pw	8.2	wert	
dl	0	wexp	
tof	1498.1	wbs	
nt	16	wnt	
ct	16		
atlock	n		
gain	not used		
FLAGS			
il	n		
in	n		
dp	y		
hs	nm		
DISPLAY			
sp	5.7		
wd	3993.9		
vs	224		
sc	0		
wc	250		
hzmm	15.98		
is	33.57		
rfl	4639.0		
rffp	3633.1		
th	7		
ins	1.000		
nm	cdc	ph

STANDARD 1H OBSERVE

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 File: DP-V-E627_H
 INOVA-500 "zippy"
 PULSE SEQUENCE
 Relax: delay 1.000 sec
 Pulse: 34.1 degrees
 Acq. time 1.935 sec
 Width 4506.5 Hz
 16 repetitions
 OBSERVE H1, 300.0986336 MHz
 DATA PROCESSING
 FI size 32768
 Total time 0 min, 48 sec

STANDARD PROTON PARAMETERS

exp4 s2pu1

SAMPLE

date Oct 20 2005 DEC. & VT
 solvent CDC13 dn 125.674
 file /data/gfu/Tsm~ C13
 /Tsm-02-030-dty.fti~ 34
 1498.1
 mm

ACQUISITION

sfrq 499.749 dmf 10000
 tn H1 dseq
 at 3.277 dres 1.0
 np 65536 homo n
 sw 9998.8
 fb not used dffq2 0
 bs 16 dn2
 tpwr 56 dpwr2 1
 pw 8.2 dof2 0
 d1 0 dm2 n
 tof 1498.1 dmm2 c
 nt 16 dmf2 200
 ct 0 dseq2
 alock n dres2 1.0
 gain not used homo2 n

FLAGS

il n dffq3 0
 in n dn3
 dp y dpwr3 1
 hs mn dof3 0
 dm3
 DISPLAY -0.2 dmm3 c
 sp 3992.5 dm3 200
 wp 224 dseq3
 vs 0 dres3 1.0
 sc 250 homo3 n

PROCESSING

hzmm 15.97 wfile
 is 33.57 proc ft
 rfl 4638.2 fn 65536
 rfp 3637.1 math f
 th 7
 lns 1.000
 rim cdc ph
 weft
 wexp
 wbs
 wnt

STANDARD PROTON PARAMETERS

exp1 s2pu1

SAMPLE

DEC. & VT

date Oct 6 2005 dfreq 125.674

solvent CDC13 dn C13

file /data/export/~ dpwr 34

home/gfu/1cf/bu1lw~ dof 1498.1

inlie/sring_chlori~ dm nm

de.fid dm w

ACQUISITION sfreq 499.749 dmf 10000

tn H1 dres 1.0

at 3.277 homo n

mp 65536 wtfile n

sw 9998.8 proc ft

fb not used f 65536

bs 8 fn f

tpwr 56 math

pw 8.2 warr

dl 3.000 wexp

tof 1498.1 wbs

nt 16 wnt

ct 16

clock n

gain not used

FLAGS

il n

in n

dp y

hs nm

DISPLAY

sp -4.1

wp 4007.1

vs 224

sc 0

wc 250

hzmm 16.03

is 2031.20

rf1 1002.6

rfp 0

th 7

ins 2.000

nm cdc ph 2.000

STANDARD PROTON PARAMETERS

exp2 szpu1

SAMPLE

date Jul 11 2005 DEC. 4 VT

solvent /data/export/~dpwr C13

file /data/export/~dpwr 34

home/gfu/tcf/bu1lw~dm 1498.1

lnkle/CF6-29fr71-8~dm 10000

1.fid dm

ACQUISITION

sfrq 499.749 dmf 10000

ln H1 dseq 1.0

at 3.277 H1 homo 1.0

np 65536 dres

sw 9998.8 wfile

td not used ft

bs 8 math 65536

ipwr 56

pw 8.2

dl 3.000 werr

tof 1498.1 wexp

nt 16 wbs

ct 16 wnt

clock n

gain not used

flags

il n

in n

dp y

hs nm

DISPLAY

sp -510.9

wp 4506.3

vs 89

sc 0

wc 250

hzmm 18.03

is 511.45

rfl 4638.2

rfp 3628.1

th 2.000

lns nm

cdc ph

STANDARD PROTON PARAMETERS

expi s2pul

SAMPLE DEC. 8 VT

date Aug 17 2005 dfreq 125.674

solvent CDC13 dn C13

file /data/export/~ dpwr 34

home/gfu/Tsm/bu1lw~ dof 1498.1

inkle/Tsm-02-041we~ dm nm

inreb.fid dmf w

ACQUISITION 499.749 dseq 10000

tn H1 dres 1.0

at 3.277 homo n

np 65536 wfile PROCESSING

sw 9998.8 wfile ft

fb not used proc 65536 f

bs 16 fn math

tpwr 56 math

pw 8.2 werr

dl 0 wexp

tof 1498.1 wbs

nt 16 wnt

ct 16

atlock n

gain not used

FLAGS

i1 n

in n

dp y

hs nm

DISPLAY

sp -5.3

wp 4007.4

vs 124

sc 0

wc 250

hzmm 16.03

is 201.44

rfi 4636.9

rfd 3633.1

th 7

ins 1.000

nm cdc ph

jw08-239

expt std1h

SAMPLE
date Nov 3 2005
solvent CDCl3
file /data/gfu/tsm/
/Tsm-02-086dry.fid
ACQUISITION
sfrq 300.100
tn H1
at 1.995
np 17964
sw 4506.5
fb not used
bs 2
tpwr 54
pw 7.0
dl 1.000
tof 0
nt 64
ct 64
alock n
gain not used
flags
il n
in n
dp y
DISPLAY
sp -0.0
wp 2397.9
vs 162
sc 0
wc 250
h2mm 9.59
is 97.78
rfl 2898.3
rflp 2181.7
th 20
ins 1.000
nm

DEC. & VT
300.100
H1
dn
dpwr
dof
dm
dm
c
200
PROCESSING
wtfile
proc
fn
ft
not used
werf
wexp
wbs
wnt

STANDARD PROTON PARAMETERS

expl szpu1

SAMPLE DEC. & VT
 date Sep 6 2005 dfrq 125.674
 solvent CDCl3 dn C13
 file /data/export/~ dpvr 34
 home/gfu/tsm/bu1lw~ dof 1498.1
 inkle/tsm-01-033-cv dm nm
 hecksept05.fid dmf w
 ACQUISITION 10000
 sfrq 499.749 dres
 tn H1
 at 3.277 homo PROCESSING 1.0
 np 65536
 sw 9998.8 wifile
 fb not used ft
 bs 16 fn 65536
 tpwr 56 math f
 pw 8.2 weff
 d1 0 weff
 tof 1498.1 wexp
 nt 16 wds
 ct 16 wnt
 alock n
 gain not used
 flags n
 i1 n
 in y
 dp y
 hs nm
 DISPLAY
 sp -5.6
 wp 4007.1
 vs 123
 sc 0
 wc 250
 h2mm 16.03
 is 33.57
 rfi 4637.5
 rfp 3633.1
 th 7
 ins 1.000
 nm cdc ph

STANDARD PROTON PARAMETERS

exp1 szpu1

SAMPLE

date

Sep 30 2005

DEC. & VT

125.674

C13

34

nm

1498.1

nm

10000

1.0

n

f

65536

ft

wtfile

proc

math

8

fn

65536

f

ft

wtfile

proc

math

8

fn

65536

f

date Sep 30 2005
 solvent CDC13
 file /data/export/~
 home/gtu/tct/pu1w~
 lnkle/Cf6-098f28~
 40.fid
 ACQUISITION
 sfrq 499.749
 tn H1
 at 3.277
 np 65536
 sw 9998.8
 fd not used
 bs 8
 tpwr 56
 pw 8.2
 dl 3.000
 tof 1498.1
 nt 16
 ct 16
 alock n
 gain not used
 flags n
 in n
 dp y
 hs nm
 DISPLAY
 sp -4.1
 wp 4007.1
 vs 112
 sc 0
 wc 250
 hzmm 16.03
 is 477.08
 rfi 1002.6
 rfp 0
 th 7
 ins 2.000
 nm cdc ph

STANDARD PROTON PARAMETERS

expl szpul

SAMPLE

DEC. & VT

date Jun 30 2005 dfrq 125.674
 solvent CDCl3 dn C13
 file /data/export/~dpwr
 home/gfu/Tcf/bu1lw~dof 1498.1
 inkle/CF6-011fr37~dm nm
 50.fid dmm w
 ACQUISITION 499.749 dmf 10000
 sfrq 499.749 dres
 tn H1
 at 3.277 homo
 np 65536
 sw 9998.8 wifile
 fb not used ft
 bs 8 fn 65536
 tpwr 56 math f
 pw 8.2 weff
 dl 3.000 weff
 tof 1498.1 weff
 nt 16 wbs
 ct 16 wnt
 gain not used
 alock n
 flags not used
 il n
 in n
 dp y
 hs nm

DISPLAY

sp -4.1
 wp 4007.1
 vs 112
 sc 0
 wc 250
 hzmm 16.03
 is 1284.19
 rfi 1002.6
 rfp 0
 th 7
 ins 3.000
 nm cdc ph

STANDARD 1H OBSERVE

expi stdh

SAMPLE

DEC. & VT

date	Oct 19 2005	dfreq	300.100
solvent	CDCl3	dn	H1
file	/data/export/~	dpwr	30
home/gfu/Tsm/mrnat/~	dof	nmn	0
/Tsm-02-086-pure.f~	dm	c	
id	dmf	200	

ACQUISITION

sfrq	300.100	wtfile	
tn	H1	proc	ft
at	1.995	fn	not used
np	17984		
sw	4506.5		
fb	not used	werf	
bs	16	wexp	
tpwr	54	wbs	
pw	7.0	wnt	
d1	1.000		
tof	0		
nt	16		
ct	16		
alock	n		
gain	not used		
flags			
il	n		
in	n		
dp	y		

DISPLAY

sp	-0.3
wp	2398.2
vs	162
sc	0
wc	250
hzmm	9.59
is	303.43
rfl	2898.6
rfd	2181.7
th	20
ins	1.000

nm cdc ph

STANDARD PROTON PARAMETERS

exp2 s2pu1

Ar = *p*-anisyl

SAMPLE DEC. & VT
 date Jul 10 2005 dfrq 125.675
 solvent DMSO dn C13
 file /data/export/~ dpvr 34
 home/gfu/tcf/bu1w~ dof 1498.1
 lnkle/PMFX2carbene~ dm nnn
 H.fid w 10000

ACQUISITION
 sfrq 499.751 dmf
 tn H1 dseq
 at 3.277 H1 homo 1.0
 np 65536 dres n
 sw 9998.8 wf file
 fb not used proc
 bs 8 fn
 tpwr 56 math 65536
 pw 8.2
 dl 3.000 werr
 tof 1498.1 wexp
 nt 16 wbs
 ct 16 wnt
 atlock n
 gain not used

FLAGS
 i1 n
 in n
 dp Y
 hs nm

DISPLAY
 sp -4.1
 wp 6004.2
 vs 112
 sc 0
 wc 250
 hzmm 24.02
 is 515.44
 rfl 1002.5
 rfp 0
 th 7
 tns 4.000
 nm cdc ph

PROCESSING
 ft
 f

STANDARD PROTON PARAMETERS

exp2 s2pu1

SAMPLE

DEC. & VT

date Jul 7 2005 dfreq 125.675
solvent DMSO dn C13
file /data/export/~ dpwr 34
home/gfu/Tcf/buliy~ dof 1498.1
inkle/CF6-015crop2~ dm mm
.fid mm w

ACQUISITION

dfreq 499.751
ln H1 dseq 10000
at H1 dres 1.0
np 3.277 homo n
sw 65536 wfile
fb 9998.8 wfile
not used proc ft
bs 8 fn 65536
tpwr 56 math f

PROCESSING

pw 8.2 werr
dl 3.000 wexp
tof 1498.1 wds
nt 16 wnt
ct 16
a-lock n
gain not used

FLAGS

il n
in n
dp Y
hs nm

DISPLAY

sp 994.3
wp 5005.8
vs 112
sc 0
wc 250
hzmm 20.02
is 2315.43
rfl 1002.5
rfl 0
th 7
ins 9.000
nm cdc ph

Ar = p-anisyl

STANDARD PROTON PARAMETERS

exp4 s2put1

SAMPLE		DEC. & VT	
date	Aug 22 2005	dfrq	125.674
solvent	CDCl3	dn	C13
file	/data/gfu/Tsm~	dpwr	34
	/Tsm-02-046.fid	dof	1498.1
ACQUISITION			
sfrq	499.749	dm	nmn
ln	H1	dmm	w
at	3.277	dseq	10000
np	65536	dres	1.0
sw	9998.8	homo	n
fb	not used	DEC2	0
bs	16	dfrq2	0
tpwr	56	dn2	1
pw	8.2	dpwr2	1
dl	0	dof2	n
tof	1498.1	dm2	n
nt	16	dmm2	n
ct	16	dmt2	200
alock	n	dseq2	1.0
gain	not used	dres2	n
FLAGS	not used	homo2	DEC3
il	n	dfrq3	0
in	n	dn3	1
dp	y	dpwr3	0
hs	nm	dof3	n
DISPLAY			
sp	-6.6	dmm3	n
wp	4007.2	dmt3	200
vs	224	dseq3	1.0
sc	0	dres3	n
wc	250	homo3	PROCESSING
hzm	16.03	wtfile	ft
is	68.08	proc	fn
rfl	4838.2	math	65536
rtp	3633.1	werr	wexp
th	7	wbs	wnt
ins	1.000		
nm	cdc		
ph

Table 2, entry 1

STANDARD 1H OBSERVE

exp3 std1h

SAMPLE DEC. & VT
date Oct 24 2005 dfrq 300.100
solvent CDCl3 dn H1
file /data/gfu/Tsm~ dpwr 30
/Tsm-02-096.fid dof 0
ACQUISITION nm
sfrq 300.100 dnm 200
tp H1 dmf
at 1.995 wtfile PROCESSING
np 17984 wfile ft
sw 4506.5 proc not used
fb not used fn
bs 16
tpwr 54 werr
pw 7.0 wexp
d1 1.000 wbs
tof 0 wnt
nt 16
ct 16
alock n
gain not used
FLAGS
il n
in n
dp y
DISPLAY
sp -1.1
wp 2400.1
vs 153
sc 0
wc 250
hznm 9.60
is 438.43
rfl 2898.6
rfp 2181.7
th 20
tms nm
nm cdc ph 2.000

Table 2, entry 2

STANDARD PROTON PARAMETERS

expt s2pul

Table 2, entry 3

SAMPLE		DEC. & VT	
date	Sep 16 2005	dfrq	125.674
solvent	CDCl3	dn	C13
file	/data/export/~	dpwr	34
home/gfu/tcf/bu11w	~	dof	1498.1
inkle/Cf6-088-2fr2~		dm	mm
5-34.fid		dm	mm
ACQUISITION		dmf	10000
sfrq	499.749	dseq	
tn	H1	dres	1.0
at	3.277	homo	
np	65536	wtfile	n
sw	9998.8	proc	
fb	not used	fn	ft
bs	not used	math	f
tpwr	56	werf	
pw	8.2	wexp	
d1	3.000	wbs	
tof	1498.1	wnt	
nt	16		
ct	8		
atock	n		
gain	not used		
FLAGS			
il	n		
in	n		
dp	y		
hs	nm		
DISPLAY			
sp	-11.8		
wd	4001.6		
vs	224		
sc	0		
wc	250		
hzmm	16.01		
is	1906.76		
rfl	4637.5		
rffp	3626.1		
th	7		
ins	4.000		
nm	cdc ph

STANDARD PROTON PARAMETERS

expl szpul

SAMPLE

date Aug 11 2005 DEC. & VT

solvent CDCl3 dn 125.674

file /data/export/~ dpwr 34

home/gfu/Tcf/bu1lw~ dof 1498.1

inle/CF6-074fr16~ dm 10000

32.fid dm 10000

ACQUISITION

sfrq 499.749 dres 1.0

tn H1 dres 1.0

at 3.277 homo

np 65536 wfile

sw 9998.8 wfile

fb not used ft

bs 8 fn 65536

tpwr 56 math

pw 8.2 weff

dl 3.000 wexp

tof 1498.1 wds

nt 16 wnt

ct 16

alock n

gain not used

flags

il n

in n

dp y

hs nm

DISPLAY

sp 488.0

wp 3507.9

vs 69

sc 0

wc 250

hzm 14.03

is 710.64

rfl 4687.6

rtp 3628.1

th 2.000

ins nm

STANDARD PROTON PARAMETERS

expi szpu1

SAMPLE		DEC. & VT	
date	Sep 9 2005	dfrq	125.674
solvent	CDCl3	dn	C13
file	/data/export/~	dpwr	34
home/gfu/Tsm/bu1lw~		dof	1498.1
lnkle/Tsm-02-054.f~		dm	nmn
id		dmm	w
dmf		df	10000
ACQUISITION		dseq	
sfrq	499.749	dres	1.0
tn	H1	homo	n
at	3.277	wtfile	
np	65536	proc	ft
sw	9998.8	fn	65536
fb	not used	math	f
bs	16	weff	
tpwr	56	wexp	
pw	8.2	wbs	
dl	0	wnt	
tof	1498.1		
nt	16		
ct	16		
alock	n		
gain	not used		
flags			
il	n		
in	y		
dp	nm		
hs			
DISPLAY			
sp	-5.9		
wd	4007.4		
vs	147		
sc	0		
wc	250		
hzmm	16.03		
is	38.57		
rfl	4637.5		
rffp	3633.1		
th	7		
ins	3.000		
nm	cdc		
ph

STANDARD 1H OBSERVE

expi std1h

SAMPLE
date Sep 24 2005
solvent CDCl3
file /data/export/~
home/gfu/Tsm/mrhat~
/Tsm-02-073-dry.f1~
ACQUISITION
sfrq 300.100
in H1
at 1.995
np 17984
sw 4506.5
fb not used
bs 16
tpr 54
pw 7.0
d1 1.000
tof 0
nt 16
ct 16
alock n
gain not used
FLAGS
f1 n
in n
dp y
DISPLAY
sp 265.9
wp 2037.1
vs 162
sc 0
wc 250
hzm 8.15
ls 198.03
rfi 2898.6
rfp 2181.7
th 20
ins 1.000
nm cdc ph

DEC. & VT
dfrq 300.100
dn H1
dpr 30
dof 0
dm nnn
dmm c
dmf 200
PROCESSING
ft
proc not used
werr
wexp
wbs
wnt

Table 2, entry 6

STANDARD 1H OBSERVE

expi std1h

SAMPLE DEC. & VT
 date Oct 7 2005 dfreq 300.100
 solvent CDC13 dn H1
 file /data/export/~ dpwr 30
 home/gfu/Tsm/mrnat/~ dof 0
 /Tsm-02-076nittrile~ dm nm
 .fid dmm c
 ACQUISITION .fid dmf 200

PROCCESSING
 tn 300.100 wtfile
 at H1 proc ft
 np 1.995 fn not used
 sw 17984
 fb 4506.5 wefr
 bs not used wexp
 tpwr 16 wbs
 pw 54 wnt
 dl 7.0
 1.000
 tof 0
 nt 16
 ct 16
 alock n
 gain not used
 FLAGS
 il n
 in n
 dp y

DISPLAY
 sp -0.0
 wp 2398.2
 vs 162
 sc 0
 wc 250
 hzmm 9.59
 is 243.70
 rf1 2898.3
 rfp 2181.7
 th 20
 ins 1.000
 nm cdc ph

Table 2, entry 7

STANDARD PROTON PARAMETERS

exp1 szpul

SAMPLE		DEC. & VT	
date	Oct 2 2005	dfreq	125.674
solvent	CDCl3	dn	C13
file	/data/export/~	dpwr	34
home/gfu/Tcf/bu1lw~		dof	1498.1
inkle/CF6-099f38~		dm	nm
49.fid		dmm	w
ACQUISITION		daf	10000
sfrq	499.749	dseg	
tn	H1	dres	1.0
at	3.277	homo	n
np	65536	wtfile	ft
sw	9398.8	proc	65536
fb	not used	fn	f
bs	8	math	
tpwr	56	werr	
pw	8.2	wexp	
dl	3.000	wbs	
tof	1498.1	wnt	
nt	16		
ct	16		
alock	n		
gain	not used		
FLAGS			
il	n		
in	n		
dp	y		
hs	nm		
DISPLAY			
sp	-4.1		
wp	4007.1		
vs	112		
sc	0		
wc	250		
hzmm	16.03		
is	520.36		
rfl	1002.6		
rfp	0		
th	7		
ins	2.000		
nm	cdc ph

STANDARD PROTON PARAMETERS

exp1 s2pu1

SAMPLE DEC. & VT
 date Oct 3 2005 dfreq 125.795
 solvent CDCl3 dn C13
 file /data/export/~ dpwr 37
 home/gfu/tcf/rocky~ dof 0
 /Cf6-100ft20-31.ft~ dm nm
 d dnm C
 10000
 ACQUISITION
 sfrq 500.235 dmf
 tn H1 dseq
 at 3.200 dres 1.0
 np 64000 homo n
 sw 10000.0 wlfite
 fb not used proc ft
 bs 8 fn 131072
 ss 1 math f
 tpwr 59 weff
 pw 9.8 weff
 dl 3.000 wexp
 tof 1498.2 wds
 nt 16 wnt
 ct 16
 clock n
 gain not used
 flags
 il n
 in n
 dp y
 hs nm
 DISPLAY
 sp -4.5
 wp 4007.7
 vs 200
 sc 0
 wc 250
 hzmm 16.03
 is 1150.85
 rfl 4834.8
 rfp 3631.7
 th 7
 ins 2.000
 nm cdc ph

STANDARD PROTON PARAMETERS

exp1 s2pul

SAMPLE

DEC. & VT

date Jun 27 2005 dfrq 125.674

solvent CDC13 dn C13

file /data/export/~ dpwr 34

home/gfu/tsm/bu1lw~ dof 1498.1

inkle/tsm-01-111-2~ dm nm

ACQUISITION .fid dmf 10000

sfrq 499.749 dseq 1.0

tn HI dres n

at 3.277 homo

np 65536 wtfile

sw 9998.8 wfile ft

fb not used PROC 65536 f

bs 16 fn

tpwr 56 math

pw 8.2 werr

di 0 wexp

tof 1498.1 wbs

nt 16 wnt

ct 16

atlock n

gain not used

FLAGS

il n

in n

dp y

hs nm

DISPLAY

sp 6.3

wd 3993.9

vs 224

sc 0

wc 250

hzmm 15.98

is 33.57

rfl 4638.4

rfl 3633.1

th 7

ins 2.000

nm cdc ph