Supplementary Content

A unified approach to the asymmetric synthesis of higher homologues of (3-quinolyl)-alanine

Shital K Chattopadhyay* and Indranil Kundu

*Department of Chemistry, University of Kalyani,
Kalyani-741235,
WestBengal, India*

E-mail: skchatto@yahoo.com

CONTENTS

1. Experimental details of the compounds not included
in experimental section	pages 2-9
2. ^1H and ^{13}C -NMR spectra of all compounds	pages 10-42

Synthesis of the olefins **1b-c**:

(*S*)-tert-butyl 4-allyl-2,2-dimethyloxazolidine-3-carboxylate (**1b**)

n-BuLi (2 M in hexane, 1.7 mL, 3.45 mmol, 1.5 eqv.) was added to a stirred suspension of methyltriphenylphosphonium bromide (1.23 g, 3.45 mmol, 1.5 eqv.) in dry THF (10 mL) under argon at 0 °C dropwise over 5 min, and the resulting solution was allowed to stir for 20 min at 0 °C. A solution of the aldehyde **4** (560 mg, 2.30 mmol) in THF (5 mL) was then added dropwise over 5 min with stirring at the same temperature. After 20 min, the solution was allowed to come to room temperature and stirred for another 4 h. The reaction mixture was quenched by addition of aq NH₄Cl solution (3 mL) and then extracted with ethyl acetate (2 × 50 mL). The combined organic extract was washed successively with H₂O (40 mL) and brine solution (40 mL) and then dried over MgSO₄. It was then filtered, and the filtrate was concentrated under reduced pressure to leave the crude product which was purified by column chromatography over silica gel (EtOAc–PE, 1:19) to give the olefin **1b** as a colourless liquid (410 mg, 74 %) . $[\alpha]_D^{25} +17.1$ (c 0.41, CHCl₃). IR (CHCl₃): 2927, 1695, 1591, 1484, 1439, 1391, 1191 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 5.74 (t, *J* = 6.4 Hz, 1H), 5.09 (d, *J* = 17.2 Hz, 2H), 3.97-3.77 (m, 3H), 2.58-2.43 (m, 1H), 2.26-2.24 (m, 1H), 1.59 (1.55) (s, 3H), 1.47 (merged singlets, 12H). ¹³C NMR (100MHz, CDCl₃): δ 152.1 (151.7), 134.6, 117.7, 93.8 (93.3), 80.0 (79.5), 66.4 (66.0), 56.9 (56.8), 38.1 (37.1), 28.5, 27.5 (26.7), 24.5 (23.2). Anal.Calcd for C₁₃H₂₃NO₃: C, 64.70; H, 9.61; N, 5.80; Found: C, 64.91; H, 9.78; N, 5.68. MS (TOF, ES⁺): *m/z* (%) 264 (100) [M+Na].

(*S*)-tert-butyl 4-(but-3-enyl)-2,2-dimethyloxazolidine-3-carboxylate (**1c**)

n-BuLi (2 M in hexane, 1.7 mL, 3.45 mmol, 1.5 eqv.) was added to a stirred suspension of methyltriphenylphosphonium bromide (1.23 g, 3.45 mmol, 1.5 eqv.) in dry THF (10 mL)

under argon at 0 °C dropwise over 5 min, and the resulting solution was allowed to stir for 20 min at 0 °C. A solution of the aldehyde **5** (590 mg, 2.30 mmol) in THF (5 mL) was then added dropwise over 5 min with stirring at the same temperature. After 20 min, the solution was allowed to come to room temperature and stirred for another 4 h. The reaction mixture was quenched by addition of aq NH₄Cl solution (3 mL) and then extracted with ethyl acetate (2 × 50 mL). The combined organic extract was washed successively with H₂O (40 mL) and brine solution (40 mL) and then dried over MgSO₄. It was then filtered, and the filtrate was concentrated under reduced pressure to leave the crude product which was purified by column chromatography over silica gel (EtOAc–PE, 1:19) to give the olefin **1b** as a colourless liquid (419 mg, 76 %) . $[\alpha]_D + 15.4$ (c 0.34, CHCl₃). IR (CHCl₃): 2979, 2933, 2872, 1698, 1641, 1388, 1365, 1259 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 5.90-5.70 (m, 1H), 5.03 (dd, $J = 17.2, 1.6$ Hz, 1H), 4.97 (d, $J = 8.4$ Hz, 1H), 3.94-3.74 (m, 3H), 2.05-1.80 (m, 3H), 1.74-1.47 (m, 16H). ¹³C NMR (100MHz, CDCl₃): δ 152.7, 137.9, 114.9, 93.7, 80.0, 66.8 (66.6), 57.3 (56.8), 32.7 (31.9), 30.6, 28.5, 27.6 (26.8), 24.6. Anal.Calcd for C₁₄H₂₅NO₃: C, 65.85; H, 9.87; N, 5.49; Found: C, 65.77; H, 9.75; N, 5.58. MS (TOF, ES⁺): m/z (%) 278 (100) [M+Na].

Synthesis of the coupled products **11** & **12**:

(*S,E*)-tert-butyl 2,2-dimethyl-4-(3-(quinolin-3-yl)allyl)oxazolidine-3-carboxylate (**11**)

A solution of the olefin **1b** (180 mg, 0.75 mmol), 3-bromoquinoline (171 mg, 0.82 mmol, 1.1 eqv.), palladium acetate (8 mg, 0.04 mmol, 5 mol %), NaHCO₃ (94 mg, 1.12 mmol, 1.5 eqv.) and *n*-Bu₄N⁺Br⁻ (723 mg, 2.24 mmol, 3eqv.) in DMF (5 mL) was taken in a seal tube and stirred at 90 °C for 18h. The reaction mixture was allowed to come to r. t.. and then diluted with H₂O (60 mL). It was then extracted with EtOAc (2 × 50 mL). The combined organic extract was washed successively with H₂O (2 × 50 mL) and brine (50 mL), and then dried

(MgSO₄). It was filtered, and then concentrated in vacuo, to leave a residue which was purified over silica gel (EtOAc–PE, 3:7) to provide **11** as a viscous liquid; (221 mg, 80 %); [α]_D + 19.3 (c 0.46, CHCl₃). IR (CHCl₃): 2980, 2934, 1695, 1570, 1391, 1366, 1258 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.98 (d, *J* = 3.6 Hz, 1H), 8.08 (d, *J* = 8.0 Hz, 1H), 8.02 (s, 1H), 7.79 (d, *J* = 8.0 Hz, 1H), 7.75-7.65 (m, 1H), 7.54 (t, *J* = 6.8 Hz, 1H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.44 (td, *J* = 15.2, 7.2 Hz, 1H), 4.14 (s, 1H), 3.99 (dd, *J* = 8.8, 4.8 Hz, 1H), 3.88 (d, *J* = 8.0 Hz, 1H), 2.78-2.68 (m, 1H), 2.62-2.54 (m, 1H), 1.53-1.50 (merged singlets, 15H). ¹³C NMR (100MHz, CDCl₃): δ 152.2 (151.7), 149.3 (149.1), 147.3, 132.0 (131.9), 130.3 (130.1), 129.6, 129.4, 129.2, 129.1 (129.0), 128.0, 127.7, 126.9 (126.8), 94.1 (93.5), 80.2 (79.7), 66.6 (66.4), 57.1 (56.9), 37.7 (36.9), 28.6 (28.4), 27.5 (26.7), 24.5 (23.2). Anal.Calcd for C₂₂H₂₈N₂O₃: C, 71.71; H, 7.66; N, 7.60; Found: C, 71.56; H, 7.85; N, 7.71. MS (TOF, ES⁺): *m/z* (%) 369 (100) [M+H].

(*R,E*)-tert-butyl 2,2-dimethyl-4-(4-(quinolin-3-yl)but-3-enyl)oxazolidine-3-carboxylate (12)

A solution of the olefin **1c** (190 mg, 0.75 mmol), 3-bromoquinoline (171 mg, 0.82 mmol, 1.1 eqv.), palladium acetate (8 mg, 0.04 mmol, 5mol %), NaHCO₃ (94 mg, 1.12 mmol, 1.5 eqv.) and *n*-Bu₄N⁺Br⁻ (723 mg, 2.24 mmol, 3eqv.) in DMF (5 mL) was taken in a seal tube and stirred at 90 °C for 18h. The reaction mixture was allowed to come to r. t.. and then diluted with H₂O (60 mL). It was then extracted with EtOAc (2 × 50 mL). The combined organic extract was washed successively with H₂O (2 × 50 mL) and brine (50 mL), and then dried (MgSO₄). It was filtered, and then concentrated in vacuo, to leave a residue which was purified over silica gel (EtOAc–PE, 3:7) to provide **12** as a viscous liquid (219 mg, 77 %); [α]_D + 17.7 (c 0.46, CHCl₃). IR (CHCl₃): 2979, 2934, 2874, 2575, 1694, 1568, 1392, 1377, 1259 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.97 (s, 1H), 8.06 (d, *J* = 8.4 Hz, 1H), 7.99 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 1H), 7.66 (s, 1H), 7.53 (d, *J* = 6.8 Hz, 1H), 6.59-6.49 (m, 2H), 3.98-

3.81 (m, 3H), 2.32-2.30 (m, 2H), 2.10-1.90 (m, 1H), 1.7-1.49 (merged singlets, 16H). ^{13}C NMR (100MHz, CDCl_3): δ 152.3 (151.8), 149.3 (149.1), 147.2, 132.6 (132.3), 131.7, 130.5 (130.3), 129.2, 129.0 (128.9), 127.7, 127.4, 127.1, 126.8, 93.8 (93.3), 80.2 (79.6), 66.9 (66.7), 57.3 (56.8), 33.0 (32.3), 30.1 (29.7), 28.6 (28.5), 27.6 (26.8), 24.5 (23.2). Anal.Calcd for $\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_3$: C, 72.22; H, 7.91; N, 7.32; Found: C, 72.43; H, 7.85; N, 7.57. MS (TOF, ES⁺): m/z (%) 383 (100) [M+H].

Hydrogenation of the coupled products **11** & **12**:

(*S*)-tert-butyl 2,2-dimethyl-4-(3-(quinolin-3-yl)propyl)oxazolidine-3-carboxylate (**13**)

Pd-C (10 mol %; 14 mg) was added to a solution of the coupled product **11** (200 mg, 0.54 mmol) in MeOH (6 mL) at r.t. and the heterogeneous mixture was vigorously stirred under hydrogen atmosphere for 2 h. It was then filtered through celite and the filter cake was thoroughly washed with MeOH (10 mL). The combined filtrate was concentrated in vacuo to leave a crude mass which was purified by chromatography over silica gel (EtOAc-PE, 3:7) to furnish the product **13** as a viscous liquid (179 mg, 90 %); $[\alpha]_{\text{D}} + 4.49$ (c 0.22, CHCl_3). IR (CHCl_3): 2979, 2935, 2870, 1694, 1605, 1496, 1391, 1258 cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ 8.78 (d, $J = 1.6$ Hz, 1H), 8.07 (d, $J = 8.4$ Hz, 1H), 7.93 (s, 1H), 7.77 (d, $J = 8.0$ Hz, 1H), 7.66 (t, $J = 6.8$ Hz, 1H), 7.52 (d, $J = 7.2$ Hz, 1H), 3.95-3.72 (m, 3H), 2.84 (br s, 2H), 1.89-1.60 (m, 4H), 1.59-1.35 (m, 15H). ^{13}C NMR (100MHz, CDCl_3): δ 150.9, 145.9, 133.9, 133.6, 133.2, 128.1, 127.6, 127.1, 126.3, 125.6, 92.7 (92.2), 79.1, 65.9, 56.3, 32.1 (31.8), 28.7, 27.4, 26.9 (26.6), 25.7, 23.5 (22.2). Anal.Calcd for $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_3$: C, 71.32; H, 8.16; N, 7.56; Found: C, 71.47; H, 8.35; N, 7.73. MS (TOF, ES⁺): m/z (%) 371 (100) [M+H].

(*R*)-tert-butyl 2,2-dimethyl-4-(4-(quinolin-3-yl)butyl)oxazolidine-3-carboxylate (**14**)

Pd-C (10 mol %; 14 mg) was added to a solution of the coupled product **12** (210 mg, 0.55 mmol) in MeOH (6 mL) at r.t. and the heterogeneous mixture was vigorously stirred under hydrogen atmosphere for 2 h. It was then filtered through celite and the filter cake was thoroughly washed with MeOH (10 mL). The combined filtrate was concentrated in vacuo to leave a crude mass which was purified by chromatography over silica gel (EtOAc-PE, 3:7) to furnish the product **13** as a viscous liquid (196 mg, 92 %); $[\alpha]_D^{25} +15.1$ (c 0.29, CHCl₃). IR (CHCl₃): 2977, 2925, 2853, 1694, 1456, 1390, 1366, 1258 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.78 (d, J = 6.0 Hz, 1H), 8.08 (d, J = 8.4 Hz, 1H), 7.92 (s, 1H), 7.77 (d, J = 7.2 Hz, 1H), 7.66 (t, J = 6.4 Hz, 1H), 7.53 (t, J = 7.2 Hz, 1H), 3.93-3.89 (m, 2H), 3.72-3.70 (m, 2H), 2.99-2.75 (m, 2H), 1.79-1.71 (m, 5H), 1.59-1.45 (merged singlets, 15H). ¹³C NMR (100MHz, CDCl₃): δ 152.1, 146.8, 134.9, 134.1, 130.3, 129.2, 128.6, 128.2, 127.3, 126.6, 93.7 (93.1), 79.9 (79.4), 67.0 (66.7), 57.6 (57.3), 33.6 (33.1), 32.7, 30.9, 28.5, 27.6 (26.8), 25.9, 24.5 (23.3). Anal.Calcd for C₂₃H₃₂N₂O₃: C, 71.84; H, 8.39; N, 7.29; Found: C, 71.61; H, 8.46; N, 7.38. MS (TOF MS ES⁺): m/z (%) 385 (100) [M+H].

Jones' oxidation of the oxazolidines **13 & **14**:**

(S)-2-(tert-butoxycarbonylamino)-5-(quinolin-3-yl)pentanoic acid (15**)**

Jones' reagent (2.67 M, 366 μ L, 0.66 mmol) was added to a solution of the oxazolidine derivative **13** (95 mg, 0.26 mmol) in acetone (3 mL) at 0 °C. The reaction mixture was stirred for 5 h and then quenched by addition of 2-propanol (2.5 mL). The resulting reaction mixture was stirred for 10 min and then neutralized with sat aq NaHCO₃ solution. (pH=4-5). It was then extracted with ethyl acetate (2 $\times$ 25 mL) and the combined organic extract was washed successively with H₂O (20 mL) and brine solution (15 mL) and then dried over MgSO₄. It was then filtered and the filtrate was concentrated under reduced pressure to leave the crude product which was purified by column chromatography over silica gel (EtOAc-PE, 7:3) to

obtain compound **15** as a colourless liquid (65 mg, 74%). $[\alpha]_D + 28.2$ (c 0.29, CHCl₃). IR (CHCl₃): 3430, 2925, 2853, 1705, 1608, 1499, 1393, 1255, 1170 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 12.51 (s, 1H), 8.82 (d, J = 1.0 Hz, 1H), 8.14 (d, J = 8.4 Hz, 1H), 8.06 (s, 1H), 7.78 (d, J = 8.4 Hz, 1H), 7.67 (t, J = 7.6 Hz, 1H), 7.55 (t, J = 7.6 Hz, 1H), 5.49 (d, J = 7.6 Hz, 1H), 4.45 (d, J = 5.6 Hz, 1H), 2.86 (s, 2H), 1.88-1.61 (m, 4H), 1.45-1.37 (m, 9H). ¹³C NMR (100MHz, CDCl₃): δ 175.6, 155.6, 149.8, 144.1, 136.8, 135.2, 129.7, 128.3, 127.5, 127.4, 126.7, 79.6, 53.4, 32.7, 29.7, 28.4, 26.8. HRMS (TOF, ES⁺): m/z [M^+ + H] calcd for C₁₉H₂₅N₂O₄: 345.1814; found: 345.1807.

(R)-2-(tert-butoxycarbonylamino)-6-(quinolin-3-yl)hexanoic acid (16)

Jones' reagent (2.67 M, 366 μ L, 0.66 mmol) was added to a solution of the oxazolidine derivative **14** (100 mg, 0.26 mmol) in acetone (3 mL) at 0 °C. The reaction mixture was stirred for 5 h and then quenched by addition of 2-propanol (2.5 mL). The resulting reaction mixture was stirred for 10 min and then neutralized with sat aq NaHCO₃ solution. (pH=4-5). It was then extracted with ethyl acetate (2 $\times$ 25 mL) and the combined organic extract was washed successively with H₂O (20 mL) and brine solution (15 mL) and then dried over MgSO₄. It was then filtered and the filtrate was concentrated under reduced pressure to leave the crude product which was purified by column chromatography over silica gel (EtOAc-PE, 7:3) to obtain compound **16** as a colourless liquid (72 mg, 77%). $[\alpha]_D + 14.0$ (c 0.26, CHCl₃). IR (CHCl₃): 3436, 2926, 2856, 1702, 1500, 1456 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.12-8.78 (m, 1H), 8.14-8.02 (m, 2H), 7.82-7.56 (m, 1H), 7.68-7.62 (m, 1H), 7.58-7.49 (m, 1H), 5.48-5.39 (m, 1H), 5.31 (d, J = 7.2 Hz, 1H), 4.38 (d, J = 6.4 Hz, 1H), 2.89-2.82 (m, 2H), 2.05-1.78 (m, 4H), 1.47-1.39 (merged singlets, 10H). ¹³C NMR (100MHz, CDCl₃): δ 176.7, 155.6, 150.3, 144.6, 136.2, 135.4, 129.5, 128.3, 127.4, 127.2, 127.0, 79.6, 53.6, 32.8, 30.5, 29.7, 28.4, 24.4. HRMS (TOF, ES⁺): m/z [M^+ + H] calcd for C₂₀H₂₇N₂O₄: 359.1971; found: 359.1981.

Peptide coupling:**(S)-methyl 2-((S)-2-(tert-butoxycarbonylamino)-5-(quinolin-3-yl)pentanamido)-3-phenylpropanoate (17)**

N-Methylmorpholine (75 μ L, 0.75 mmol) was added dropwise to a stirred suspension of *L*-Phe-OMe.HCl (50 mg, 0.3 mmol, 1eqv.) in dry CH₂Cl₂ (4 mL) at 0 °C under argon. After 10 minutes, a solution of the carboxylic acid **15** (104 mg, 0.3 mmol, 1 eqv) in dry CH₂Cl₂ (4 mL) was added dropwise and stirred for 10 min. Then EDC.HCl (64 mg, 0.34 mmol, 1.1 eqv.) and HOBT (46 mg, 0.34 mmol, 1.1 eqv.) was added successively with 10 minutes interval at the same temperature. The reaction mixture was allowed to come to room temperature and stirred for 18 h. It was then diluted with water (25 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layer was washed successively with H₂O (2 $\times$ 20 mL) and brine (15 mL). It was then dried over Na₂SO₄, filtered and the filtrate was concentrated in vacuo to leave a pale yellow liquid, which was purified by chromatography over silica gel (EtOAc–PE, 1:1) as eluent to give the product **17** as a colourless solid (114 mg, 78%). Mp. 98-100 °C. $[\alpha]_D^{25} +26.9$ (c 0.24, CHCl₃). IR (CHCl₃): 3317, 2954, 2927, 2855, 1745, 1665, 1498, 1367, 1251, 1170 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.74 (s, 1H), 8.07 (d, *J* = 8.4 Hz, 1H), 7.90 (s, 1H), 7.76 (d, *J* = 8.0 Hz, 1H), 7.66 (t, *J* = 7.6 Hz, 1H), 7.52 (t, *J* = 7.6 Hz, 1H), 7.24 (d, *J* = 7.2 Hz, 2H), 7.08 (d, *J* = 8.4 Hz, 2H), 6.50 (d, *J* = 7.6 Hz, 1H), 5.02 (d, *J* = 7.2 Hz, 1H), 4.87-4.84 (m, 1H), 4.14 (d, *J* = 5.6 Hz, 1H), 3.70 (s, 3H), 3.17-3.06 (m, 2H), 2.83-2.75 (m, 2H), 2.10-1.80 (m, 2H), 1.74 (t, *J* = 7.2 Hz, 2H), 1.63-1.62 (s, 1H), 1.43 (s, 9H). ¹³C NMR (100 MHz, CDCl₃): δ 171.6, 171.5, 155.6, 151.8, 146.8, 135.6, 134.4, 129.2, 129.1, 128.7, 128.6, 128.1, 127.4, 127.2, 126.7, 80.2, 54.2, 53.1, 52.4, 37.8, 32.7, 32.1, 28.3, 27.0. Anal. Calcd for C₂₉H₃₅N₃O₅: C, 68.89; H, 6.98; N, 8.31; Found: C, 68.77; H, 7.05; N, 8.52. MS (TOF, ES⁺): *m/z* (%) 506 (100) [M+H].

(*R*)-methyl 2-((*R*)-2-(tert-butoxycarbonylamino)-6-(quinolin-3-yl)hexanamido)-3-phenylpropanoate (18)

N-Methylmorpholine (40 μ L, 0.4 mmol) was added dropwise to a stirred suspension of *L*-Phe-OMe.HCl (25 mg, 0.15 mmol, 1eqv.) in dry CH₂Cl₂ (2 mL) at 0 °C under argon. After 10 minutes, a solution of the carboxylic acid **16** (54 mg, 0.15 mmol, 1 eqv) in dry CH₂Cl₂ (2 mL) was added dropwise and stirred for 10 min. Then EDC.HCl (32 mg, 0.17 mmol, 1.1 eqv.) and HOBT (23 mg, 0.17 mmol, 1.1 eqv.) was added successively with 10 minutes interval at the same temperature. The reaction mixture was allowed to come to room temperature and stirred for 18 h. It was then diluted with water (25 mL) and extracted with EtOAc (2 x 25 mL). The combined organic layer was washed successively with H₂O (2 $\times$ 20 mL) and brine (15 mL). It was then dried over Na₂SO₄, filtered and the filtrate was concentrated in vacuo to leave a pale yellow liquid, which was purified by chromatography over silica gel (EtOAc–PE, 1:1) as eluent to give the product **18** as a colourless liquid (58 mg, 75%). [α]_D +12.4 (*c* 0.25, CHCl₃). IR (CHCl₃): 3318, 2956, 2926, 2851, 1750, 1666, 1499, 1368 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ 8.75 (d, *J* = 2.0 Hz, 1H), 8.07 (d, *J* = 8.0 Hz, 1H), 7.90 (s, 1H), 7.77 (d, *J* = 7.6 Hz, 1H), 7.66 (t, *J* = 8.4 Hz, 1H), 7.52 (t, *J* = 7.6 Hz, 1H), 7.31-7.18 (m, 5H), 7.09 (d, *J* = 6.8 Hz, 2H), 6.43 (d, *J* = 7.6 Hz, 1H), 4.92 (d, *J* = 5.6 Hz, 1H), 4.85 (dd, *J* = 13.6, 6.0 Hz, 1H), 4.06 (d, *J* = 7.2 Hz, 1H), 3.63 (s, 3H), 3.16-3.07 (m, 2H), 2.78 (t, *J* = 7.6 Hz, 2H), 1.84-1.58 (m, 2H), 1.42 (s, 9H), 1.28-1.25 (m, 2H). ¹³C NMR (100MHz, CDCl₃): δ 171.7, 155.5, 151.9, 151.0, 146.8, 135.7, 134.8, 134.2, 129.2, 129.1, 128.6, 128.1, 127.5, 127.3, 127.1, 126.6, 80.1, 54.4, 53.1, 52.3, 37.9, 32.9, 32.3, 30.7, 28.3, 25.1. Anal. Calcd for C₃₀H₃₇N₃O₅: C, 69.34; H, 7.18; N, 8.09; Found: C, 69.57; H, 7.25; N, 8.21. MS (TOF, ES⁺): *m/z* (%) 520 (100) [M+H].

IK-Ser-13

22.11.2012

^1H NMR spectrum (400 MHz, CDCl_3) of **1a**

SER-WITTIG

21.10.2013

¹H NMR spectrum (400 MHz, DMSO-d₆) of **1a**

IK-Ser-13

22.11.12

^{13}C NMR spectrum (100 MHz, CDCl_3) of **1a**

IK-SER-14

26.11.2012

^1H NMR spectrum (400 MHz, CDCl_3) of **7**

Heck SER

25.10.2013

¹H NMR spectrum (400 MHz, DMSO-d₆) of 7

^{13}C NMR spectrum (100 MHz, CDCl_3) of **7**

SER-14 13 at 343K

Current Data Parameters
 NAME SER-14
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20131008
 Time 15.53
 INSTRUM spect
 PROBHD 5 mm PABBO BB-
 PULPROG zgpg30
 TD 32768
 SOLVENT DMSO
 NS 64
 DS 2
 SWH 21929.824 H
 FIDRES 0.669245 H
 AQ 0.7471104 s
 RG 64.79
 DW 22.800 u
 DE 6.50 u
 TE 343.0 K
 D1 2.00000000 s
 D11 0.03000000 s
 TD0 1

===== CHANNEL f1 =====
 SFO1 75.4752949 MHz
 NUC1 13C
 P1 9.25 u
 PLW1 34.00000000 u

===== CHANNEL f2 =====
 SFO2 300.1312005 MHz
 NUC2 1H
 CPDPRG[2] waltz16
 PCPD2 90.00
 PLW2 6.99840021 u
 PLW12 0.17545000 u
 PLW13 0.14211001 u

F2 - Processing parameters
 SI 32768
 SF 75.4677380 MHz
 WDW EM
 SSB 0
 LB 1.00
 GB 0
 PC 1.40

^{13}C NMR spectrum (100 MHz, DMSO- d_6) of **7** at 70 °C

^1H NMR spectrum (400 MHz, CDCl_3) of **8**

IK-SER-15

11.05.12

^{13}C NMR spectrum (100 MHz, CDCl_3) of **8**

IK-Ser-16

27.12.2012

¹H NMR spectrum (400 MHz, CDCl₃) of **9**

^{13}C NMR spectrum (100 MHz, CDCl_3) of **9**

IK-SER-17

19.12.2012

¹H NMR spectrum (400 MHz, CDCl₃) of **10**

^1H NMR spectrum (400 MHz, CDCl_3) of **1b**

^{13}C NMR spectrum (100 MHz, CDCl_3) of **1b**

¹H NMR spectrum (400 MHz, CDCl₃) of **11**

IK-ASP-14

18.4.2011

152.20
151.71
149.28
149.10
147.28
132.01
131.87
130.30
130.06
129.64
129.40
129.16
129.08
128.96
128.04
127.74
126.93
126.85

94.09
93.54

80.22
79.72
77.40
77.08
76.76

66.63
66.44

57.05
56.93

37.68
36.86
29.68
28.55
28.42
27.53
26.74
24.50
23.23

11 ¹³C NMR spectrum (100 MHz, CDCl₃) of **11**

¹H NMR spectrum (400 MHz, CDCl₃) of **13**

^{13}C NMR spectrum (100 MHz, CDCl_3) of **13**

¹H NMR spectrum (400 MHz, CDCl₃) of **15**

^{13}C NMR spectrum (100 MHz, CDCl_3) of **15**

¹H NMR spectrum (400 MHz, CDCl₃) of **17**

¹³C NMR spectrum (100 MHz, CDCl₃) of **17**

^1H NMR spectrum (400 MHz, CDCl_3) of **1c**

IK-Ghi-13

20.11.12

^{13}C NMR spectrum (100 MHz, CDCl_3) of **1c**

^1H NMR spectrum (400 MHz, CDCl_3) of **12**

^{13}C NMR spectrum (100 MHz, CDCl_3) of **12**

¹H NMR spectrum (400 MHz, CDCl₃) of **14**

^1H NMR spectrum (400 MHz, CDCl_3) of **16**

IK-Giu-17

16.01.2013

^1H NMR spectrum (400 MHz, CDCl_3) of **18**

¹³C NMR spectrum (100 MHz, CDCl₃) of **18**