Unsupervised Change Detection Driven by Floating References

A Pattern Analysis Approach

Rogério G. Negri · Alejandro C. Frery

Received: 24 June 2019 / Accepted: date

Abstract The Earth's environment is continually changing due to both human and natural factors. Timely identification of the location and kind of change is of paramount importance in several areas of application. Because of that, Remote Sensing change detection is a topic of great interest. The development of precise change detection methods is a constant challenge. This study introduces a novel unsupervised change detection method based on data clustering and optimization. The proposal is less dependent on radiometric normalization than classical approaches. We carried experiments with remote sensing images and simulated data sets to compare the proposed method with other unsupervised well-known techniques. At its best, the proposal improves by 50 % the accuracy concerning the second best technique. Such improvement is most noticeable with uncalibrated data. Experiments with simulated data reveal that the proposal is better than all other compared methods at any practical significance level. The results show the potential of the proposed method.

Keywords Unsupervised change detection · Pattern Analysis · Remote Sensing

The authors thank FAPESP (Grant 2018/01033-3), and CNPq (Grants 303267/2019-4 and 405364/2018-0).

Rogério G. Negri
Institute of Science and Technology – ICT
São Paulo State University – UNESP
Department of Environmental Engineering
São José dos Campos, Brazil
ORCID 0000-0002-4808-2362
E-mail: rogerio.negri@unesp.br

Alejandro C. Frery
School of Mathematics and Statistics
Victoria University of Wellington, 6140 New Zealand
ORCID 0000-0002-8002-5341
E-mail: alejandro.frery@vuw.ac.nz

1 Introduction

Automatic land cover change detection using remote sensing images is fundamental, for example, in environmental sciences, cartography, geography, ecology, and hydrography (Srivastava et al., 2012). This class of techniques aims to identify the changes between two or more dates through the images acquired in those or close instants (Liu et al., 2015).

The detection of changes cannot rely solely on data but should be supported by the comprehension of the interactions between anthropic and natural phenomena. Because of this, the development of accurate change detection techniques is a constant challenge (Lu et al., 2004; Ross and Bhadauria, 2015).

All the methods discussed below rely on registered images for comparison.

The most successful change detection techniques are deeply rooted in concepts from Pattern Recognition and Digital Image Processing. According to Wu et al. (2014), change detection techniques may be categorized as: (i) image algebra; (ii) image transformations; (iii) feature extraction; (iv) image classification; and (v) other methods. These techniques may also be distinguished between supervised and unsupervised (Volpi et al., 2012). While supervised methods require labeled data for training, the unsupervised approach is automatically guided by clustering and thresholding processes. The independence of prior knowledge makes unsupervised methods more attractive in comparison to supervised.

Change Vector Analysis (CVA) is a classic example of an unsupervised technique. It appeared in the early 80s (Colwell and Weber, 1981) and became popular later in the 90s (Johnson and Kasischke, 1998). CVA uses the amplitude difference between the attribute vectors of pixels from two images acquired in distinct instants over the same spatial position. In the sequence, the resulting amplitude difference image is thresholded, for instance, by the well-known Otsu's (1979) method, to produce a map of land cover changes.

The use of morphological filters also yields change detection maps with low complexity algorithms. Mura et al. (2008) showed the potential of Alternating Sequential Filters (ASF), based on iterative opening-closing operations over an amplitude difference image, likewise computed by CVA, to highlight change regions and smooth the non-changes. Lastly, and similarly to CVA, the resulting filtered data is thresholded using the Kittler and Illingworth's (1986) method.

Unlike those approaches, Celik (2009) proposed an "spectral expansion" of the amplitude difference image using contextual information, i.e., the neighborhood, followed by a re-projection of this extended space using Principal Component Analysis (PCA). The pixels of the resulting image, whose center is closer to the feature space's origin, are clustered into non-change using all or some Principal Components, while the others are assigned to changes.

Motivated by the behavior of the change and non-change amplitude difference values, the Gustafson and Kessel's (1978) Clustering (GKC) algorithm analyses two bands: the amplitude difference information and the context of amplitude differences (computed through a spatial averaging of pixels) in order to identify the two clusters. The GKC parameters are tuned using Simulated Annealing (Kirkpatrick et al., 1983), considering as objective function the Xie and Beni's (1991) clustering assessment

measure. Similarly to the previous method, the cluster closest to the feature space's origin is assigned to pixels without change.

The methods above require a radiometric correction or a previous feature extraction from the input data. Such a requirement may be a limiting factor in some cases, such as when there are no available models to correct or normalize the images, or when such models are inadequate or too demanding.

This research proposes a new unsupervised change detection method based on non-fixed (floating) references identified utilizing clustering. The use of non-fixed references is a crucial feature that makes this technique independent of requirements like radiometric normalization and other pre-processing. Such floating references act as an attention model.

Attention models in remotely sensed data have been successfully employed, mostly in classification and segmentation, aided with neural networks (Mou and Zhu, 2020). Our approach is simpler, in the sense that we find the regions that deserve attention by explicit spectral and spatial operations and, thus, are fully interpretable and adaptable to specific problems.

The main contribution of this paper is an innovative unsupervised change detection method, which

- relies on simple clustering and statistical concepts;
- is robust before the lack of data calibration;
- is reproducible and extensible, as we provide its implementation.

We report experiments using multispectral remote sensing images and simulated data, both radiometrically processed and not, to check the proposed method's potential. We also include comparisons with the above-cited methods.

2 Background

We introduce a new unsupervised change detection technique in Section 2.2, after recalling some preliminary concepts and notations in Section 2.1. Section 2.3 presents remarks to complete the discussions and formalization.

2.1 Preliminary concepts and notations

Let $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ be images defined over the same support $\mathcal{S} \subset \mathbb{N}^2$ and acquired over the same scene region in distinct instants. Denote $\mathcal{X}$ the set of their possible values, and $\mathbf{x}_i^{(j)} = \mathcal{I}^{(j)}(s_i)$ the observation $\mathbf{x}_i^{(j)} \in \mathcal{X}$ at position $s_i \in \mathcal{S}$ of $\mathcal{I}^{(j)}$, for $j = 1, 2$. The components of $\mathbf{x}_i^{(j)}$ are the values measured by the sensor, or derived features, over a specific Earth surface position.

We define data clusters as a partition of the support $\mathcal{S}$, obtained by means of a function $\mathcal{G}^{(j)}: \mathcal{S} \rightarrow \{\mathcal{G}_1^{(j)}, \dots, \mathcal{G}_c^{(j)}\}$. The c clusters $\mathcal{G}_k^{(j)}$, $k = 1, \dots, c$, satisfy $\mathcal{G}_k^{(j)} \neq \emptyset$, $\bigcup_{k=1}^c \mathcal{G}_k^{(j)} = \mathcal{S}$ and $\bigcap_{k=1}^c \mathcal{G}_k^{(j)} = \emptyset$. The partition function is such that $\mathcal{G}_k^{(j)}$, $k = 1, \dots, c$, represents a set of positions s_i that verify (Theodoridis and Koutroumbas, 2008)

$$\mathcal{G}_k^{(j)} = \{\mathbf{x}_i : m(\mathbf{x}_i, \mathbf{x}_k^*) < m(\mathbf{x}_i, \mathbf{x}_\ell^*)\}; \forall k \neq \ell, \quad (1)$$

where $\mathbf{x}_k^*$ is the medoid for $\mathcal{G}_k^{(j)}$ and $m(\cdot, \cdot)$ is a dissimilarity function. The dissimilarity function and the medoids specify a partition. Obtaining a partition function able to define clusters that consistently express the data behavior may comprise a difficult task. It may be alleviated, for instance, by specifying the number of clusters in advance.

We will use $z_i \in \mathbb{R}$ as a measure of the possible changes between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ at s_i . We denote $z_i = u(\mathbf{x}_i^{(1)}, \mathbf{x}_i^{(2)})$ where $u: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ is a dissimilarity function between elements of $\mathcal{X}$. The collection of all change measures is also an image defined on the same support $\mathcal{Z}: \mathcal{S} \rightarrow \mathbb{R}$. A possible measure of change frequently used in the literature (Johnson and Kasischke, 1998) is the L_2 norm between $\mathbf{x}_i^{(1)}$ and $\mathbf{x}_i^{(2)}$:

$$\mathcal{Z}(s_i) = \|\mathbf{x}_i^{(1)} - \mathbf{x}_i^{(2)}\|_2. \quad (2)$$

Finally, we define neighborhoods on the support. Denote $\mathcal{V}_\rho(s_i) = \{t : d(s_i, t) \leq \rho\}$ all the coordinates $t \in \mathcal{S}$ that are at most at distance ρ from s_i . The distance d may be, for instance, the infinite norm: $d(s, t) = \|s - t\|_\infty = \max\{|s_1 - t_1|, |s_2 - t_2|\}$. The neighborhood information may be coded into the feature vector of s_i as:

$$\mathbf{x}_i^{(j)} = \frac{\sum_{t \in \mathcal{V}_\rho(s_i)} \mathbf{x}_t^{(j)}}{\#\mathcal{V}_\rho(s_i)}, \quad (3)$$

where $\#$ denotes the cardinality of a set.

2.2 Floating references for unsupervised change detection

The following premises are the starting point for the formalization of the proposed unsupervised change detection method:

- (i) External factors related to the data acquisition process may influence the target representation, and such influence may be manifested differently depending on the target;
- (ii) There may be differences between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ unrelated to external acquisition factors.

The first premise assumes that some factors may influence the scene representation so that similar targets express differently when observed in two distinct instants. Atmospheric interference, seasonal variations, the sensor's angle/type of view, use of different channels (i.e., bands or polarizations), resolutions, or even different sensors are examples of external factors that may influence the target representation. Furthermore, this premise assumes that there is no known function able to model and correct the variations of target representation between two distinct instants. These factors are *irrelevant* to the change detection process. The second premise imposes that there might be differences or variations between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ that cannot be explained by external factors; such differences and variations are temporal changes whose detection is desirable.

The first premise motivates the use of *non-fixed references* to compare $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ so that differences/variations related to irrelevant factors are discarded. For this

purpose, considering the initial instant $\mathcal{I}^{(1)}$, the clusters $\mathcal{G}_k^{(1)}$, $k = 1, \dots, c$ isolate different targets into separate sets. These sets model the regions that deserve attention and are our version of an attention model for the problem at hand.

The possible changes that occur between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ are verified in the clusters $\mathcal{G}_k^{(1)}$, $k = 1, \dots, c$. We, thus, define the sets:

$$\mathcal{Z}(\mathcal{G}_k^{(1)}) = \{\mathcal{Z}(s_i) : s_i \in \mathcal{G}_k^{(1)}\}, k = 1, \dots, c. \quad (4)$$

In order to obtain this, we need to smooth the data. We achieve this by using the neighboring information of each element of the partition in the two epochs.

The values in $\mathcal{Z}(\mathcal{G}_k^{(1)})$, by construction, are concentrated around a central tendency. Deviations from this tendency is related to either relevant (changes) or irrelevant factors. Irrelevant changes cause mild deviations, while changes induce substantial alterations. The concepts of mild and significant will be described with respect to a central tendency in a nonparametric manner.

Denote f_k the probability density function that characterizes the distribution of $\mathcal{Z}(\mathcal{G}_k^{(1)})$. Given an arbitrary $0 < \alpha < 1$, then the tendency, denoted by τ_k , may be determined as the middle point of $[a, b]$ that arises from the solution of:

$$\min_{a < b} \left\{ b - a : \int_a^b f_k(z) dz = \alpha \right\}. \quad (5)$$

This way of computing the tendency is similar to Chernoff's (1964) kernel-based definition of the mode. Chacón (2020) praises the use of modal approaches in problems where it is not sensible to make strong hypothesis about the data distribution, as is the case we are handling here.

Note that τ_k acts as a *reference* for the elements of $\mathcal{Z}(\mathcal{G}_k^{(1)})$. As such, a large difference between an element of $\mathcal{Z}(\mathcal{G}_k^{(1)})$ and τ_k suggests that such an element is a temporal change. Once the reference τ_k is found for each cluster $\mathcal{Z}(\mathcal{G}_k^{(1)})$, for $k = 1, \dots, c$, we define a new difference image $\mathcal{F}$, with support $\mathcal{S}$:

$$\mathcal{F}(s_i) = |\mathcal{Z}(s_i) - \tau_k| = \tilde{z}_i; \forall s_i \in \mathcal{G}_k^{(1)}; k = 1, \dots, c. \quad (6)$$

It is worth observing that $\mathcal{F}$ measures the deviation from relative tendencies. Such tendencies are established with basis on the behavior of elements in $\mathcal{S}$ that belong to the same cluster (and consequently have similar behavior) in the first epoch $\mathcal{I}^{(1)}$.

As the comparison is made in objects that depend on the observed data, we call this *Floating Reference*.

Fig. 1(a) depicts hypothetical values in $\mathcal{Z}(\mathcal{G}_k^{(1)})$, where it is possible to observe a general tendency τ_k as well as values far from such tendency. In order to determine τ_k , we model a probability density function f_k from $\mathcal{Z}(\mathcal{G}_k^{(1)})$, and then we find the interval $[a, b]$ containing τ_k , as shown in Fig 1(b). The optimization problem (5) determines the smallest interval $[a, b]$ whose values present maximum probability of occurrence. This approach amounts to finding a robust tendency measure without committing to any particular probabilistic model, as would be the case if we used the mean or the median. In this way, we can cope with multimodal observations.

Fig. 1 The elements that define the tendency τ_k .

We measure the evidence of change by the absolute difference between τ_k and the elements of $\mathcal{Z}(\mathcal{G}_k^{(1)})$. We denote as $\mathcal{F}$ all the absolute differences found for each $\mathcal{Z}(\mathcal{G}_k^{(1)})$ with respect its τ_k , $k = 1, \dots, c$. Fig.2 shows a histogram of such values.

We then generate a change/non-change map from $\mathcal{F}$ by choosing a threshold η , above which we consider there has been a change. We do this by selecting two sets of differences: short-range low-variance and long-range high-variance.

Let g be the probability density function that models $\mathcal{F}$. The following empirical rule finds η which splits the values:

$$\eta = \arg \max_{w \in [\tilde{z}_{\min}, \tilde{z}_{\max}]} \left\{ \frac{C_1 C_2 (C_1 + C_2)}{A_1 A_2 (A_1 + A_2)} \right\}, \quad (7)$$

where

$$\begin{aligned} C_1 &= \int_{\tilde{z}_{\min}}^w g(\tilde{z}) d\tilde{z}, & C_2 &= \int_w^{\tilde{z}_{\max}} g(\tilde{z}) d\tilde{z}, \\ A_1 &= (w - \tilde{z}_{\min}) \max_{\tilde{z} \in [\tilde{z}_{\min}, w]} \{g(\tilde{z})\}, & A_2 &= (\tilde{z}_{\max} - w) \max_{\tilde{z} \in [w, \tilde{z}_{\max}]} \{g(\tilde{z})\}. \end{aligned}$$

Fig. 2 Geometry of empirical rule to determine the threshold η .

While C_1 and C_2 are the areas under g up to and above η , respectively, A_1 and A_2 are the areas of the rectangles that enclose g in $[\tilde{z}_{\min}, \eta]$ and $[\eta, \tilde{z}_{\max}]$, resp. The purpose is determining the best balance for both ratios C_1/A_1 and C_2/A_2 . An unbalanced choice for η implies $C_1/A_1 \rightarrow 0$ or $C_2/A_2 \rightarrow 0$, and consequently $\frac{C_1 C_2 (C_1 + C_2)}{A_1 A_2 (A_1 + A_2)} \rightarrow 0$. Fig. 2 depicts these elements, where C_1 and C_2 are the dotted and dashed areas under g ; A_1 and A_2 are the green and blue rectangles over g . We find the best balance when the areas under the curve are closest to the rectangle area.

Fig. 3 presents a general overview of the proposed method. The images at different epochs are $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$. The relevant changes are related to (i) modifications in the target structure, depicted by the change of position of small squares in the bottom of both images, and (ii) the shortening verified in the upper-right rectangle from $\mathcal{I}^{(1)}$ to $\mathcal{I}^{(2)}$. Irrelevant changes correspond to intensity difference and, consequently the “color aspect” distinction regarding some objects when comparing $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$. The neighborhood information is computed, cf. Equation (3), and coded in the feature vector of its pixels. The data are smoothed as a consequence of such operation. Then, a partition $\mathcal{G}^{(1)}$ of $\mathcal{I}^{(1)}$ is found along with its data clusters $\mathcal{G}_1^{(1)}$, $\mathcal{G}_2^{(1)}$ and $\mathcal{G}_3^{(1)}$. Such partition comes from $\mathcal{I}^{(1)}$ after the neighborhood information coding. Additionally, it is worth observing that $\mathcal{G}_1^{(1)}$ corresponds to two spatially disconnected objects, but with similar spectral behavior in $\mathcal{I}^{(1)}$. The difference image $\mathcal{Z}$, as defined in Equation (2), is also computed from neighborhoods coded in $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$. As shown, $\mathcal{Z}$ expresses the amplitude differences derived between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$. We fuse geometrical information (the clusters) with radiometric information (the difference image) in the next step. We compute central tendencies of differences $\mathcal{Z}$ in each cluster $\mathcal{G}_k^{(1)}$, and obtain τ_k for each with Equation (5). It is important to note that the geometric bounds defined by these clusters allow to isolate/mask similar objects (even spatially unconnected) of $\mathcal{I}^{(1)}$ in $\mathcal{Z}$, and to check their behavior. The deviation from each τ_k

yields the image $\mathcal{F}$ (Equation (6)). Since the deviations are computed with references that “float” as a function of the object’s behavior in $\mathcal{I}^{(1)}$, the resulting $\mathcal{F}$ tends to highlight the differences from $\mathcal{I}^{(1)}$ to $\mathcal{I}^{(2)}$.

Fig. 3 General overview of proposed method: from input images $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$, to the change/no-change map $\mathcal{F}$.

2.3 Remarks

The previous formalization was simplified for illustration purposes. The following information completes the proposal.

- Among the clustering methods available in the literature for finding $\mathcal{G}_k^{(1)}$, a divisive hierarchical schema combined with binary K-Means algorithm (Theodoridis and Koutroumbas, 2008) proved to be convenient for our proposal. The data set is initially comprised of all the feature vectors of $\mathcal{I}^{(1)}$. Such a dataset is split recursively into two subsets building a binary tree structure. The splitting process stops when each leaf's cardinality attains an inferior bound β , specified as a percentage of the problem size. The leaves are the clusters found in $\mathcal{I}^{(1)}$.
- The neighborhood radius ρ and distance d are chosen to smooth the change/non-change results. Larger values of ρ yield a less sensitive procedure, while smaller ones may lead to an increased false alarm. Consequently, ρ should be set considering the input image characteristics as well as the application purpose. The expression for d defines the neighborhood geometry. It is usually chosen as the infinity norm to ease the implementation, as it implies square-shaped neighborhoods.
- One of the key points in this *Floating Reference* method is the use of probability distributions to model the magnitude differences between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ conditioned on the clusters $\mathcal{G}_k^{(1)}$, i.e., f_k , for $k = 1, \dots, c$, as well as the model g for $\mathcal{F}$. These distributions are inferred using Gaussian kernel density estimators. Furthermore, we used Silverman's (1986) rule-of-thumb to select the kernel bandwidth;
- Regarding Equation (5), different choices of α lead to different intervals $[a, b]$. Preliminary analyses revealed that $\alpha \in [0.05, 0.25]$ leads to similar values of τ_k . In the face of this, we fixed α to 0.05.
- At last, beyond the geometric motivation, the proposed thresholding criterion, Equation (7), was also tested in several controlled experiments, including the datasets of Section 3. In all these cases, the values of η were close to cut-points that maximize the change/non-change detection accuracy. We also compared such a proposal with Otsu's (1979) and Kittler and Illingworth's (1986) algorithms, with favorable results to Equation (7).

3 Experiments and Results

In the following, we show the results of applying the proposed technique to real remote sensing and simulated data. The case studies presented in Section 3.1 focus on mapping areas affected by an environmental disaster and by extensive deforestation. Section 3.2 presents a quantitative analysis of the performance of the proposed technique. This is achieved using simulated data, where the correct results are known.

We measured the performance in both applications (i.e., with actual remote sensing images and simulated data), considering two plausible situations: with and without temporal radiometric normalization.

We included well-known unsupervised change detection methods for comparison, reported in terms of mapping accuracy (Precision, Recall, and F_1 -Score) and compu-

tational run-time. These are the methods discussed in Section 1: alternating sequential filter (ASF, Mura et al., 2008); spatial expansion with PCA followed by clustering with K-Means (PCA+K-Means, Celik, 2009); change vector analysis (CVA, Johnson and Kasischke, 1998); also, the Gustafson-Kessel clustering optimally parameterized using Simulated Annealing (GKC+SA, Ghosh et al., 2011). The proposed *Floating Reference* method will be referred to as FR.

The compared methods require specifying tuning parameters. Regarding the studies with actual remote sensing images, our choices were:

ASF: The number of opening-closing iterations is expressed by the parameter $\lambda \in \{1, 2, \dots, 20\}$.

PCA+K-Means: The neighborhood radius $\rho \in \{1, 2, \dots, 6\}$ to promote a spectral expansion (each value generates squared neighborhoods of side 1, 3, 5, $\dots$, 13), and $1 \leq \theta \leq \rho$ that defines the number of principal components considered.

CVA: The Otsu (*OT*) or Kittler-Illingworth (*KI*) thresholding options.

GKC+SA: The SA cooling factor $\alpha \in [0.8, 0.9]$.

We used the same set of remotely sensed images with each procedure and found the best parameter configuration. These parameters were then employed in the experiment with simulated data.

The experiments run on a computer with an Intel Core i7 processor, and 16 GB of RAM running the Debian Linux version 8.1 operating system. The platform was IDL (Interactive Data Language) version 8.5. Following the reproducibility guidelines discussed by Frery et al. (2020), we make the code and data available at <https://github.com/rogerionegri/FloatingReferences>.

3.1 Experiments with remote sensing images

This section presents two case studies regarding the mapping of temporal land cover changes through unsupervised change detection techniques. The first study aims at detecting the changes caused by the environmental disaster that occurred on November 5th, 2015, as a consequence of a dam rupture in Mariana, state of Minas Gerais, Brazil. According to do Carmo et al. (2017), the “Mariana’s Dam” case is the most massive worldwide environmental/technological disaster. The second study focuses on the mapping of extensive deforestation between 1988 to 1999 over the rain forest region near Belterra, state of Pará, Brazil. Fig. 4 illustrates the two study areas.

Both studies employed images acquired by LANDSAT sensors. Regarding Mariana’s Dam case, the 760×400 pixels wide images were obtained on September 25th, 2015, and August 10th, 2016, by the LANDSAT-8 OLI multispectral sensor, with 30 m spatial resolution, six multispectral bands from blue to short wave infrared. Similarly, the images adopted for Belterra Deforestation study were acquired by LANDSAT-5 TM sensor in August 22th 1988, and August 2nd, 1999, 30 m spatial resolution, the same six multispectral bands considered for the first study, and correspond to a portion of 900×500 pixels wide. Both image sets are georeferenced.

Figs. 5(a)-5(b) and 6(a)-6(b) present the images used in Mariana’s Dam and Belterra deforestation studies, respectively. Furthermore, Figs. 5(c) and 6(c) identify samples of change and non-change over the considered periods, represented by

Fig. 4 Study areas location.

polygons in green and red, respectively, comprising 7,712 and 7,258 pixels resp. in Mariana's Dam, and 15,014 and 32,270 in Belterra Deforestation areas. Such reference samples, obtained by visual inspection, were adopted to measure the accuracy of the methods in Precision, Recall, and F_1 -Score (Rijsbergen, 1979).

These measures are especially useful in binary classification problems. Based on the reference information, the correct identification of cases is expressed in True Positives (TP) and True Negatives (TN) rates. Wrong identifications lead to False Positive (FP) and False Native (FN) rates. Precision and Recall are:

$$\text{Pr} = \frac{\text{TP}}{\text{TP} + \text{FP}}, \text{ and } \text{Re} = \frac{\text{TP}}{\text{TP} + \text{FN}}.$$

The F_1 -Score is the harmonic mean of Pr and Re:

$$F_1 = 2 \frac{\text{Pr} \cdot \text{Re}}{\text{Pr} + \text{Re}}.$$

It is worth observing that more accurate results occur when Pr, Re, and F_1 simultaneously tend to 1. Low accuracy is observed if such measures are near 0.

In order to assess the robustness of the methods in terms of temporal radiometric normalization, both data sets were employed with and without atmospheric correction. Mariana's Dam images were processed with the FLAASH model Adler-Golden et al. (1999), and the Belterra data were transformed using the Tasseled Cap method Crist and Ciccone (1984).

Table 1 Parameter space search. Areas 1 and 2 correspond to Mariana’s Dam and Belterra Deforestation data sets, respectively. The symbols $\oplus$ and $\ominus$ denote, resp., whether the data sets were or were not corrected to reduce atmospheric effects.

Method	Parameters				
	Space search	Area 1		Area 2	
		$\oplus$	$\ominus$	$\oplus$	$\ominus$
ASF	$\lambda = \{1, 2, \dots, 20\}$	5	6	3	18
PCA+K-Means	$\rho = \{1, \dots, 6\}$	2	3	2	5
	$\theta = \{1, 2, \dots, 6\}$	3	4	2	3
CVA	threshold = $\{OT, KI\}$	2	2	2	2
GKC+SA	$\alpha = \{0.8, 0.81, \dots, 0.9\}$	0.8	0.82	0.8	0.88
FR	$\rho = \{0, 1, 2, \dots, 6\}$	1	1	2	1
	$\beta = \{1, 2.5, 5, 10, 25\} (\%)$	2.5	1	25	25

Table 1 shows the space search for the configuration that produces the best classification, over which we performed an exhaustive search.

Fig. 7 shows the spread of the F_1 -Score for all tested parameters. PCA+K-Means, CVA, and FR are the least sensitive procedures (i.e., the smallest F_1 -Score variation) to parameters changes. Moreover, FR achieves high accuracy values in comparison to the other methods regardless of the study area and the radiometric normalization, especially for Belterra dataset whose original images (Figs. 6(a) and 6(b)) strongly affects the performance of ASF, PCA+K-Means, CVA, and GKC+SA methods.

Table 2 presents the configurations that led to the best F_1 -Scores, along with the processing run times.

The values of Table 2 show that FR has the best F_1 -Score. ASF and PCA+K-Means were the most competitive methods in comparison to FR. Regarding the computational cost, FR was the second most expensive method. The clustering stage is the main reason behind the high computational cost.

The Precision and Recall values, shown in Table 2, suggest that FR has balanced Precision values through analyzed areas and conditions. Regarding the Recall, FR has intermediate values and reveals a conservative behavior concerning change events.

Figs. 8 and 9 present the best change/non-change maps. In general, it is possible to observe FR’s robustness when the input data are not radiometrically normalized, while the other methods are more suited for normalized input data. Furthermore, FR maps are closer to expected results regarding the spatial distribution of change/non-change areas.

3.2 Experiments with simulated data

We built two simulated datasets and adopted them as benchmark to access the methods. Section 3.2.1 presents a brief discussion about the simulation process. Section 3.2.2 presents the results and discussions obtained using the simulated images as input data.

Table 2 F_1 -Scores and run-times. Areas 1 and 2 correspond to Mariana’s Dam and Belterra Deforestation datasets, respectively. The symbols $\oplus$ and $\ominus$ denote, resp., whether the data sets were or were not corrected to reduce atmospheric effects.

Method	F_1 -Score				Time [sec.]			
	Area 1		Area 2		Area 1		Area 2	
	$\oplus$	$\ominus$	$\oplus$	$\ominus$	$\oplus$	$\ominus$	$\oplus$	$\ominus$
ASF	0.864	0.812	0.883	0.580	104.2	162.4	50.0	4655.6
PCA+K-Means	0.909	0.859	0.790	0.202	55.1	55.8	70.4	75.4
CVA	0.859	0.808	0.801	0.548	3.1	3.3	4.7	4.8
GKC+SA	0.739	0.749	0.851	0.542	7575.7	11245.7	7285.5	10344.7
FR	0.914	0.879	0.885	0.870	801.5	657.7	902.2	748.3

Method	Precision				Recall			
	Area 1		Area 2		Area 1		Area 2	
	$\oplus$	$\ominus$	$\oplus$	$\ominus$	$\oplus$	$\ominus$	$\oplus$	$\ominus$
ASF	0.792	0.702	0.948	0.408	0.951	0.963	0.826	1.000
PCA+K-Means	0.967	0.776	0.989	0.190	0.858	0.962	0.658	0.215
CVA	0.773	0.694	0.993	0.404	0.967	0.968	0.671	0.853
GKC+SA	0.790	0.662	0.980	0.383	0.694	0.861	0.752	0.928
FR	0.995	0.907	0.942	0.963	0.845	0.852	0.835	0.794

3.2.1 Simulation of change detection

We structured the image simulation process in two stages: classes and observations. The first defines a spatial distribution of elements/objects over two images, defining the pair “initial–posterior” phantoms. In the second, we simulate observations (spectral behavior) as Gaussian deviates. The choice of a Gaussian model as the basis in the simulation process is chosen in agreement with the spectral behavior observed in optical remote sensing imagery, as discussed by Schowengerdt (2006); Mather and Koch (2011), but can be replaced by any suitable model.

Let $\mathcal{S} \subset \mathbb{N}^2$ be the image support, and R'_j , $j = 1, \dots, n$, subsets of $\mathcal{S}$ such that $\bigcup_{j=1}^n R'_j = \mathcal{S}$. Each R'_j may be expressed as a geometric object in $\mathcal{S}$, composing then an initial phantom image $\mathcal{I}^{(1)}$. The change $\mathcal{I}^{(1)}$ to the posterior ($\mathcal{I}^{(2)}$) stage is modeled by geometric transformations (rotations and scale transformations) over the regions R'_j . We denote R''_j the region R'_j after the transformation. The actual change is the collection of differences between R'_j and R''_j , for every j .

We used circular, rectangular, and triangular regions, being their parameters the position over $\mathcal{S}$, the initial rotation, and the edges or radius size. These shapes and their parameters are randomly chosen as samples of uniformly distributed random variables. Furthermore, the changes from R'_j and R''_j are achieved multiplying the edges or radius of R'_j by a uniformly distributed scale factor in $[1/2, 3/2]$, allowing to express limited effects of increasing or decreasing its size.

Fig. 10 illustrates the step-by-step procedure for simulating a pair of images in which changes have occurred. Figs. 10(a) and 10(b) depict a typical example of initial-posterior pair of phantom images and Fig. 10(c) the resulting change/non-change map.

The spectral behavior of R'_j , $j = 1, \dots, n$, is simulated with basis on statistical parameters estimated from real targets observed in the image used in the previous section. For the sake of simplicity, let us denote $N(\mu_j, \Sigma_j)$ the p -variate Gaussian distribution that models region R'_j . Denote V_j the eigenvectors of Σ_j and V_j^{-1} its inverse; L_j is the diagonal matrix defined by the square root of the eigenvalues of Σ_j . In order to sample from this law, one starts from $\mathbf{N} = (N_1, \dots, N_p)$ a vector of p independent identically distributed standard Gaussian random variables, and computes $V_j^{-1} \mathbf{N} L_j + \mu_j$. Since we want to model a certain amount of variability in R'_j , the observations in R'_j will be samples from $V_j^{-1} \mathbf{N} L_j \xi_j + \mu_j \psi_j$, where ξ_j and ψ_j are independent random variables with uniform distribution in $[3/4, 5/4]$. Notice that ξ_j introduces variability in each observation variance, while ψ_j perturbs each sample's mean for region R'_j . Frery et al. (2009) and Negri et al. (2019) described the user uncertainty about the true nature of the scene with this heterogeneous model.

We adopted a similar approach for the simulation of the regions R''_j , $j = 1, \dots, n$ of the posterior image. The observations in R''_j are samples from $(V_j W)^{-1} \mathbf{N} L_j \xi_j + \mu_j \psi_j$, where W is a orthogonal matrix with $w_{pp} = 1$ and $w_{pp} > w_{pq}$, for $p, q = 1, \dots, \dim(\mu_j)$ and $p \neq q$, was adopted to insert a spectral distortion on the simulation results though changes on the correlation structure of V_j . Observe that if W is the identity matrix, then R'_j and R''_j have the same distribution. Furthermore, W is fixed in the simulation of all R''_j along with the posterior image.

The parameters μ_j and Σ_j are estimated from a set of observations $\mathcal{D}_j = \{\mathbf{x}_d \in \mathcal{X} : d = 1, \dots, n_j\}$ related to a land cover type ω_j in $\mathcal{I}$. Assuming the existence of t sets $\mathcal{D}_c$, $c = 1, \dots, t$, a (uniform) random choice is made to select one observation set to estimate the parameters that model the spectral behavior of each region R'_j and R''_j (i.e., both regions are simulated with basis on a same land cover class).

Fig. 11 depicts the spectral distortion trough

$$W = \begin{pmatrix} 1 & 0.393 & 0.279 \\ -0.085 & 1 & -0.965 \\ -0.360 & 0.855 & 1 \end{pmatrix}$$

using the same data set of Fig. 3. The top row shows the initial image and its 2-dimensional scatter plots. The bottom row shows the result of applying the spectral distortion on the data distribution. It can be noted that the W amounts to a linear transformation on the features.

We identified nine land covers in the Mariana's Dam image (Fig. 5(b)): two types of vegetation, four types of soil, two mixtures of rocks and vegetation, river and sandbank. We used these samples to estimate the parameters needed in the simulation process.

The initial phantom (Fig. 10(a)) is transformed into the initial image (Fig. 10(d)). The posterior phantom, i.e., the one that has changed, becomes an intermediate (unobserved) posterior image (Fig. 10(d)). Before being observed, the intermediate posterior image suffers from spectral distortions, yielding the image shown in Fig. 10(f), which is the observed data. These three sources of change (spatial change, spectral change, and spectral distortion) comprise our model for the process that transforms the data along observation epochs.

We used models inferred from Mariana’s Dam image to sample the observations in Figs. 10(d) and 10(e).

Using the above-discussed method and parameters estimated from remotely sensed data (i.e., the Mariana’s Dam image of 2016 and land cover samples), we simulated 50 pairs of spectrally normalized images, as well as 50 pairs of non-normalized images. These images have 500×500 pixels.

3.2.2 Change detection on simulated datasets

Similarly to Section 3.1, we applied the analyzed methods to each pair of simulated images, and then assessed the results in terms of F_1 -Score, Precision, and Recall using the change/non-change map as reference truth. We also computed the computational run-times. We used fixed parameters in this experiment as follows: for ASF was adopted $\lambda = 6$; for PCA+K-Means, both ρ and θ equal to 2; the threshold was the Kittler-Illingworth method in CVA; the cooling parameter α was set as 0.82 in GKC+SA; finally, ρ and β were fixed as 1 and 2.5 % in FR method.

Fig. 12 presents the notched boxplots of the three figures of merit (F_1 , Precision, and Recall) observed over 50 independent replications. The results describe two situations: with and without spectral distortion. The F_1 -Score of FR is, in both situations, better than those produced by the competitors: its median is higher, with little to no overlap of the notches. The Precision of FR is, in median, also better than the other procedures. Notice that under the presence of spectral distortion, FR’s Precision is indisputably the best. When it comes to Recall, there is no clear winner.

To check the significance of results, we computed bilateral Wilcoxon tests between the F_1 -Scores of each pair of methods; Table 3 shows the p -values of the null hypothesis that they are equal. These results verify that the proposed technique is statistically different from all the other methods, regardless of the considered dataset. Concerning the other methods, two pairs of techniques are equivalent when dealing with non-normalized data sets (p -values above the diagonal): CVA and ASF, and GKC+SA. When the data sets are normalized (p -values above the diagonal), the equivalence is between GKC+SA and ASF.

Table 3 Two-sided Wilcoxon test results. Above diagonal are the comparisons of non-normalized datasets, and below it are the results of spectral normalized simulated datasets.

	ASF	PCA+K-Means	CVA	GKC+SA	FR
ASF		0.0184	0.7407	0.0006	0.0264
PCA+K-Means	0		0.0009	0.3520	0
CVA	0.0002	0.0899		0	0.0393
GKC+SA	0.8740	0.0007	0.0143		0
FR	0	0.0020	0	0	

Figs. 13 and 14 show the change detection map produced by each method with the simulated examples shown in Fig. 10. It can be observed that PCA+K-Means and FR provide outputs closer to the expected result. On the other hand, ASF and GKC+SA return completely inaccurate change maps.

Fig. 15 shows the notched boxplots of the times, in seconds, required to run the simulations. We present these results in a logarithmic scale since some of the medians differ orders of magnitude. On the one hand, GKC+SA and FR are the most expensive methods; on the other hand, the fastest methods are CVA, ASF, and PCA+K-Means. Additionally, the spectral distortion does not lead to relevant changes in computational times of any method.

4 Conclusions

We proposed a novel method for unsupervised change detection based on clustering and optimization. As part of this proposal, we performed experiments with remotely sensed imagery and simulated data to prove its effectiveness and compare it with well-known methods.

One of our proposal's significant strengths is the use of floating references, which are automatically detected and, thus, require no user intervention. Our proposal is less sensitive to temporal radiometric normalization than the other techniques. Also, the proposed method is less sensitive to parameter tuning in comparison to other methods face to different input data (*vide* Fig. 1). Furthermore, our method is better at preserving the geometric structure of objects providing, therefore, more realistic change maps; (*vide* Figs. 8 and 9).

The computational time is the price to pay for the proposed method. However, this weakness is minimized since the temporal radiometric normalization is no longer needed.

As a perspective for future works, we intend to: (i) eliminate the *minimum cluster size* parameter with the use of a new initial clustering; (ii) exploit other similarity measures, in particular, stochastic and geodesic distances; (iii) adapt the method in order to allow temporal comparisons with images acquired by different sensors; (iv) develop a computational strategy to reduce the computational cost of the proposed method; and (v) extend the proposed simulation framework in order to allow other statistical distributions beyond the Gaussian.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Adler-Golden SM, Matthew MW, Bernstein LS, Levine RY, Berk A, Richtsmeier SC, Acharya PK, Anderson GP, Felde JW, Gardner JA, Hoke ML, Jeong LS, Pukall B, Ratkowski AJ, Burke HhK (1999) Atmospheric correction for shortwave spectral imagery based on MODTRAN4. In: Imaging Spectrometry V, vol 3753, pp 3753 – 3753 – 9, DOI 10.1117/12.366315, URL <https://doi.org/10.1117/12.366315>

- do Carmo FF, Kamino LHY, Junior RT, de Campos IC, do Carmo FF, Silvino G, da Silva Xavier de Castro KJ, Mauro ML, Rodrigues NUA, de Souza Miranda MP, Pinto CEF (2017) Fundão tailings dam failures: the environment tragedy of the largest technological disaster of brazilian mining in global context. *Perspectives in Ecology and Conservation* 15(3):145 – 151, DOI <https://doi.org/10.1016/j.pecon.2017.06.002>, URL <http://www.sciencedirect.com/science/article/pii/S1679007316301566>
- Celik T (2009) Unsupervised change detection in satellite images using principal component analysis and *k*-means clustering. *IEEE Geoscience and Remote Sensing Letters* 6(4):772–776, DOI 10.1109/LGRS.2009.2025059
- Chacón JE (2020) The modal age of statistics. *International Statistical Review* DOI 10.1111/insr.12340
- Chernoff H (1964) Estimation of the mode. *Annals of the Institute of Statistical Mathematics* 16(1):31–41, DOI 10.1007/bf02868560
- Colwell J, Weber F (1981) Forest change detection. In: *Fifteenth International Symposium on Remote Sensing of Environment*, ERIM, Ann Arbor, pp 839–852.
- Crist EP, Cicone RC (1984) A physically-based transformation of thematic mapper data—the tm tasseled cap. *IEEE Transactions on Geoscience and Remote Sensing* GE-22(3):256–263, DOI 10.1109/TGRS.1984.350619
- Frery AC, Ferrero S, Bustos OH (2009) The influence of training errors, context and number of bands in the accuracy of image classification. *International Journal of Remote Sensing* 30(6):1425–1440, DOI 10.1080/01431160802448919
- Frery AC, Gomez L, Medeiros AC (2020) A badging system for reproducibility and replicability in remote sensing research. *IEEE Journal of Selected Topics on Applied Earth Observations and Remote Sensing* 13:4988–4995, DOI 10.1109/JSTARS.2020.3019418
- Ghosh A, Mishra NS, Ghosh S (2011) Fuzzy clustering algorithms for unsupervised change detection in remote sensing images. *Information Sciences* 181(4):699 – 715, DOI <https://doi.org/10.1016/j.ins.2010.10.016>
- Gustafson DE, Kessel WC (1978) Fuzzy clustering with a fuzzy covariance matrix. In: *1978 IEEE Conference on Decision and Control including the 17th Symposium on Adaptive Processes*, pp 761–766, DOI 10.1109/CDC.1978.268028
- Johnson RD, Kasischke ES (1998) Change vector analysis: A technique for the multispectral monitoring of land cover and condition. *International Journal of Remote Sensing* 19(3):411–426, DOI 10.1080/014311698216062
- Kirkpatrick S, Gelatt CD, Vecchi MP (1983) Optimization by simulated annealing. *Science* 220(4598):671–680, DOI 10.1126/science.220.4598.671, URL <http://science.sciencemag.org/content/220/4598/671>
- Kittler J, Illingworth J (1986) Minimum error thresholding. *Pattern Recognition* 19(1):41 – 47, DOI [https://doi.org/10.1016/0031-3203\(86\)90030-0](https://doi.org/10.1016/0031-3203(86)90030-0)
- Liu S, Bruzzone L, Bovolo F, Du P (2015) Hierarchical Unsupervised Change Detection in Multitemporal Hyperspectral Images. *IEEE Transactions on Geoscience and Remote Sensing* 53(1):244–260, DOI 10.1109/TGRS.2014.2321277
- Lu D, Mausel P, Brondízio E, Moran E (2004) Change detection techniques. *International Journal of Remote Sensing* 25(12):2365–2401, DOI 10.1080/0143116031000139863

- Mather PM, Koch M (2011) *Computer Processing of Remotely-Sensed Images: An Introduction*, 3rd edn. Wiley
- Mou L, Zhu XX (2020) Learning to pay attention on spectral domain: A spectral attention module-based convolutional network for hyperspectral image classification. *IEEE Transactions on Geoscience and Remote Sensing* 58(1):110–122, DOI 10.1109/tgrs.2019.2933609
- Mura MD, Benediktsson JA, Bovolo F, Bruzzone L (2008) An unsupervised technique based on morphological filters for change detection in very high resolution images. *IEEE Geoscience and Remote Sensing Letters* 5(3):433–437, DOI 10.1109/LGRS.2008.917726
- Negri RG, Frery AC, Silva WB, Mendes TSG, Dutra LV (2019) Region-based classification of PolSAR data using radial basis kernel functions with stochastic distances. *International Journal of Digital Earth* 12(6):699–719, DOI 10.1080/17538947.2018.1474958
- Otsu N (1979) A threshold selection method from gray-level histograms. *IEEE Trans on Systems, Man, and Cybernetics* 9(1):62–66, DOI 10.1109/TSMC.1979.4310076
- Rijsbergen CJV (1979) *Information Retrieval*, 2nd edn. Butterworth-Heinemann, USA
- Ross S, Bhadauria HS (2015) A Review of Change Detection Techniques of Land-Cover Using Remote Sensing Data. *IOSR Journal of Computer Engineering Ver IV* 17(3):2278–661, DOI 10.9790/0661-17341721
- Schowengerdt RA (2006) *Remote Sensing, Third Edition: Models and Methods for Image Processing*. Academic Press, Inc., Orlando, FL, USA
- Silverman BW (1986) *Density Estimation for Statistics and Data Analysis*. Chapman & Hall, London
- Srivastava PK, Han D, Rico-Ramirez MA, Bray M, Islam T (2012) Selection of classification techniques for land use/land cover change investigation. *Advances in Space Research* 50(9):1250–1265, DOI 10.1016/j.asr.2012.06.032
- Theodoridis S, Koutroumbas K (2008) *Pattern Recognition*, 4th edn. Academic Press, San Diego
- Volpi M, Tuia D, Camps-Valls G, Kanevski M (2012) Unsupervised change detection with kernels. *IEEE Geoscience and Remote Sensing Letters* 9(6):1026–1030, DOI 10.1109/LGRS.2012.2189092
- Wu C, Du B, Zhang L (2014) Slow Feature Analysis for Change Detection in Multispectral Imagery. *IEEE Transactions on Geoscience and Remote Sensing* 52(5):2858–2874, DOI 10.1109/TGRS.2013.2266673
- Xie XL, Beni G (1991) A validity measure for fuzzy clustering. *IEEE Transactions on Pattern Analysis and Machine Intelligence* 13(8):841–847, DOI 10.1109/34.85677

(a) September 25th 2015

(b) and August 10th 2016

(c) Ground truth change/non-change areas

Fig. 5 Mariana's Dam dataset: (a) and (b) represented in natural color composition; (c) change and non-change areas denoted in green and red, respectively.

(a) August 22th 1988

(b) August 2nd 1999

(c) Ground truth change/non-change areas

Fig. 6 Belterra deforestation dataset: (a) and (b) represented in natural color composition; (c) change and non-change areas denoted in green and red, respectively.

Fig. 7 F_1 -Scores obtained for each parameter configuration presented in Table 1. Box-and-whisker plots, with whiskers representing the minimum-maximum accuracy range, notches express the 90 % confidence range around the median, and black dots the individual accuracy values.

Fig. 8 Best change/non-change maps obtained by each analyzed method for Mariana's Dam dataset. Changes and non-changes are in green and red, respectively.

Fig. 9 Best change/non-change maps obtained by each analyzed method for Belterra Deforestation dataset. Changes and non-changes are in green and red, respectively.

Fig. 10 Step-by-step procedure for image simulation.

Fig. 11 Spectral differences between $\mathcal{I}^{(1)}$ and $\mathcal{I}^{(2)}$ caused by W on the feature components (bands) x_1 , x_2 , and x_3 .

Fig. 12 F_1 -Scores, Precision and Recall values regarding the experiments with simulated images.

Fig. 13 Example of result for simulated data without spectral distortion.

Fig. 14 Example of result for simulated data with spectral distortion.

Fig. 15 Run-time on experiments with simulated datasets.