Unusual Gold(I)-Catalyzed Isomerisation of 3-hydroxylated 1,5-enynes : Highly Substrate Dependant Reaction Manifolds

Fabien Gagosz*

Département de Chimie et de Synthèse Organique, Ecole Polytechnique, 91128 Palaiseau, France

Supporting Information

General Information.

All reactions were carried out under an atmosphere of dry argon. Commercial reagents were used as received without further purification. $(\text{Ph}_3)\text{PAuCl}$ was synthesized following a literature procedure.¹ All products were purified by flash column chromatography using silica gel SDS 60 C.C. 40-63 or neutral alumina SDS C.C. 50-200. NMR spectra were recorded in CDCl_3 with TMS as an internal standard at ambient temperature on a Bruker Avance 400 operating at 400 MHz for ^1H and 100 MHz for ^{13}C . Infrared Absorption spectra were recorded as a solution in CCl_4 with a Perkin-Elmer 1600 Fourier Transform Spectrophotometer. Mass spectra were recorded with a HP 5989B mass spectrometer. Melting points were determined by Reichert microscope apparatus and were uncorrected.

Experimental Section.

I- synthesis of the substrates

General procedure for the synthesis of 3-hydroxylated 1,5-enynes

Enyne substrates were synthesized according to the general following sequence:

(1) Bruce, M. I. ; Nicholson, B. K. ; Shawkataly, O. B. *Inorg. Synth.* **1989**, 26, 324-330.

(E)-2,4-Dimethyl-7-phenyl-hept-4-en-3-one (5)

To a stirred solution of dry diisopropylamine (3.36 ml, 24 mmol) in 40ml of THF under Argon at -10°C was added dropwise a solution of n-BuLi (1.35M in hexanes, 16.3 ml, 22 mmol) and the reaction mixture was stirred for 30 min. The mixture was then cooled to -78°C and 2-methyl pentan-3-one (2.47 ml, 20 mmol) was added dropwise. After stirring for 45 min at -78°C , hydrocinnamaldehyde (3.15 ml, 24 mmol) was added dropwise and the reaction mixture was stirred for a further 45 min. The reaction mixture was diluted with 40ml of dry THF and acetic anhydride (3.77 ml, 40 mmol) was then added dropwise at -78°C followed by DMAP (2.44 g, 20 mmol) and the mixture was slowly cooled to room temperature. Water was then added and the mixture was extracted with ether. The combined organic layer was washed with aqueous 1.0M HCl, dried over magnesium sulfate, filtered, and the solvents were removed under reduced pressure. The crude resulting oil was dissolved in 40ml of dry THF at RT under argon and DBU (4.48 ml, 30 mmol) was added. The mixture was stirred for 12h after which time the reaction was complete. Aqueous 1.0M HCl was added and the mixture was extracted with ether. The separated organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography (silica gel, Petroleum Ether-Ether, 97.5:2.5) to give **5** (3.51 g, 81%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.29 (m, 2H), 7.24-7.20 (m, 3H), 6.65 (tq, $J = 7.2, 1.2$ Hz, 1H), 3.28 (sept, $J = 6.8$, 1H), 2.81 (t, $J = 7.7$ Hz, 2H), 2.58 (td, $J = 7.7, 7.1$ Hz, 2H), 1.75 (bs, 3H), 1.08 (d, $J = 6.8$, 6H). ^{13}C NMR (50 MHz, CDCl_3) δ 206.2, 141.1, 140.5, 136.6, 128.6, 128.4, 126.2, 34.9, 33.8, 30.9, 19.7, 11.7. IR (CCl_4) 2970, 2931, 1669, 1639, 1467, 1454, 1043 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 234 (MNH_4^+), 217 (MH^+). Anal. calcd. for $\text{C}_{15}\text{H}_{20}\text{O}$: C, 83.28; H, 9.32, found: C, 83.33; H, 9.47.

(E)-2-Methyl-7-phenyl-hept-4-en-3-one (6)

To a stirred solution of dry diisopropylamine (3.36 ml, 24 mmol) in 40ml of THF under Argon at -10°C was added dropwise a solution of n-BuLi (1.35M in hexanes, 16.3 ml, 22 mmol) and the reaction mixture was stirred for 30 min. The mixture was then cooled to -78°C and 3-methyl butan-2-one (2.14 ml, 20 mmol) was added dropwise. After stirring for 45 min

at -78°C, hydrocinnamaldehyde (3.15 ml, 24 mmol) was added dropwise and the reaction mixture was stirred for a further 45 min. The reaction mixture was diluted with 40ml of dry THF and acetic anhydride (3.77 ml, 40 mmol) was then added dropwise at -78°C followed by DMAP (2.44 g, 20 mmol) and the mixture was slowly cooled to room temperature. Water was then added and the mixture was extracted with ether. The combined organic layer was washed with aqueous 1.0M HCl, dried over magnesium sulfate, filtered, and the solvents were removed under reduced pressure. The crude resulting oil was dissolved in 40ml of dry THF at RT under argon and DBU (4.48 ml, 30 mmol) was added. The mixture was stirred for 12h after which time the reaction was complete. Aqueous 1.0M HCl was added and the mixture was extracted with ether. The separated organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography (silica gel, Petroleum Ether-Ether, 97.5:2.5) to give **6** (3.19 g, 79%) as a colorless oil. $^1\text{H NMR}^2$ (400 MHz, CDCl_3) δ 7.32-7.28 (m, 2H), 7.24-7.18 (m, 3H), 6.90 (dt, $J = 15.7, 6.9$ Hz, 1H), 6.17 (dt, $J = 15.7, 1.5$ Hz, 1H), 2.77-2.84 (m, 3H), 2.52-2.58 (m, 2H), 1.09 (d, $J = 6.9$, 6H).

(E)-2-Methyl-5-phenyl-pent-2-enal (7)

A mixture of hydrocinnamaldehyde (0.880 ml, 6 mmol) and (α -formylethylidene) triphenylphosphorane³ (2.32 g, 7.3 mmol) in dry toluene was heated at 80°C under argon with stirring for 13 h. The solution was then allowed to cool to room temperature, the toluene was removed in vacuo, and the concentrated material was diluted with ether. The solid was removed by filtration, and evaporation of the filtrate gave a crude product which was purified by flash column chromatography (silica gel, Petroleum Ether-Ether, 97.5:2.5) to give **7** (0.94 g, 90%) as a colorless oil. $^1\text{H NMR}^4$ (400 MHz, CDCl_3) δ 9.39 (s, 1H), 7.29-7.34 (m, 2H), 7.20-7.25 (m, 3H), 6.52 (tq, $J = 7.2, 1.3$ Hz, 1H), 2.84 (t, $J = 7.5$ Hz, 2H), 2.69 (td, $J = 7.5, 7.2$ Hz, 2H), 1.70 (bs, 3H).

(2) Nakamura, S.; Hayakawa, T; Nishi, T.; Watanabe, Y.; Toru, T. *Tetrahedron* **2001**, 57, 6703-6712

(3) Kiyooka, S-I.; Hena, M. A. *J. Org. Chem.* 1999, 64, 5511-5523.

(4) Hon, Y.-S.; Lu, L.; Chang, R.-C.; Lin, S.-W.; Sun, P.-P.; Lee, C.-F. *Tetrahedron* **2000**, 56, 9269-9280.

General procedure for the synthesis of 3-hydroxylated 1,5-enynes from **5,6** and **7**

To a stirred solution of the α,β -unsaturated carbonyl compound **5,6** or **7** (1 mmol) in 4ml of dry THF at -78°C under argon was added diiodomethane (2 mmol). A solution of n-BuLi (1.6M in hexanes, 1.75 mmol) was then added dropwise and the reaction mixture was stirred for 1h at -78°C . Saturated aqueous ammonium chloride was added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. Excess of diiodomethane was removed from the crude highly labile epoxide by quick filtration of the residue through a small column of neutral alumina deactivated with 7% of water (petroleum ether to petroleum ether-ether 99:1). The crude epoxide was then diluted with 7ml of dry ether under argon and magnesium bromide ethyl etherate (1 mmol) was added. The reaction mixture was stirred at RT for 45 min. Saturated aqueous ammonium chloride was then added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. A solution of the resulting crude aldehyde in 3 ml of dry THF was then added to a preformed solution of the lithium acetylide anion of the corresponding alkyne (3mmol) at -78°C and the mixture was slowly cooled to room temperature. After stirring at RT for a further 30 min, saturated aqueous ammonium chloride was added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography to give the corresponding 3-hydroxylated 1,5-enyne.

(E)-5-Isopropyl-4-methyl-1-phenyl-tridec-3-en-7-yn-6-ol

Isolated as an inseparable 6:1 mixture of *syn* and *anti* isomers. Yield 70%. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.31 (m, 2H), 7.18-7.21 (m, 3H), 5.34 (tq, $J = 7.0, 0.9$ Hz, 1H), 4.42-4.47 (m, 1H), 2.66-2.77 (m, 2H), 2.43 (q, $J = 7.4$, 2H), 2.19 (td, $J = 7.1, 2.1$ Hz, 2H), 1.84-1.96 (m, 3H), 1.66 (bs, 3H), 1.45-1.54 (m, 2H), 1.27-1.40 (m, 4H), 0.95 (d, $J = 6.2$, 3H), 0.91 (t, $J = 7.1$, 3H), 0.80 (d, $J = 6.2$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.9, 134.2, 131.3, 128.5, 128.4, 126.0, 86.4, 79.8, 61.9, 61.5, 35.8, 31.2, 29.7, 28.4, 27.4, 22.3, 21.8, 20.6, 18.8, 14.2, 14.1. IR (CCl_4) 3542, 2958, 2931, 2871, 1603, 1495, 1453, 1030 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 344 (MNH_4^+), 309, 253.

(E)-4-Isopropyl-5-methyl-1,8-diphenyl-oct-5-en-1-yn-3-ol

Isolated as an inseparable 5:1 mixture of *syn* and *anti* isomers. Yield 73%. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.39-7.42 (m, 2H), 7.28-7.33 (m, 5H), 7.20-7.22 (m, 3H), 5.41 (tq, $J = 7.1, 1.1$ Hz, 1H), 4.68 (dd, $J = 9.9, 5.2$ Hz, 1H), 2.74 (td, $J = 7.2, 3.3$ Hz, 2H), 2.47 (td, $J = 7.2, 7.1$ Hz, 2H), 1.96-2.08 (m, 3H), 1.73 (bs, 3H), 1.02 (d, $J = 6.1$, 3H), 0.84 (d, $J = 6.1$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.8, 134.0, 131.8, 131.6, 128.5 (x2), 128.3 (x2), 126.0, 123.1, 89.2, 85.7, 62.1, 61.9, 35.8, 29.7, 27.5, 21.8, 20.7, 14.2. IR (CCl_4) 3536, 2959, 2927, 2870, 1589, 1489, 1453, 1385, 1029 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 350 (MNH_4^+), 332, 315, 259.

(E)-4-Isopropyl-5-methyl-8-phenyl-1-trimethylsilanyl-oct-5-en-1-yn-3-ol

Isolated as an inseparable 5:1 mixture of *syn* and *anti* isomers. Yield 68%. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.32 (m, 2H), 7.18-7.22 (m, 3H), 5.35 (tq, $J = 7.1, 1.0$ Hz, 1H), 4.42 (dd, $J = 9.9, 5.5$ Hz, 1H), 2.71 (td, $J = 7.2, 2.1$ Hz, 2H), 2.43 (td, $J = 7.2, 7.1$ Hz,

2H), 1.85-1.98 (m, 3H), 1.65 (bs, 3H), 0.96 (d, $J = 6.2$, 3H), 0.81 (d, $J = 6.4$, 3H), 0.16 (s, 9H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.9, 134.0, 131.6, 128.5 (x2), 126.0, 105.8, 90.5, 61.8, 61.7, 35.8, 29.7, 27.3, 21.8, 20.6, 14.1, -0.1. IR (CCl_4) 3537, 2960, 2870, 2167, 1603, 1496, 1453, 1385, 1250, 1030 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 346 (MNH_4^+), 311.

(E)-5-Isopropyl-1-phenyl-tridec-3-en-7-yn-6-ol

Isolated as a separable 5:1 mixture of *syn* and *anti* isomers. Yield 83%. *syn* isomer **1d** ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.32 (m, 2H), 7.18-7.22 (m, 3H), 5.56 (dt, $J = 15.3$, 6.7 Hz, 1H), 5.31 (dd, $J = 15.3$, 9.7 Hz, 1H), 4.36-4.40 (m, 1H), 2.69-2.82 (m, 2H), 2.38-2.51 (m, 2H), 2.20 (td, $J = 7.0$, 2.0 Hz, 2H), 1.84 (td, $J = 7.0$, 2.0 Hz, 1H), 1.77 (d, $J = 10.3$ Hz, 1H), 1.65-1.74 (m, 1H), 1.48-1.55 (m, 2H), 1.30-1.42 (m, 4H), 0.93 (t, $J = 7.0$, 3H), 0.92 (d, $J = 6.6$, 3H), 0.84 (d, $J = 6.7$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.6, 135.3, 129.2, 128.5, 128.4, 125.9, 86.4, 79.0, 62.9, 57.1, 35.8, 34.5, 31.0, 29.0, 28.5, 22.2, 21.2, 20.6, 18.7, 14.1. IR (CCl_4) 3549, 2958, 2932, 2871, 1496, 1454, 1383, 1366, 1026 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 330 (MNH_4^+), 312 (MH^+), 295. *anti* isomer **1f** ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.31 (m, 2H), 7.17-7.21 (m, 3H), 5.59 (dt, $J = 15.0$, 6.8 Hz, 1H), 5.22 (dd, $J = 15.3$, 9.7 Hz, 1H), 4.19-4.22 (m, 1H), 2.66-2.81 (m, 2H), 2.36-2.50 (m, 2H), 2.22 (td, $J = 7.1$, 1.9 Hz, 2H), 2.01-2.09 (m, 1H), 1.93-1.99 (m, 1H), 1.60 (d, $J = 3.8$ Hz, 1H), 1.48-1.55 (m, 2H), 1.27-1.40 (m, 4H), 0.90 (t, $J = 7.1$, 3H), 0.86 (d, $J = 6.8$, 3H), 0.78 (d, $J = 6.8$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.7, 135.6, 128.6, 128.4, 127.3, 125.9, 86.5, 79.8, 64.0, 56.6, 35.9, 34.7, 31.1, 28.4, 28.3, 22.3, 21.8, 18.8, 17.7, 14.1. IR (CCl_4) 3619, 3027, 2958, 2931, 2871, 1603, 1496, 1454, 1385, 1030 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 330 (MNH_4^+), 312 (MH^+), 295.

(E)-4-Methyl-1-phenyl-tridec-3-en-7-yn-6-ol

Yield 51%. ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.32 (m, 2H), 7.19-7.22 (m, 3H), 5.33 (tq, $J = 7.2$, 1.1 Hz, 1H), 4.39-4.40 (m, 1H), 2.63-2.75 (m, 2H), 2.32-2.41 (m, 4H), 2.21 (td, $J = 7.2$, 1.9 Hz, 2H), 1.81 (d, $J = 4.2$, 2H), 1.61 (bs, 3H), 1.48-1.55 (m, 2H), 1.28-1.41 (m, 4H), 0.92

(t, $J = 7.1$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.0, 131.4, 128.6, 128.5, 128.4, 125.9, 85.5, 80.9, 60.4, 48.5, 35.8, 31.1, 30.0, 28.4, 22.2, 18.7, 16.3, 14.0. IR (CCl_4) 3617, 2957, 2931, 2859, 1603, 1495, 1453, 1379 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 302 (MNH_4^+), 284 (MH^+), 284, 267.

(E)-4-Isopropyl-5-methyl-8-phenyl-oct-5-en-1-yn-3-ol

To a solution of (E)-4-Isopropyl-5-methyl-8-phenyl-1-trimethylsilanyl-oct-5-en-1-yn-3-ol (200 mg, 0.61 mmol) in 4 ml of DMSO at RT was added potassium fluoride (350 mg, 6.1 mmol) and a few drops of water. After stirring for 2h at RT, saturated aqueous ammonium chloride was added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography (silica gel, Petroleum Ether-Ether, 95: 5) to give **1k** (150 mg, 96%) as a colorless oil. Isolated as an inseparable 5:1 mixture of *syn* and *anti* isomers. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.32 (m, 2H), 7.18-7.22 (m, 3H), 5.36 (tq, $J = 7.1$, 1.1 Hz, 1H), 4.46 (ddd, $J = 9.8$, 5.6, 2.3 Hz, 1H), 2.72 (td, $J = 7.3$, 4.1 Hz, 2H), 2.44 (td, $J = 7.3$, 7.1 Hz, 2H), 2.41 (d, $J = 2.3$ Hz, 1H), 1.85-2.02 (m, 3H), 1.67 (bs, 3H), 0.97 (d, $J = 6.3$, 3H), 0.81 (d, $J = 6.3$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 141.8, 133.9, 131.8, 128.5, 128.4, 126.0, 83.7, 74.0, 61.5, 61.3, 35.7, 29.6, 27.3, 21.7, 20.6, 14.1. IR (CCl_4) 3536, 3309, 2960, 2927, 2870, 1495, 1453, 1386, 1046, 1026 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 274 (MNH_4^+), 257 (MH^+).

General procedure for the benzylation of 3-hydroxylated 1,5-enynes

To a solution of the alcohol (1 mmol) in 3 ml of dry THF under argon at RT was added NaH (60% in mineral oil, 1.2 mmol) and the solution was stirred until hydrogen evolution has stopped. Benzyl bromide (1.4 mmol) was then added followed by a catalytic amount of tetrabutylammonium iodide (0.1 mmol). The reaction mixture was stirred until full consumption of the starting material (as attested by TLC analysis, 3 to 12h). Saturated aqueous ammonium chloride was then added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography (silica gel, Petroleum Ether-Ether) to give the corresponding benzyl ether as a colorless oil.

Isolated as an inseparable 6:1 mixture of *syn* and *anti* isomers. Yield 95%. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.39 (m, 4H), 7.27-7.30 (m, 3H), 7.17-7.23 (m, 3H), 5.28 (bt, $J = 7.0$ Hz, 1H), 4.82 (d, $J = 12.0$ Hz, 1H), 4.52 (d, $J = 12.0$ Hz, 1H), 4.25 (dt, $J = 6.1, 1.7$ Hz, 1H), 2.69 (td, $J = 7.5, 2.1$ Hz, 2H), 2.39 (td, $J = 7.5, 7.0$ Hz, 2H), 2.24 (td, $J = 6.9, 1.91$ Hz, 2H), 1.98-2.11 (m, 2H), 1.68 (bs, 3H), 1.49-1.56 (m, 2H), 1.30-1.45 (m, 4H), 0.93 (d, $J = 6.8$, 3H), 0.91 (t, $J = 7.0$, 3H), 0.78 (d, $J = 6.4$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.6, 138.8, 134.7, 128.5, 128.3 (x2), 127.7, 127.6, 127.4, 125.7, 87.5, 78.4, 70.5, 70.3, 59.4, 36.1, 31.2, 29.9, 28.5, 27.8, 22.3, 21.5, 20.7, 18.9, 15.7, 14.1. IR (CCl_4) 2957, 2931, 2869, 1604, 1496, 1453, 1090, 1069 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 435 (MNH_4^+), 307.

Isolated as an inseparable 5:1 mixture of *syn* and *anti* isomers. Yield 95%. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.16-7.45 (m, 15H), 5.36 (tq, $J = 7.0, 1.0$ Hz, 1H), 4.90 (d, $J = 12.0$ Hz, 1H), 4.61 (d, $J = 12.0$ Hz, 1H), 4.49 (d, $J = 12.0$ Hz, 1H), 2.69 (td, $J = 7.6, 2.0$ Hz, 2H), 2.41 (td, $J = 7.6, 7.0$ Hz, 2H), 2.22 (dd, $J = 8.7, 6.8$ Hz, 1H), 2.05-2.15 (m, 1H), 1.74 (bs, 3H), 1.00 (d, $J = 6.6$, 3H), 0.82 (d, $J = 6.7$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.5, 138.5, 134.6, 131.7, 128.6, 128.4, 128.3 (x2), 128.2, 127.9, 127.7, 127.5, 125.7, 123.3, 88.1, 87.0, 70.9, 70.7, 59.6, 36.1, 29.9, 27.9, 21.5, 20.8, 15.7. IR (CCl_4) 2959, 2933, 2865, 1490, 1451, 1070, 1005 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 441 (MNH_4^+), 313, 257.

Yield 98%. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.39 (m, 4H), 7.27-7.31 (m, 3H), 7.17-7.22 (m, 3H), 5.55 (dt, $J = 15.2, 6.6$ Hz, 1H), 5.38 (dd, $J = 15.2, 9.5$ Hz, 1H), 4.81 (d, $J = 12.0$ Hz, 1H), 4.53 (d, $J = 12.0$ Hz, 1H), 4.12 (dt, $J = 6.7, 1.9$ Hz, 1H), 2.67-2.79 (m, 2H), 2.34-2.47 (m, 2H), 2.26 (td, $J = 7.0, 1.9$ Hz, 2H), 2.10 (td, $J = 9.4, 6.4$ Hz, 1H), 1.89-2.01 (m, 1H), 1.51-

1.58 (m, 2H), 1.30-1.46 (m, 4H), 0.93 (t, $J = 7.1$, 3H), 0.85 (d, $J = 6.8$, 3H), 0.76 (d, $J = 6.8$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.3, 138.6, 133.1, 128.6, 128.3 (x2), 128.1, 127.9, 127.5, 125.8, 87.8, 78.1, 70.6, 70.2, 54.3, 36.3, 34.8, 31.2, 28.6, 27.9, 22.3, 21.4, 18.9, 18.7, 14.2. IR (CCl_4) 2957, 2930, 2870, 1496, 1454, 1090, 1069 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 421 (MNH_4^+), 293.

Yield 94%. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.39 (m, 4H), 7.27-7.31 (m, 3H), 7.17-7.23 (m, 3H), 5.53 (dt, $J = 15.2$, 6.6 Hz, 1H), 5.37 (dd, $J = 15.2$, 9.1 Hz, 1H), 4.81 (d, $J = 12.0$ Hz, 1H), 4.50 (d, $J = 12.0$ Hz, 1H), 4.15 (dt, $J = 5.7$, 1.9 Hz, 1H), 2.68-2.79 (m, 2H), 2.35-2.48 (m, 2H), 2.27 (td, $J = 7.0$, 1.9 Hz, 2H), 1.98-2.06 (m, 2H), 1.52-1.59 (m, 2H), 1.317-1.46 (m, 4H), 0.94 (t, $J = 7.1$, 3H), 0.85 (d, $J = 6.4$, 3H), 0.78 (d, $J = 6.5$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.3, 138.6, 132.9, 128.6, 128.5, 128.3, 128.2, 127.9, 127.5, 125.8, 87.6, 78.6, 71.1, 70.2, 54.7, 36.3, 34.8, 31.1, 28.6, 28.4, 22.3, 21.5, 19.1, 18.8, 14.2. IR (CCl_4) 2957, 2931, 2870, 1496, 1454, 1090, 1069 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 421 (MNH_4^+), 293.

Yield 99%. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.40 (m, 4H), 7.27-7.31 (m, 3H), 7.18-7.22 (m, 3H), 5.34 (tq, $J = 6.7$, 0.9 Hz, 1H), 4.81 (d, $J = 11.9$ Hz, 1H), 4.53 (d, $J = 11.9$ Hz, 1H), 4.19 (tt, $J = 6.9$, 1.8 Hz, 1H), 2.67 (t, $J = 7.5$ Hz, 2H), 2.49 (dd, $J = 13.6$, 7.2 Hz, 1H), 2.38 (dd, $J = 13.6$, 6.5 Hz, 1H), 2.35 (td, $J = 7.5$, 6.7 Hz, 2H), 2.26 (td, $J = 7.0$, 1.8 Hz, 2H), 1.60 (bs, 3H), 1.52-1.57 (m, 2H), 1.32-1.45 (m, 4H), 0.94 (t, $J = 7.1$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.4, 138.4, 131.8, 128.5, 128.4, 128.3, 128.0, 127.6, 127.1, 125.8, 86.9, 79.2, 70.2, 68.2, 46.2, 36.1, 31.2, 30.2, 28.6, 22.3, 18.8, 16.5, 14.1. IR (CCl_4) 3064, 3028, 2931, 2859, 1496, 1453, 1336, 1094, 1071 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 392 (MNH_4^+), 267.

Synthesis of acetate **1c** from **1a**

To a solution of **1a** (200 mg, 0.61 mmol) in 2 ml of dry dichloromethane at RT under argon were sequentially added triethylamine (130 μ l, 0.92 mmol), acetic anhydride (76 μ l, 0.80 mmol) and a catalytic amount of DMAP (7 mg, 0.06 mmol). After stirring for 1h at RT, saturated aqueous ammonium chloride was added and the mixture was extracted with ether. The combined organic layer was dried over magnesium sulfate, filtered and the solvents were removed under reduced pressure. The residue was purified by flash column chromatography (silica gel, Petroleum Ether-Ether, 95: 5) to give **1c** (216 mg, 96%) as a colorless oil.

Isolated as an inseparable 6:1 mixture of *syn* and *anti* isomers. *syn* isomer ^1H NMR (400 MHz, CDCl_3) δ 7.26-7.30 (m, 2H), 7.17-7.21 (m, 3H), 5.60 (dt, J = 6.4, 2.1 Hz, 1H), 5.20 (tq, J = 6.9, 0.8 Hz, 1H), 2.66 (t, J = 7.8 Hz, 2H), 2.36 (td, J = 7.8, 6.9 Hz, 2H), 2.21 (td, J = 7.0, 2.1 Hz, 2H), 2.09 (dd, J = 9.9, 6.4 Hz, 1H), 2.00 (s, 3H), 1.88-1.98 (m, 1H), 1.67 (bs, 3H), 1.45-1.55 (m, 2H), 1.27-1.41 (m, 4H), 0.99 (d, J = 6.5, 3H), 0.90 (t, J = 7.1, 3H), 0.77 (d, J = 6.7, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 169.9, 142.3, 133.9, 128.4 (x2), 128.3, 125.7, 87.6, 76.4, 65.8, 57.9, 35.9, 31.0, 29.7, 28.2, 27.8, 22.2, 21.3, 21.2, 20.9, 18.8, 14.6, 14.0. IR (CCl_4) 2958, 2931, 1738, 1453, 1370, 1236, 1020 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 386 (MNH_4^+), 326, 309.

II- Au(I) catalysed isomerisation of 3-hydroxylated 1,5-enynes

General procedure for the isomerisation of 3-hydroxylated 1,5-enynes

To a 0.1 M solution of $(\text{PPh}_3)\text{AuCl}$ (2 mol%) and AgBF_4 (2 mol%) in dry dichloromethane under argon and at the stated conditions of temperature (cf eq 2,3,5 and table 1) was added the enyne dissolved in a minimum amount of dry dichloromethane. The reaction was stirred until full consumption of the starting material (as attested by TLC analysis). The resulting mixture was filtered through a pad of silica pre-impregnated with dichloromethane, evaporated and purified by flash column chromatography.

Rq: Some of the benzyl enol ether products obtained when the reaction was performed starting from benzyl derivatives of enynes were relatively instable (i.e. **2e**, **2g**, **3g**, **2i** and **3i**). In that special cases, the reaction mixture was preferentially filtered through silicagel pre-impregnated with wet ether to cleave the enol function and furnish the corresponding keto products.

Isomerisation of enyne 1a

2-Isopropyl-3-methyl-4-pentyl-5-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enol (3a)

Yield 55%. ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.31 (m, 2H), 7.18-7.22 (m, 3H), 5.33 (bt, $J = 7.5$ Hz, 1H), 4.16 (bs, 1H), 2.74 (t, $J = 7.7$ Hz, 2H), 2.53-2.67 (m, 2H), 2.27-2.31 (m, 3H), 1.98 (septd, $J = 6.9, 3.6$ Hz, 1H), 1.71 (bs, 3H), 1.25-1.43 (m, 7H), 1.05 (d, $J = 6.9$, 3H), 0.89 (t, $J = 6.9$, 3H), 0.62 (d, $J = 6.9$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 148.7, 143.8, 142.0, 136.4, 128.5, 128.4, 125.9, 119.9, 75.4, 61.7, 36.9, 32.0, 29.8, 29.1, 27.9, 26.9, 22.7, 20.9, 16.7, 14.2, 13.5. IR (CCl_4) 3600, 2958, 2929, 2871, 1495, 1454, 1379, 1119 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 344 (MNH_4^+), 327 (MH^+), 309.

Isomerisation of enyne 1b

2-Isopropyl-3-methyl-4-pentyl-5-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enyl benzyl ether (3b)

Yield 80%. ^1H NMR (400 MHz, CDCl_3) δ 7.19-7.36 (m, 10H), 5.35 (bt, $J = 7.4$ Hz, 1H), 4.52 (d, $J = 12.1$ Hz, 1H), 4.48 (d, $J = 12.1$ Hz, 1H), 4.02 (bs, 1H), 2.74-2.80 (m, 2H), 2.57-2.72 (m, 2H), 2.43 (d, $J = 3.1$ Hz, 1H), 2.33 (bt, $J = 7.4$ Hz, 2H), 1.95 (septd, $J = 6.9, 3.6$ Hz, 1H), 1.75 (bs, 3H), 1.29-1.46 (m, 6H), 1.04 (d, $J = 6.9$, 3H), 0.91 (t, $J = 6.9$, 3H), 0.62 (d, $J = 6.9$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 144.6, 144.0, 142.0, 139.3, 137.1, 128.6, 128.4, 128.3, 127.9, 127.3, 125.8, 121.6, 82.3, 68.1, 58.9, 36.9, 32.0, 29.8, 29.1, 28.3, 27.1, 22.7, 21.3, 17.0, 14.3, 13.7. IR (CCl_4) 2957, 2929, 2871, 1496, 1453, 1084, 1065 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 435 (MNH_4^+), 418 (MH^+), 307.

Isomerisation of enyne 1c

Acetic acid 2-isopropyl-3-methyl-4-pentyl-5-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enyl ester (3c)

Yield 74%. ^1H NMR (400 MHz, CDCl_3) δ 7.26-7.30 (m, 2H), 7.17-7.20 (m, 3H), 5.51 (bs, 1H), 5.33 (bt, $J = 7.7$ Hz, 1H), 2.65-2.77 (m, 2H), 2.53-2.60 (m, 2H), 2.34 (bs, 1H), 2.30 (t, $J = 7.2$ Hz, 2H), 2.05 (s, 3H), 1.97 (septd, $J = 6.9, 3.5$ Hz, 1H), 1.73 (bs, 3H), 1.25-1.44 (m, 6H), 0.99 (d, $J = 6.9$, 3H), 0.89 (t, $J = 6.9$, 3H), 0.67 (d, $J = 6.9$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 170.9, 144.0, 143.4, 142.0, 137.3, 128.5, 128.4, 125.9, 122.0, 77.2, 59.2, 36.8, 31.9, 29.9, 28.9, 28.0, 27.0, 22.7, 21.8, 20.8, 16.8, 14.2, 13.5. IR (CCl_4) 2958, 2930, 2872, 1732, 1496, 1370, 1240, 1019 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 369 (MH^+), 309.

Acetic acid (1R,5S,6S)-4-isopropyl-5-methyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-en-2-yl ester (2c)

Yield 10%. ^1H NMR (400 MHz, CDCl_3) δ 7.31-7.35 (m, 2H), 7.21-7.26 (m, 3H), 5.21 (d, $J = 2.0$ Hz, 1H), 2.66-2.81 (m, 2H), 2.36 (dd, $J = 9.6, 1.7$ Hz, 1H), 2.21 (s, 3H), 1.63-1.73 (m, 2H), 1.53-1.62 (m, 1H), 1.43-1.52 (m, 1H), 1.23-1.42 (m, 6H), 1.21 (s, 3H), 1.10-1.18 (m, 1H), 1.09 (d, $J = 6.7$, 3H), 0.96 (d, $J = 6.6$, 3H), 0.71 (t, $J = 7.1$, 3H), 0.90-0.95 (m, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 168.5, 155.7, 142.8, 128.5, 128.4, 125.8, 112.6, 57.9, 37.9, 36.5, 33.2, 32.5, 32.7, 28.8, 27.5, 26.8, 23.1, 22.7, 22.6, 21.9, 21.2, 16.0, 14.1. IR (CCl_4) 2956, 2928, 2869, 1762, 1454, 1366, 1209 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 369 (MH^+), 327, 309.

Isomerisation of enyne 1d

(1R,4R,5R,6S)-4-Isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (2d)

Yield 66%. ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.31 (m, 2H), 7.17-7.21 (m, 3H), 2.66-2.79 (m, 2H), 2.32 (dd, $J = 19.0, 0.2$ Hz, 1H), 2.21 (d, $J = 19.0$ Hz, 1H), 1.86-2.01 (m, 3H), 1.24-1.62 (m, 9H), 0.93 (d, $J = 6.8$, 3H), 0.90 (t, $J = 7.0$, 3H), 0.89 (d, $J = 6.8$, 3H), 0.72 (d, $J = 3.9$ Hz, 1H), 0.28-0.32 (m, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 142.2, 128.57, 128.4, 125.9, 59.6, 47.3, 36.2, 32.4, 31.7, 30.9, 30.8, 30.7, 28.6, 28.2, 27.9, 22.7, 20.8, 19.4, 14.1. IR (CCl_4) 2959, 2929, 2857, 1737, 1495, 1464 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 330 (MNH_4^+), 313 (MH^+).

(1S,4R,5S,6R)-4-Isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (3d)

Yield 14%. ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.31 (m, 2H), 7.18-7.22 (m, 3H), 2.67-2.80 (m, 2H), 2.50-2.55 (m, 2H), 2.19 (d, J = 18.9 Hz, 1H), 1.87-1.99 (m, 1H), 1.78-1.87 (m, 1H), 1.55-1.65 (m, 1H), 1.48-1.53 (m, 2H), 1.24-1.46 (m, 6H), 1.11 (d, J = 6.7, 3H), 1.07 (dd, J = 4.9, 4.3 Hz, 1H), 0.94 (d, J = 6.8, 3H), 0.90 (t, J = 7.0, 3H), 0.40-0.44 (m, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 217.6, 142.2, 128.5, 128.4, 126.0, 57.4, 46.7, 36.4, 32.3, 31.0, 30.6, 28.6, 28.0, 27.7, 27.6, 24.3, 22.8, 21.9, 30.6, 14.2. IR (CCl_4) 2957, 2928, 2857, 1740, 1454 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 330 (MNH_4^+), 313 (MH^+).

Isomerisation of enyne 1e

(1S,5R)-5-Isopropyl-3-pentyl-2-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enyl benzyl ether (3e)

Yield 27%. ^1H NMR (400 MHz, CDCl_3) δ 7.18-7.41 (m, 10H), 5.77 (bs, 1H), 5.49 (bt, J = 7.5 Hz, 1H), 4.53 (d, J = 11.9 Hz, 1H), 4.49 (d, J = 11.9 Hz, 1H), 4.13 (bs, 1H), 2.72-2.82 (m, 2H), 2.61-2.72 (m, 2H), 2.45-2.48 (m, 1H), 2.35 (bt, J = 7.6 Hz, 2H), 1.56-1.68 (m, 1H), 1.49-1.59 (m, 2H), 1.29-1.39 (m, 4H), 0.94 (d, J = 6.7, 3H), 0.92 (t, J = 7.1, 3H), 0.87 (d, J = 6.7, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 143.7 (x2), 141.95, 139.2, 134.6, 128.6, 128.4, 128.4, 127.9, 127.4, 125.9, 123.9, 85.7, 68.5, 54.9, 36.8, 31.9, 31.3, 30.7, 30.1, 28.3, 22.7, 20.9, 19.8, 14.2. IR (CCl_4) 2957, 2929, 2870, 1604, 1496, 1453, 1089, 1066 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 421 (MNH_4^+), 293.

(1R,5R,6S)-3-Benzyloxy-4-isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-ene (2e)

Isolated as the deprotected form of the benzyl enol ether, i.e. ketone **2d**, Yield 54%

Isomerisation of enyne 1f

(1R,4R,5R,6S)-4-Isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (2d)

Yield 17%.

(1S,4R,5S,6R)-4-Isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (3d)

Yield 70%.

Isomerisation of enyne 1g

(1R,5R,6S)-3-Benzyloxy-4-isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-ene (2e)

Isolated as the deprotected form of the benzyl enol ether, i.e. ketone **2d**, Yield 15%

(1S,5S,6R)-3-Benzyloxy-4-isopropyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-ene (3g)

Isolated as the deprotected form of the benzyl enol ether, i.e. ketone **3d**, Yield 63%.

Isomerisation of enyne 1h

4-Methyl-3-pentyl-2-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enol (4h)

Yield 29%. ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.32 (m, 2H), 7.18-7.22 (m, 3H), 5.34 (bt, J = 7.5 Hz, 1H), 4.45 (bd, J = 6.2 Hz, 1H), 2.75 (t, J = 7.8 Hz, 2H), 2.55-2.67 (m, 3H), 2.29 (t, J = 7.5 Hz, 2H), 2.20 (d, J = 17.2 Hz, 1H), 1.77 (bs, 3H), 1.24-1.49 (m, 7H), 0.89 (t, J = 6.9, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 149.1, 142.0, 141.2, 135.5, 128.5, 128.4, 125.9, 119.8, 74.2, 45.3, 37.0, 31.9, 29.7, 29.0, 26.8, 22.7, 14.9, 14.2. IR (CCl_4) 3597, 2929, 1454, 1263 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 302 (MNH_4^+), 285 (MH^+), 267.

(1S,5R,6S)-1-Methyl-5-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (2h) and (1S,5R,6R)-1-Methyl-5-pentyl-6-phenethyl-bicyclo[3.1.0]hexan-3-one (3h)

Yield 63% / inseparable mixture of isomers **2h** and **3h** (d.r. 2.3:1). Isomer **2h** ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.31 (m, 2H), 7.15-7.22 (m, 3H), 2.59-2.72 (m, 2H), 2.49 (d, $J = 18.5$ Hz, 1H), 2.44 (d, $J = 18.5$ Hz, 1H), 2.36 (d, $J = 18.5$ Hz, 1H), 2.25 (d, $J = 18.5$ Hz, 1H), 1.75-1.84 (m, 1H), 1.54-1.67 (m, 1H), 1.24-1.54 (m, 8H), 1.08 (s, 3H), 0.90 (t, $J = 6.9$, 3H), 0.30 (dd, $J = 8.1, 5.8$ Hz, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ 217.8, 142.2, 128.5, 128.4, 126.0, 50.1, 47.2, 36.5, 32.6, 32.4, 29.9, 27.4, 27.3, 27.2, 25.7, 22.7, 14.1, 12.5. IR (CCl_4) 2928, 2858, 1745, 1454, 1406, 1153 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 302 (MNH_4^+), 285 (MH^+).

Isomerisation of enyne 1i

4-Methyl-3-pentyl-2-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enyl benzyl ether (4i)

Yield 57%. ^1H NMR (400 MHz, CDCl_3) δ 7.20-7.42 (m, 10H), 5.40 (bt, $J = 7.5$ Hz, 1H), 4.57 (d, $J = 11.9$ Hz, 1H), 4.50 (d, $J = 11.9$ Hz, 1H), 4.35 (bd, $J = 6.5$ Hz, 1H), 2.75-2.81 (m, 2H), 2.52-2.66 (m, 3H), 2.30-2.37 (m, 3H), 1.78 (bs, 3H), 1.37-1.46 (m, 2H), 1.25-1.37 (m, 4H), 0.90 (t, $J = 6.8$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 144.7, 142.2, 140.8, 139.0, 136.3, 128.5, 128.4, 128.3, 127.9, 127.4, 125.9, 120.7, 80.2, 69.3, 42.2, 37.0, 31.9, 29.9, 29.0, 26.9, 22.7, 14.9, 14.2. IR (CCl_4) 2957, 2929, 2857, 1496, 1454, 1068, 1028 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 392 (MNH_4^+), 375 (MH^+), 357, 267.

(1S,5S,6S)-3-Benzyloxy-5-methyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-ene (2i) and (1S,5S,6R)-3-benzyloxy-5-methyl-1-pentyl-6-phenethyl-bicyclo[3.1.0]hex-2-ene (3i)

Isolated as a mixture of the deprotected form of the benzyl enol ether, i.e. ketone **2h** and **3h**, (d.r. 2.3:1), Yield 30%.

Isomerisation of enyne 1j

(1S,2R)-2-Isopropyl-3-methyl-4-phenyl-5-[3-phenyl-prop-(E)-ylidene]-cyclopent-3-enyl benzyl ether (4j)

Yield 54%. ^1H NMR (400 MHz, CDCl_3) δ 7.27-7.41 (m, 8H), 7.10-7.21 (m, 5H), 6.90 (d, $J = 6.9$ Hz, 2H), 5.41 (t, $J = 7.9$ Hz, 1H), 4.61 (s, 2H), 4.13 (s, 1H), 2.56 (d, $J = 3.5$ Hz, 2H), 2.39-2.51 (m, 2H), 1.99-2.07 (m, 1H), 1.82-1.95 (m, 2H), 1.67 (bs, 3H), 1.08 (d, $J = 6.9$, 3H), 0.74 (d, $J = 7.0$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 146.1, 144.9, 141.9, 139.3, 138.5, 138.4, 129.4, 128.5, 128.4, 128.2, 128.1, 127.9, 127.4, 126.8, 125.6, 123.5, 82.5, 68.3, 59.9, 36.2, 29.4, 28.8, 21.3, 17.4, 14.6. IR (CCl_4) 3028, 2958, 2927, 1601, 1495, 1453, 1386, 1065, 1028 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 441 (MNH_4^+), 424 (MH^+), 402, 313.

(1R,5S,6R)-3-Benzyl-4-isopropyl-5-methyl-6-phenethyl-1-phenyl-bicyclo[3.1.0]hex-2-ene (2j) and 3j

Isolated as a mixture of **2j** and **3j**, (d.r. 1:3.3), Yield 33%.

Structure of compound **3j** was confirmed by deprotection of the enol ether form in acidic medium (THF / HCl 1N):

(S)-6-Isopropyl-5-methyl-4-phenethyl-3-phenyl-cyclohex-3-enone (8)

^1H NMR (400 MHz, CDCl_3) δ 7.36 (t, $J = 7.3$ Hz, 2H), 7.28 (t, $J = 7.3$ Hz, 1H), 7.22 (t, $J = 7.2$ Hz, 2H), 7.11-7.17 (m, 3H), 7.00 (d, $J = 7.0$ Hz, 2H), 3.08 (d, $J = 21.8$ Hz, 1H), 2.99 (dd, $J = 21.8, 1.5$ Hz, 1H), 2.86 (qd, $J = 7.0, 1.3$ Hz, 1H), 2.72-2.80 (m, 1H), 2.57-2.64 (m, 1H),

2.38-2.46 (m, 1H), 2.13-2.21 (m, 1H), 2.03-2.12 (m, 1H), 1.97 (dd, $J = 10.3, 1.3$ Hz, 1H), 1.12 (d, $J = 7.2$, 3H), 1.08 (d, $J = 6.4$, 3H), 0.93 (d, $J = 6.6$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 212.6, 141.7, 141.3, 137.1, 130.6, 128.6, 128.5, 128.2, 128.0, 127.0, 126.0, 64.6, 44.4, 37.9, 35.5, 34.3, 28.0, 21.6, 21.3, 21.0. IR (CCl_4) 2961, 2927, 1714, 1495, 1454, 1389 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 350 (MNH_4^+), 333 (MH^+).

Isomerisation of enyne 1k

(2E,5E)-5,7-Dimethyl-4-phenethyl-octa-2,5-dienal (2k)

Yield 68%. ^1H NMR (400 MHz, CDCl_3) δ 9.53 (d, $J = 7.9$ Hz, 1H), 7.28-7.32 (m, 2H), 7.17-7.23 (m, 3H), 6.76 (dd, $J = 15.7, 7.1$ Hz, 1H), 6.76 (dd, $J = 15.7, 7.1$ Hz, 1H), 6.10 (ddd, $J = 15.7, 7.8, 1.2$ Hz, 1H), 5.15 (d, $J = 9.1$ Hz, 2H), 2.89 (td, $J = 7.3, 7.3$ Hz, 1H), 2.51-2.65 (m, 3H), 1.80-1.97 (m, 2H), 1.59 (s, 3H), 0.99 (d, $J = 6.7$, 3H), 0.97 (d, $J = 6.7$, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ 193.1, 160.0, 140.6, 135.5, 131.1, 129.9, 127.4 (x2), 124.9, 49.9, 32.4, 32.1, 26.2, 22.0, 21.9, 12.4. IR (CCl_4) 2958, 2931, 1694, 1630, 1496, 1454, 1127 cm^{-1} . MS ($\text{CI}^+ \text{NH}_3$) m/z 274 (MNH_4^+), 257 (MH^+).

5

1j

8

2k