Unusual Tandem Alkynylation and *trans*-Hydrosilylation to Form Oxasilacyclopentenenes

Sarah V. Maifeld and Daesung Lee*
Department of Chemistry, University of Wisconsin,
Madison, WI 53706
dlee@chem.wisc.edu

Supporting Information

Table of Contents:

Materials and Methods.....	S2
Experimental procedures and spectroscopic data.....	S2–S11
¹ H NMR and ¹³ C NMR spectra.....	S12–S74

Materials and Methods. Unless otherwise noted, all reagents were purchased without further purification. THF was distilled over Na and benzophenone. Reactions were monitored by thin layer chromatography (TLC) using Fisher Silica Gel 60 F₂₅₄ precoated plates. Silica gel (40-63 micrometer particle size) was purchased from Silicycle. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker AC-300 spectrometer. Chemical shifts are reported in δ values relative to the internal standard, tetramethylsilane (TMS). High resolution mass spectra were obtained using Leu5-enkephalin or erythromycin as a lock mass on a Micromass LCT ESI-TOF spectrometer equipped with Z-sprayTM and a reflectron.

Representative procedure for the synthesis of alkynylsilanes:

Compound 4b, diphenyl(ethynyl)silane. To a stirred solution of phenylacetylene (2 g, 0.020 mol) in 20 mL of dry THF at -78°C was added *n*-BuLi (8.8 mL of a 2.5 M solution in hexanes, 0.022 mol). After stirring for 15 minutes, chlorodiphenyl silane (4.4 mL, 0.024 mol) was added. Reaction was stirred for 30 minutes at -78°C , then allowed to warm to ambient temperature and stir for an additional two hours. The reaction was then diluted with diethyl ether and quenched with saturated NH_4Cl solution. The ethereal layer was washed with brine, dried over MgSO_4 and concentrated to a light yellow oil. Flash column chromatography with hexanes yielded, upon concentration, 5.12 g of white solid. (90% yield). ¹H NMR (CDCl_3 , 300 MHz) δ 7.73 (m, 4H), 7.57 (m, 2H), 7.46 – 7.28 (m, 9H), 5.31 (s, 1H); ¹³C NMR (CDCl_3 , 75 MHz) δ 135.1, 134.2, 132.1, 130.0, 129.0, 128.2, 128.0, 122.4, 109.4, 87.0; HRMS (EI) calcd for $\text{C}_{20}\text{H}_{16}\text{Si}$ (M)⁺: 284.1021, found 284.1034.

Compound 4c, diisopropyl(phenylethynyl)silane. ¹H NMR (CDCl_3 , 300 MHz) δ 7.51 (m, 2H), 7.33 (m, 3H), 3.84 (br s, 1H), 1.12 (m, 14H); ¹³C NMR (CDCl_3 , 75 MHz) δ

131.9, 128.4, 128.0, 122.9, 107.6, 87.7, 18.4, 18.1, 10.7; HRMS (EI) calcd for $C_{14}H_{20}Si$ (M) $^{*+}$: 216.1334, found 216.1330.

Representative procedure for the synthesis of oxasilacyclopentenones:

Compound 6. To a stirred solution of acetophenone (0.050 g, 0.42 mmol) in 3 mL THF was added dimethyl(phenylethynyl)silane (0.080 g, 0.50 mmol) followed by potassium *tert*-butoxide (95%, 5 mg, 0.042 mmol). After stirring for twenty minutes, the reaction was quenched with an aqueous solution of saturated NH_4Cl and extracted twice with ether. The ethereal layer was washed with brine, dried over $MgSO_4$ and concentrated to an orange oil. Flash column chromatography with 1% diethyl ether in hexanes yielded, upon concentration, 57 mg of a slightly yellow oil (48%). 1H NMR ($CDCl_3$, 300 MHz) δ 7.49 (d, $J = 7.2$, 2H), 7.32 (m, 6H), 7.23 (m, 2H), 7.15 (s, 1H), 1.74 (s, 3H), 0.52 (s, 3H), 0.47 (s, 3H); ^{13}C NMR ($CDCl_3$, 75 MHz) δ 149.3, 146.3, 138.0, 137.0, 128.6, 128.1, 127.1, 126.7, 126.5, 124.5, 86.1, 30.7, 1.6, 0.9; HRMS (EI) calcd for $C_{18}H_{20}OSi$ (M) $^{*+}$: 280.1283, found 280.1282.

Compound 7a. 1H NMR ($CDCl_3$, 300 MHz) δ 7.33 (m, 4H), 7.24 (m, 1H), 7.04 (s, 1H), 1.73 (m, 3H), 1.58 (m, 6H), 1.37 (m, 1H), 0.40 (s, 6H); ^{13}C NMR ($CDCl_3$, 75 MHz) δ 149.6, 139.0, 137.6, 128.5, 126.8, 126.5, 84.5, 38.6, 25.2, 22.5, 1.7; HRMS (EI) calcd for $C_{16}H_{22}OSi$ (M) $^{*+}$: 258.1440, found 258.1441.

Compound 7b. 1H NMR ($CDCl_3$, 300 MHz) δ 7.67 (m, 4H), 7.45 – 7.33 (m, 8H), 7.29 (s, 1H), 7.26 – 7.15 (m, 3H), 1.91 – 1.72 (m, 6H), 1.61 (m, 3H), 1.40 (m, 1H). ^{13}C NMR

(CDCl₃, 75 MHz) δ 151.8, 137.4, 136.5, 135.0, 134.6, 130.1, 128.4, 127.8, 127.1, 127.0, 85.2, 38.4, 25.3, 22.4; HRMS (EI) calcd C₂₆H₂₆OSi (M)⁺ : 382.1753, found 382.1734.

Compound 8. ¹H NMR (CDCl₃, 300 MHz) δ 7.44 (m, 2H), 7.31 (m, 3H), 2.10 (br d, 1H), 2.00 (m, 2H), 1.65 (m, 7H), 1.27 (m, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 131.5, 128.04, 128.0, 122.7, 92.6, 84.2, 68.9, 39.9, 25.0, 23.2; HRMS (EI) calcd C₁₄H₁₆O (M)⁺ : 200.1201, found 200.1204.

Compound 9c. ¹H NMR (CDCl₃, 300 MHz) δ 7.42 (m, 2H), 7.30 (m, 3H), 4.48 (br s, 1H), 2.00 (m, 2H), 1.73 (m, 7H), 1.32 (1H), 1.06 (m, 14H); ¹³C NMR (CDCl₃, 75 MHz) δ 131.3, 128.0, 127.8, 123.1, 92.7, 85.1, 70.7, 40.7, 25.1, 23.0, 17.5, 12.6; HRMS (EI) calcd C₂₀H₃₀OSi (M)⁺ : 314.2066, found 314.2064.

Compound 11a. ¹H NMR (CDCl₃, 300 MHz) δ 7.33 (m, 10H), 7.02 (m, 1H), 5.85 (m, 1H), 0.54 (s, 3H), 0.50 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 144.4, 142.0, 140.1, 136.9, 128.6, 128.4, 127.5, 127.2, 126.7, 126.0, 83.7, 1.3, 0.6; HRMS (EI) calcd C₁₇H₁₈OSi (M)⁺ : 266.1127, found 266.1121.

Compound 11b. ¹H NMR (CDCl₃, 300 MHz) δ 7.36 (m, 4H), 7.26 (m, 1H), 7.01 (br d, *J* = 2.4, 1H), 6.32 (dd, *J* = 3, 1.8, 1H), 6.25 (br d, *J* = 3.3, 1H), 5.84 (br d, *J* = 1.5, 1H), 0.462 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 154.6, 142.3, 142.2, 140.4, 136.8, 128.7,

127.4, 126.7, 110.1, 106.4, 77.3, 1.3, 0.4; HRMS (EI) calcd for C₁₅H₁₆O₂Si (M)⁺ : 256.0920, found 256.0921.

Compound 11c. ¹H NMR (CDCl₃, 300 MHz) δ (reported as a 1.3:1 mixture of *E*- and *Z*-isomers) 7.33 (m, 4H), 7.25 (m, 1H), 6.83 (br d, *J* = 1.8, 0.57 H), 6.80 (br d, *J* = 1.8, 0.44H), 5.54 (m, 1H), 5.09 (m, 2H), 2.33 (m, .44H), 2.10 (m, 3.56 H), 1.80 (m, 1.71H), 1.76 (m, 1.32H), 1.72 (br s, 1.32H), 1.68 (br s, 1.71H), 1.63 (br s, 1.32H), 1.61 (br s, 1.71H), 0.44 (br s, 3.42H), 0.41 (m, 2.64 H); δ ¹³C NMR (CDCl₃, 75 MHz) (reported as a 1.3: 1 mixture of *E*- and *Z*-isomers) 144.4, 144.1, 140.1, 140.0, 138.3, 138.1, 137.2, 131.9, 131.4, 128.5, 127.0, 126.54, 126.52, 126.5, 125.5, 123.8, 123.7, 78.7, 78.5, 39.4, 32.3, 26.7, 26.3, 25.6, 25.5, 23.3, 17.5, 17.5, 16.5, 1.66, 1.64, 0.57, 0.56; HRMS (EI) calcd C₂₀H₂₈OSi (M)⁺ : 312.1909, found 312.1910.

Compound 11d. ¹H NMR (CDCl₃, 300 MHz) δ (major diastereomer reported) 7.34 (m, 4H), 7.26 (m, 1H), 7.04 (s, 1H), 5.57 (m, 1H), 4.74 (m, 2H), 2.52 (m, 1H), 2.20 (m, 1H), 2.01 (m, 2H), 1.89 (m, 1H), 1.74 (m, 3H), 1.60 (m, 3H), 0.46 (s, 3H), 0.44 (s, 3H); (minor diastereoisomer reported) 7.34 (m, 4H), 7.26 (m, 1H), 6.77 (s, 1H), 5.71 (m, 1H), 4.74 (m, 2H), 2.52 (m, 1H), 2.20 (m, 1H), 2.01 (m, 2H), 1.89 (m, 1H), 1.76 (m, 3H), 1.63 (m, 3H), 0.47 (s, 3H), 0.42 (s, 3H); δ ¹³C NMR (CDCl₃, 75 MHz) (2.2 : 1 mixture of diastereomers) 149.2, 148.7, 148.6, 147.2, 140.9, 140.2, 137.1, 136.2, 134.6, 128.57, 128.55, 127.09, 127.06, 126.8, 126.6, 126.5, 125.8, 124.0, 109.0, 108.6, 87.0, 85.3, 43.5, 42.8, 40.3, 37.1, 31.2, 31.0, 20.8, 20.1, 18.4, 17.3, 2.0, 1.9, 0.7, 0.2; HRMS (EI) calcd C₂₀H₂₆OSi (M)⁺ : 310.1753, found 310.1744.

Compound 11e. ^1H NMR (CDCl_3 , 300 MHz) δ 7.33 (m, 4H), 7.25 (m, 1H), 6.87 (s, 1H), 5.11 (m, 1H), 2.00 (m, 2H), 1.67 (m, 5H), 1.57 (s, 3H), 1.38 (s, 3H), 0.43 (s, 3H), 0.41 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 149.0, 139.5, 137.3, 131.3, 128.5, 126.9, 126.5, 124.3, 85.5, 42.0, 29.0, 25.5, 23.0, 17.5, 1.8, 0.9; HRMS (EI) calcd $\text{C}_{18}\text{H}_{26}\text{OSi}$ (M) $^{++}$: 286.1753, found 286.1760.

Compound 11f. ^1H NMR (CDCl_3 , 300 MHz) δ 7.33 (m, 4H), 7.24 (m, 1H), 7.06 (s, 1H), 1.82 – 1.53 (m, 12H), 0.40 (s, 6H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 150.6, 137.6, 137.5, 128.5, 126.8, 126.5, 88.2, 42.1, 29.6, 22.7, 1.6; HRMS (EI) calcd $\text{C}_{17}\text{H}_{24}\text{OSi}$ (M) $^{++}$: 272.1596, found 272.1601.

Compound 12a, dimethyl((4-methoxyphenyl)ethynyl)silane. ^1H NMR (CDCl_3 , 300 MHz) δ 7.41 (d, $J = 9$, 2H), 6.81 (d, $J = 9$, 2H), 4.26 (sept, $J = 3.6$, 1H), 3.79 (s, 3H), 0.30 (d, $J = 3.6$, 6H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 159.8, 133.4, 114.8, 113.7, 106.4, 89.3, 55.1, -3.0; HRMS (EI) calcd for $\text{C}_{11}\text{H}_{14}\text{OSi}$ (M) $^{++}$: 190.0814, found 190.0815.

Compound 12b, dimethyl(o-tolyethynyl)silane. ^1H NMR (CDCl_3 , 300 MHz) δ 7.43 (br d, $J = 7.5$, 1H), 7.21 (m, 2H), 7.11 (td, $J = 7.2$, 2.1, 1H), 4.30 (sept, $J = 3.5$, 1H), 2.44 (s, 3H), 0.32 (d, $J = 3.5$, 6H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 140.5, 132.0, 129.2, 128.5, 125.3, 122.4, 105.2, 94.9, 20.4, -3.0; HRMS (EI) calcd for $\text{C}_{11}\text{H}_{14}\text{Si}$ ($\text{M}-\text{CH}_3$) $^{++}$: 159.0630, found 159.0623.

Compound 12c, dimethyl(tolyethynyl)silane. ^1H NMR (CDCl_3 , 300 MHz) δ 7.39 (d, $J = 8$, 2H), 7.12 (d, $J = 8$, 2H), 4.30 (sept, $J = 3.5$, 1H), 2.34 (s, 3H), 0.33 (d, $J = 3.5$, 6H); ^{13}C

NMR (CDCl₃, 75 MHz) δ 138.7, 131.7, 128.8, 119.6, 106.5, 90.5, 21.3, -3.1; HRMS (EI) calcd for C₁₁H₁₄Si (M)⁺: 174.0865, found 174.0866.

Compound 12d, dimethyl((4-fluorophenyl)ethynyl)silane. ¹H NMR (CDCl₃, 300 MHz) δ 7.46 (m, 2H), 7.00 (m, 2H), 4.26 (sept, J = 3.6, 1H), 0.31 (d, J = 3.6, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 162.7 (d, ¹ J_{C-F} = 249), 133.9 (d, ³ J_{C-F} = 8), 118.9, 115.5 (d, ² J_{C-F} = 22), 105.2, 90.8, -3.1; HRMS (EI) calcd for C₁₀H₁₁FSi (M-CH₃)⁺: 163.0379, found 163.0379.

Compound 13c. ¹H NMR (CDCl₃, 300 MHz) δ 7.24 (d, J = 8, 2H), 7.12 (d, J = 8, 2H), 6.99 (s, 1H), 2.33 (m, 3H), 1.73 (m, 3H), 1.57 (m, 6H), 1.42 (m, 1H) 0.40 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 148.7, 138.7, 136.6, 134.6, 129.2, 126.4, 84.5, 38.7, 25.3, 22.5, 21.0, 1.7; HRMS (EI) calcd C₁₇H₂₄OSi (M)⁺ = 272.1596, found 272.1601.

Compound 13d. ¹H NMR (CDCl₃, 300 MHz) δ 7.28 (m, 2H), 7.00 (m, 2H), 6.95 (s, 1H), 1.73 (m, 2H), 1.57 (m, 7H), 1.35 (m, 1H), 0.39 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 161.8 (d, ¹ J_{C-F} = 246), 149.4, 138.0, 133.7, 127.9, (d, ³ J_{C-F} = 8), 115.3 (d, ² J_{C-F} = 22), 84.5, 38.6, 25.2, 22.4, 1.6; HRMS (EI) calcd C₁₆H₂₁FOSi (M)⁺: 276.1346, found 276.1355.

Compound 13e. ¹H NMR (CDCl₃, 300 MHz) δ 7.58 (d, J = 8.5, 2H), 7.42 (d, J = 8.5, 2H), 7.10 (s, 1H), 1.64 (m, 2H), 1.61 (m, 7H), 1.58 (m, 1H), 0.42 (s, 6H); ¹³C NMR

(CDCl₃, 75 MHz) δ 152.2, 141.4, 138.3, 128.7 (q, $^2J_{C-F} = 32$), 126.6, 125.4 (q, $^3J_{C-F} = 4$), 124.1 (q, $^1J_{C-F} = 270$), 86.7, 38.4, 25.2, 22.3, 1.5; HRMS (EI) calcd C₁₇H₂₁OSi (M)⁺: 326.1314, found 326.1303.

Compound 14a. ¹H NMR (CDCl₃, 300 MHz) δ 7.36 (m, 2H), 6.83 (m, 2H), 3.80 (s, 3H), 2.11 (br s, 1H), 1.98 (m, 2H), 1.74 – 1.58 (m, 7H), 1.25 (m, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 159.3, 132.9, 114.8, 113.7, 91.2, 84.0, 68.9, 55.1, 40.0, 25.1, 23.2; HRMS (EI) calcd C₁₅H₁₈O₂ (M)⁺: 230.1307, found = 230.1296.

Compound 14b. ¹H NMR (CDCl₃, 300 MHz) δ 7.39 (br d, $J = 7.2$, 1H), 7.18 (m, 2H), 7.11 (m, 1H), 2.43 (s, 3H), 2.32 (br s, 1H), 2.02 (m, 2H), 1.67 (m, 7H), 1.26 (m, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 139.9, 131.7, 129.2, 128.0, 125.3, 122.5, 96.7, 83.0, 69.1, 40.0, 25.1, 23.3, 20.6; HRMS (EI) calcd C₁₅H₁₈OSi (M)⁺: 214.1358, found 214.1353.

Compound 14c. ¹H NMR (CDCl₃, 300 MHz) δ 7.32 (d, $J = 8$, 2H), 7.12 (d, $J = 8$, 2H), 2.34 (s, 3H), 2.02 (m, 3H), 1.66 (m, 7H), 1.27 (m, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 138.1, 131.2, 128.8, 119.6, 91.9, 84.3, 68.9, 39.9, 25.1, 23.2, 21.2; HRMS (EI) calcd C₁₅H₁₈O (M)⁺: 214.1358, found 214.2.

Compound 14e. ^1H NMR (CDCl_3 , 300 MHz) δ 7.54 (m, 4H), 2.31 (br s, 1H), 2.03 (m, 2H), 1.65 (m, 7H), 1.31 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 131.7, 129.8 (q, $^2J_{\text{C-F}} = 32$), 126.5, 125.0 (q, $^3J_{\text{C-F}} = 4$), 123.7 (q, $^1J_{\text{C-F}} = 271$), 96.2, 82.8, 68.9, 39.7, 24.9, 23.1; HRMS (EI) calcd $\text{C}_{15}\text{H}_{15}\text{F}_3\text{O}$ (M) $^{+}$: 268.1075, found 268.1063.

Compound 17, diphenyl(tolylethynyl)silane. ^1H NMR (CDCl_3 , 300 MHz) δ 7.73 (m, 4H), 7.39 (m, 8H), 7.13 (d, $J = 8$, 2H) 5.29 (s, 1H), 2.35 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 139.3, 135.0, 132.2, 132.0, 129.9, 128.9, 127.9, 119.3, 109.7, 86.1, 21.4; HRMS (EI) calcd for $\text{C}_{21}\text{H}_{18}\text{Si}$ (M) $^{+}$: 298.1178, found 298.1185.

Compound 18. ^1H NMR (CDCl_3 , 300 MHz) δ 7.68 (m, 4H), 7.37 (m, 6H), 7.25 (m, 3H), 7.02 (d, $J = 7.5$, 2H), 2.62 (s, 3H), 1.74 (m, 6H), 1.59 (m, 3H), 1.36 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 151.0, 136.9, 136.3, 135.2, 134.9, 134.7, 130.2, 129.3, 127.9, 127.1, 85.3, 38.6, 25.5, 22.6, 21.1; HRMS (EI) calcd $\text{C}_{27}\text{H}_{28}\text{OSi}$ (M) $^{+}$: 396.1909, found 396.1849.

Compound 19. To a stirred solution of **7a** (0.030 g, 0.12 mmol) in 1.2 mL of solvent (0.6 mL of THF and 0.6 mL of MeOH) was added KF (1.5 equiv, 0.17 mmol, 10 mg), LiOH (3 equiv, 0.36 mmol, 8.6 mg) and H_2O_2 (40 equiv, 4.6 mmol, 0.45 mL of a 30% wt in water solution). After stirring at room temperature for 2 hours, the reaction was quenched by the slow, dropwise addition of a saturated aqueous solution of $\text{Na}_2\text{S}_2\text{O}_3$ and extracted twice with diethyl ether. The ethereal layer was washed with brine, dried with magnesium sulfate and concentrated to an oil. Flash chromatography with 12% diethyl ether in hexanes yielded 20 mg of a clear oil. ^1H NMR (CDCl_3 , 300 MHz) δ 7.95 (m, 2H), 7.59 (tt, $J = 7$, 1H), 7.48 (m, 2H), 4.00 (s, 1H), 3.12 (s, 2H), 1.74 – 1.46 (m, 9H),

1.28 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 201.7, 137.3, 133.3, 128.5, 127.9, 70.8, 47.5, 37.6, 25.6, 21.8; HRMS (ESI) calcd $\text{C}_{14}\text{H}_{18}\text{O}_2$ ($\text{M}+\text{Na}$) $^+$: 241.1204, found 241.1209.

Compound 20. To a stirred solution of **7a** (0.075 g, 0.29 mmol) in 1 mL CH_2Cl_2 was added DMDO¹ (2 equiv of an approximately 0.08 M acetone solution) at room temperature. After an hour, the solvent was removed under a stream of nitrogen. Flash column chromatography with 6% diethyl ether in hexanes yielded 45 mg of a clear oil. ^1H NMR (CDCl_3 , 300 MHz) δ 7.31 (m, 5H), 3.45 (s, 1H), 1.60 (m, 6H), 1.53 (m, 3H), 1.40 (m, 1H), 0.42 (s, 3H), 0.34 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 138.0, 128.2, 126.9, 125.7, 77.9, 71.2, 71.2, 59.4, 36.4, 33.9, 25.5, 22.3, 22.0, 0.9, -3.2.

Compound 21. To a stirred solution of **7a** (0.020 g, 0.07 mmol) in 1 mL THF was added CuI (1.1 equiv, 16 mg, 0.085 mmol), allyl chloride (3 equiv, 0.019 mL, 0.23 mmol) and tetrabutylammonium fluoride (1.2 equiv, 0.092 mmol, 0.092 mL). The reaction was gently heated to 40 °C for thirty minutes. Following the consumption of **7a** on TLC, the reaction was filtered through pad of silica gel and concentrated by rotary evaporation. Flash column chromatography with 13% diethyl ether in hexanes yielded 13 mg of a light yellow oil. ^1H NMR (CDCl_3 , 300 MHz) δ 7.29 (m, 5H), 5.88 (ddt, J = 17, 10.2, 6, 1H), 5.86 (s, 1H), 5.06 (dtd, J = 17, 1.8, 1.8, 1H), 4.98 (dtd, J = 10.5, 1.8, 1.8, 1H), 3.67 (ddd, J = 6.3, 1.8, 1.8, 2H), 1.67 (m, 7H), 1.51 (m, 3H), 1.38 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 143.6, 139.6, 137.5, 136.3, 128.1, 127.0, 126.6, 115.5, 72.4, 39.5, 34.6, 25.4, 22.5; HRMS (EI) calcd $\text{C}_{17}\text{H}_{22}\text{O}$ ($\text{M}-\text{H}_2\text{O}$) $^+$: 224.1565, found 224.1563.

¹ Waldemar, A.; Bialas, J.; Hadjjarapoglou, L. *Chem. Ber.* **1991**, 2377.