Supporting Information

The Utility of Iridium-Complex of Pybox Ligand in Regio- and Enantioselective Allylic Substitution

Hideto Miyabe, Akira Matsumura, Katsuhiko Moriyama, and Yoshiji Takemoto

Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan. FAX: +81-75-753-4569; E-mail: takemoto@pharm.kyoto-u.ac.jp

Table of contents

Experimental procedure for allylic substitution of **2** with amines **1** or oximes **11**: S2.

Characterization data of all obtained compounds **3-4** and **12-13**: S2-14.

Experimental procedure for reduction of **12Aa** and characterization data of **14-17**: S14-15.

Copies of ^1H and ^{13}C NMR spectrum of all obtained compounds: S15-91.

General. Melting points were taken on a YANAGIMOTO micromelting point apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 at 500 MHz, and at 125 MHz, respectively; Tetramethylsilane (TMS) was used as an internal standard. IR spectra were recorded on a JASCO FT/IR-410 Fourier-transfer infrared spectrometer. Low and high resolution mass spectra were obtained by EI or FAB method. Preparative TLC separations were carried out on precoated silica gel plates (E. Merck 60F₂₅₄). Optical rotations were recorded on a JASCO DIP-360 polarimeter with a path length of 1 cm; concentrations are quoted in mg (2 mL). $[\alpha]_D^{25}$ values are measured in $10^{-1} \text{ deg cm}^2 \text{ g}^{-1}$. The ratios of products were determined by ^1H NMR analysis. Enantiomeric excess was determined by high performance liquid chromatography (HPLC) analysis.

1. Experimental procedure for allylic substitution of **2** with amines **1** or oximes **11**

General procedure: A mixture of amine **1** or oxime **11** (1.0 mmol) and $\text{CsOH} \cdot \text{H}_2\text{O}$ or $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ (1.0 mmol) in CH_2Cl_2 (4.0 mL) was stirred under argon atmosphere at 20 °C for 10 min. To the reaction mixture was added a solution of allylic phosphate **2** (1.5 mmol), pybox **5** (0.080 - 0.16 mmol) and $[\text{IrCl}(\text{cod})]_2$ (0.040 - 0.080 mmol) in CH_2Cl_2 (2.0 mL) at -20 °C. After the reaction was completed, the reaction mixture was diluted with saturated aqueous NH_4Cl and then extracted with AcOEt . The organic phase was dried over MgSO_4 and concentrated at reduced pressure. The ratio of products was determined by ^1H NMR analysis of crude products. Purification of the residue by preparative TLC (hexane: AcOEt =5:1 - 25:1, 2-fold development) afforded the products.

2. Characterization Data

N-(Benzyloxy)-*N*-((*S*)-1-phenylallyl)benzamide (**3Aa**, $\text{R}^1=\text{Bz}$, $\text{R}^2=\text{OBn}$).

A colorless oil. IR (CHCl_3) 1635, 1495, 1451 cm^{-1} . ^1H NMR (CDCl_3) δ 7.70 (2H, d, $J = 7.0$ Hz), 7.60-7.15 (11H, m), 6.83 (2H, d, $J = 7.0$ Hz), 6.32 (1H, ddd, $J = 17.7, 11.0, 6.1$ Hz), 6.15 (1H, d, $J = 6.1$ Hz), 5.39 (1H, d, $J = 17.7$ Hz), 5.38 (1H, d, $J = 11.0$ Hz), 4.48 (1H, d, $J = 8.5$ Hz), 4.11 (1H, d, $J = 8.5$ Hz). ^{13}C NMR (CDCl_3) δ 170.6, 138.1, 134.9, 134.2, 134.1, 130.6, 129.5, 128.8, 128.6 (2C), 128.3, 128.3, 128.1, 118.6, 78.5, 64.1. One carbon peak was missing due to overlapping. MS (EI^+) m/z : 343 (M^+ , 11), 115 (100). HRMS calcd for $\text{C}_{23}\text{H}_{21}\text{NO}_2$: 343.1572, Found: 343.1570. HPLC (Chiralcel OD-H, hexane/2-propanol=95/5, 0.5 ml/min, 254 nm) t_r (*S*) = 15.8 min, t_r (*R*) = 18.1 min. A sample of 92%ee (*S*) by HPLC analysis gave $[\alpha]_D^{29} -33.8$ (c 1.02, CHCl_3).

N-(Benzyloxy)-*N*-cinnamylbenzamide (**4Aa**, $\text{R}^1=\text{Bz}$, $\text{R}^2=\text{OBn}$).

A colorless oil. IR (CHCl₃) 1635, 1496, 1450 cm⁻¹. ¹H NMR (CDCl₃) δ 7.67 (2H, d, *J* = 7.3 Hz), 7.49-7.22 (11H, m), 7.07 (2H, d, *J* = 6.7 Hz), 6.61 (1H, d, *J* = 15.9 Hz), 6.31 (1H, dt, *J* = 15.9, 6.1 Hz), 4.72 (2H, s), 4.49 (2H, d, *J* = 6.1 Hz). ¹³C NMR (CDCl₃) δ 170.3, 136.5, 134.5, 134.3, 134.0, 130.6, 129.5, 128.8, 128.6, 128.5, 128.3, 128.1, 127.9, 126.6, 123.4, 77.1, 50.5. One carbon peak was missing due to overlapping. MS (FAB⁺) *m/z*: 344 (M+H⁺, 64), 117 (100). HRMS calcd for C₂₃H₂₂NO₂ (M+H⁺): 344.1651, Found: 344.1656.

***N*-(Benzyloxy)-*N*-((*S*)-1-(4-chlorophenyl)allyl)benzamide (3Ab, R¹=Bz, R²=OBn).**

A colorless oil. IR (CHCl₃) 1639, 1492, 1449 cm⁻¹. ¹H NMR (CDCl₃) δ 7.67 (2H, d, *J* = 7.0 Hz), 7.52-7.19 (10H, m), 6.87 (2H, d, *J* = 7.3 Hz), 6.29 (1H, ddd, *J* = 15.9, 9.5, 6.7 Hz), 6.09 (1H, d, *J* = 6.7 Hz), 5.40 (1H, d, *J* = 9.5 Hz), 5.37 (1H, d, *J* = 15.9 Hz), 4.54 (1H, br d, *J* = 8.9 Hz), 4.22 (1H, br d, *J* = 8.9 Hz). ¹³C NMR (CDCl₃) δ 170.7, 136.8, 134.7, 133.9 (2C), 130.7, 130.0, 129.4, 128.8, 128.4, 128.2, 128.1, 119.2, 78.7, 63.7. Two carbon peaks were missing due to overlapping. MS (EI⁺) *m/z*: 377 (M⁺, 2), 91 (100). HRMS calcd for C₂₃H₂₀ClNO₂ (M⁺): 377.1188, Found: 377.1187. HPLC (Chiralcel AD-H, hexane/2-propanol=90/10, 0.5 ml/min, 254 nm) *t_r* (*S*) = 28.0 min, *t_r* (*R*) = 21.9 min. A sample of 87%ee (*S*) by HPLC analysis gave [α]_D²⁶ -3.7 (*c* 1.15, CHCl₃).

***N*-(4-Chlorocinnamyl)-*N*-(benzyloxy)benzamide (4Ab, R¹=Bz, R²=OBn).**

A colorless oil. IR (CHCl₃) 1637, 1492, 1450 cm⁻¹. ¹H NMR (CDCl₃) δ 7.67 (2H, d, *J* = 7.0 Hz), 7.48 (1H, t, *J* = 7.3 Hz), 7.44-7.20 (9H, m), 7.07 (2H, d, *J* = 6.7 Hz), 6.54 (1H, d, *J* = 15.9 Hz), 6.28 (1H, dt, *J* = 15.9, 6.1 Hz), 4.71 (2H, s), 4.48 (2H, d, *J* = 6.1 Hz). ¹³C NMR (CDCl₃) δ 170.2, 134.9, 134.3, 134.2, 133.5, 132.6, 130.6, 129.5, 128.8, 128.7, 128.5, 128.3, 128.1, 127.7, 124.1, 76.9, 50.3. MS (EI⁺) *m/z*: 377 (M⁺, 0.5), 105 (100). HRMS calcd for C₂₃H₂₀ClNO₂ (M⁺): 377.1188, Found: 377.1184.

***N*-(Benzyloxy)-*N*-((*S*)-1-(naphthalen-1-yl)allyl)benzamide (3Ac, R¹=Bz, R²=OBn).**

A colorless oil. IR (CHCl₃) 1632, 1511, 1495, 1449 cm⁻¹. ¹H NMR (CDCl₃) δ 8.37 (1H, d, *J* = 8.5

Hz), 7.90 (2H, d, $J = 8.2$ Hz), 7.76 (3H, d, $J = 7.9$ Hz), 7.65 (1H, t, $J = 7.0$ Hz), 7.57-7.35 (5H, m), 7.20-7.05 (4H, m), 6.50-6.40 (3H, m), 5.59 (1H, d, $J = 17.1$ Hz), 5.48 (1H, d, $J = 10.7$ Hz), 3.94 (1H, br s), 3.36 (1H, br s). ^{13}C NMR (CDCl_3) δ 170.1, 134.8, 134.1, 133.7, 133.6, 133.4, 132.0, 130.5, 129.5, 129.3, 128.9, 128.5, 128.3, 128.1, 128.0, 127.8, 127.1, 126.1, 125.1, 123.5, 117.8, 78.4, 58.7. MS (EI^+) m/z : 393 (M^+ , 6), 167 (100). HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{NO}_2$ (M^+): 393.1729, Found: 393.1727. HPLC (Chiralcel AD-H, hexane/2-propanol=90/10, 0.5 ml/min, 254 nm) t_r (S) = 17.4 min, t_r (R) = 14.9 min. A sample of 96% ee (S) by HPLC analysis gave $[\alpha]_{\text{D}}^{28} -20.6$ (c 1.07, CHCl_3).

***N*-(Benzyloxy)-*N*-((*E*)-3-(naphthalen-1-yl)allyl)benzamide (4Ac, $\text{R}^1=\text{Bz}$, $\text{R}^2=\text{OBn}$).**

A colorless oil. IR (CHCl_3) 1637, 1496, 1450 cm^{-1} . ^1H NMR (CDCl_3) δ 8.06 (1H, d, $J = 8.9$ Hz), 7.84 (1H, d, $J = 9.2$ Hz), 7.78 (1H, d, $J = 7.9$ Hz), 7.70 (2H, d, $J = 7.0$ Hz), 7.58 (1H, d, $J = 7.0$ Hz), 7.53-7.32 (7H, m), 7.30-7.20 (3H, m), 7.10 (2H, d, $J = 7.0$ Hz), 6.34 (1H, dt, $J = 15.6, 6.1$ Hz), 4.76 (2H, s), 4.61 (2H, d, $J = 6.1$ Hz). ^{13}C NMR (CDCl_3) δ 170.3, 134.5, 134.3, 134.2, 133.6, 131.3, 131.1, 130.6, 129.4, 128.8, 128.5 (2C), 128.3, 128.2, 128.1, 126.6, 126.1, 125.8, 125.6, 124.1, 123.7, 77.1, 50.7. MS (EI^+) m/z : 393 (M^+ , 0.7), 105 (100). HRMS calcd for $\text{C}_{27}\text{H}_{23}\text{NO}_2$ (M^+): 393.1729, Found: 393.1734.

***N*-(Benzoyloxy)-*N*-((*S*)-1-phenylallyl)benzamide (3Ba, $\text{R}^1=\text{Bz}$, $\text{R}^2=\text{OBz}$).**

A colorless oil. IR (CHCl_3) 1767, 1665, 1495, 1451 cm^{-1} . ^1H NMR (CDCl_3) δ 7.84 (2H, d, $J = 7.6$ Hz), 7.68 (2H, d, $J = 7.3$ Hz), 7.55 (1H, t, $J = 7.6$ Hz), 7.45-7.22 (10H, m), 6.22 (1H, ddd, $J = 17.1, 8.9, 6.1$ Hz), 6.11 (1H, d, $J = 6.1$ Hz), 5.40 (1H, d, $J = 8.9$ Hz), 5.37 (1H, d, $J = 17.1$ Hz). ^{13}C NMR (CDCl_3) δ 170.6, 164.0, 134.0, 133.9, 133.8, 131.1, 129.8, 128.6, 128.4, 128.1, 127.8, 127.2, 119.4, 66.1. Two carbon peaks were missing due to overlapping. MS (EI^+) m/z : 357 (M^+ , 7), 116 (100). HRMS calcd for $\text{C}_{23}\text{H}_{19}\text{NO}_3$: 357.1365, Found: 357.1371. HPLC (Chiralcel AD-H, hexane/2-propanol=90/10, 1.0 ml/min, 254 nm) t_r (S) = 19.7 min, t_r (R) = 15.1 min. A sample of 87% ee (S) by HPLC analysis gave $[\alpha]_{\text{D}}^{31} +2.6$ (c 0.96, CHCl_3).

***N*-(Benzoyloxy)-*N*-cinnamylbenzamide (4Ba, $\text{R}^1=\text{Bz}$, $\text{R}^2=\text{OBz}$).**

A colorless oil. IR (CHCl₃) 1764, 1663, 1495, 1449 cm⁻¹. ¹H NMR (CDCl₃) δ 7.91 (2H, d, J = 7.6 Hz), 7.65 (2H, d, J = 7.0 Hz), 7.56 (1H, d, J = 7.3 Hz), 7.44-7.21 (10H, m), 6.61 (1H, d, J = 15.9 Hz), 6.31 (1H, dt, J = 15.9, 6.4 Hz), 4.67 (2H, d, J = 6.4 Hz). ¹³C NMR (CDCl₃) δ 170.8, 164.6, 136.3, 134.4, 134.1, 133.5, 131.1, 129.9, 128.7, 128.6, 128.3, 128.0, 127.9, 127.0, 126.6, 122.6, 52.3. MS (FAB⁺) m/z : MS (EI⁺) m/z : 357 (M⁺, 3), 116 (100). HRMS calcd for C₂₃H₁₉NO₃: 357.1365, Found: 357.1369.

(S)-N,N-dibenzyl-1-phenylprop-2-en-1-amine (3Ca, R¹=Bn, R²=Bn).

A colorless oil. IR (CHCl₃) 1493, 1451 cm⁻¹. ¹H NMR (CDCl₃) δ 7.48 (2H, d, J = 7.6 Hz), 7.40 (4H, d, J = 7.3 Hz), 7.35-7.15 (9H, m), 6.08 (1H, ddd, J = 17.1, 10.1, 8.5 Hz), 5.44 (1H, d, J = 10.1 Hz), 5.20 (1H, d, J = 17.1 Hz), 4.27 (1H, d, J = 8.5 Hz), 3.66 (2H, d, J = 14.0 Hz), 3.53 (2H, d, J = 14.0 Hz). ¹³C NMR (CDCl₃) δ 141.5, 140.1, 135.4, 128.7, 128.3, 128.2 (2C), 127.0, 126.8, 119.4, 65.2, 53.6. MS (EI⁺) m/z : 313 (M⁺, 22), 117 (100). HRMS calcd for C₂₃H₂₃N: 313.1830, Found: 313.1824. HPLC (Chiralcel OJ-H, hexane/2-propanol=95/5, 0.5 ml/min 254 nm) t_r (S) = 10.8 min, t_r (R) = 14.0 min. A sample of 95%ee (S) by HPLC analysis gave $[\alpha]_D^{31}$ -111.0 (c 1.06, CHCl₃).

(E)-N,N-Dibenzyl-3-phenylprop-2-en-1-amine (4Ca, R¹=Bn, R²=Bn).

A colorless oil. IR (CHCl₃) 1494, 1450 cm⁻¹. ¹H NMR (CDCl₃) δ 7.41-7.15 (15H, m), 6.52 (1H, d, J = 15.9 Hz), 6.29 (1H, dt, J = 15.9, 11.4 Hz), 3.63 (4H, s), 3.22 (2H, d, J = 11.4 Hz). ¹³C NMR (CDCl₃) δ 139.7, 137.3, 132.5, 128.8, 128.5, 128.3, 127.8, 127.3, 126.9, 126.3, 57.9, 55.7. MS (EI⁺) m/z : 313 (M⁺, 38), 115 (100). HRMS calcd for C₂₃H₂₃N (M⁺): 313.1830, Found: 313.1836.

4-Methoxy-N-((S)-1-phenylallyl)benzenamine (3Da, R¹=4-MeO-C₆H₄, R²=H).

A colorless oil. IR (CHCl₃) 1512, 1457 cm⁻¹. ¹H NMR (CDCl₃) δ 7.42-7.21 (5H, m), 6.72 (2H, d, J = 8.9 Hz), 6.55 (2H, d, J = 8.9 Hz), 6.02 (1H, ddd, J = 17.1, 10.1, 5.8 Hz), 5.25 (1H, d, J = 17.1 Hz), 5.19 (1H, d, J = 10.1 Hz), 4.85 (1H, d, J = 5.8 Hz), 3.70 (3H, s). ¹³C NMR (CDCl₃) δ 152.3, 142.1, 141.3, 139.5, 128.7, 127.4, 127.2, 115.9, 115.0, 114.7, 61.8, 55.6. MS (EI⁺) m/z : 239 (M⁺, 58), 115 (100). HRMS calcd for C₁₆H₁₇NO: 239.1310, Found: 239.1303. HPLC (Chiralcel AD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) t_r (S) = 15.8 min, t_r (R) = 13.8 min. A sample of 88%ee (S) by HPLC analysis gave $[\alpha]_D^{28}$ -12.1 (c 0.95, CHCl₃).

***N*-Cinnamyl-4-methoxybenzenamine (4Da, R¹=4-MeO-C₆H₄, R²=H).**

A colorless oil. IR (CHCl₃) 1513, 1467 cm⁻¹. ¹H NMR (CDCl₃) δ 7.39-7.18 (5H, m), 6.79 (2H, d, *J* = 9.2 Hz), 6.63 (2H, d, *J* = 9.2 Hz), 6.61 (1H, d, *J* = 15.9 Hz), 6.33 (1H, dt, *J* = 15.9, 6.1 Hz), 3.88 (2H, d, *J* = 6.1 Hz), 3.74 (3H, s). ¹³C NMR (CDCl₃) δ 152.4, 142.3, 136.9, 131.5, 128.6, 127.5, 127.4, 126.3, 114.9, 114.4, 55.7, 47.2. MS (EI⁺) *m/z*: 239 (M⁺, 41), 117 (100). HRMS calcd for C₁₆H₁₇NO: 239.1310, Found: 239.1320.

***(E)*-(*S*)-*O*-(1-Phenylprop-2-enyl)benzaldehyde oxime (12Aa, R¹=Ph, R²=H, Ar=Ph).**

A colorless oil. IR (CHCl₃) 2908, 1494, 1450, 1414 cm⁻¹. ¹H NMR (CDCl₃) δ 8.18 (1H, s), 7.60-7.50 (2H, m), 7.45-7.23 (8H, m), 6.16 (1H, ddd, *J* = 17.1, 10.5, 6.3 Hz), 5.71 (1H, d, *J* = 6.3 Hz), 5.34 (1H, d, *J* = 17.1 Hz), 5.29 (1H, d, *J* = 10.5 Hz). ¹³C NMR (CDCl₃) δ 149.2, 140.1, 137.6, 132.3, 129.8, 128.6, 128.4, 127.9, 127.4, 127.1, 117.3. MS (EI⁺) *m/z*: 237 (M⁺, 0.2), 117 (100). HRMS calcd for C₁₆H₁₅NO: 237.1154, Found: 237.1156. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*S*) = 8.8 min, *t_r* (*R*) = 13.2 min. A sample of 95%ee (*S*) by HPLC analysis gave [α]_D²⁶ -62.3 (*c* 1.03, CHCl₃).

***(E)*-*O*-(3-Phenylprop-2-enyl)benzaldehyde oxime (13Aa, R¹=Ph, R²=H, Ar=Ph).**

A colorless oil. IR (CHCl₃) 2926, 1494, 1448 cm⁻¹. ¹H NMR (CDCl₃) δ 8.13 (1H, s), 7.63-7.48 (2H, m), 7.46-7.16 (8H, m), 6.67 (1H, d, *J* = 15.9 Hz), 6.42 (1H, dt, *J* = 15.9, 6.2 Hz), 4.83 (2H, d, *J* = 6.2 Hz). ¹³C NMR (CDCl₃) δ 148.9, 136.6, 133.5, 132.2, 129.8, 128.7, 128.5, 127.8, 127.1, 126.6, 125.1, 74.9. MS (EI⁺) *m/z*: 237 (M⁺, 1.5), 117 (100). HRMS calcd for C₁₆H₁₅NO: 237.1154, Found: 237.1155.

***(E)*-(*S*)-*O*-[1-(1-Naphthyl)prop-2-enyl]benzaldehyde oxime (12Ac, R¹=Ph, R²=H, Ar=1-Naph).**

A colorless oil. IR (CHCl₃) 2944, 1511, 1447 cm⁻¹. ¹H NMR (CDCl₃) δ 8.21 (1H, s), 8.20 (1H, d, *J* = 11.0 Hz), 7.85 (1H, d, *J* = 7.9 Hz), 7.80 (2H, d, *J* = 8.3 Hz), 7.61-7.44 (6H, m), 7.34-7.28 (3H, m), 6.43 (1H, d, *J* = 5.5 Hz), 6.35 (1H, ddd, *J* = 17.0, 9.2, 5.5 Hz), 5.37 (1H, d, *J* = 17.0 Hz), 5.34 (1H, d, *J* = 9.2 Hz). ¹³C NMR (CDCl₃) δ 149.3, 137.1, 135.6, 134.0, 132.3, 131.2, 129.8, 128.8, 128.7, 128.6, 127.2, 126.1, 125.6, 125.5, 125.3, 124.3, 117.8, 83.6. MS (EI⁺) *m/z*: 287 (M⁺, 0.2), 167 (100). HRMS calcd for C₂₀H₁₇NO: 287.1310, Found: 287.1313. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*S*) = 12.0 min, *t_r* (*R*) = 13.6 min. A sample of 90% ee (*S*) by HPLC analysis gave [α]_D²⁸ -17.1 (*c* 1.04, CHCl₃).

(*E*)-*O*-[3-(1-Naphthyl)prop-2-enyl]benzaldehyde oxime (13Ac, R¹=Ph, R²=H, Ar=1-Naph).

A colorless oil. IR (CHCl₃) 2925, 1507, 1493, 1446 cm⁻¹. ¹H NMR (CDCl₃) δ 8.17 (1H, s), 8.12 (1H, d, *J* = 7.9 Hz), 7.82 (1H, d, *J* = 6.4 Hz), 7.75 (1H, d, *J* = 8.2 Hz), 7.68-7.55 (3H, m), 7.51-7.30 (7H, m), 6.44 (1H, dt, *J* = 15.8, 6.1 Hz), 4.94 (2H, d, *J* = 6.1 Hz), 3.78 (3H, s). ¹³C NMR (CDCl₃) δ 149.0, 134.5, 133.6, 132.3, 131.2, 130.6, 129.9, 128.7, 128.5, 128.4, 128.1, 127.1, 126.1, 125.8, 125.6, 124.1, 123.8, 75.0. MS (EI⁺) *m/z*: 287 (M⁺, 10.1), 167 (100). HRMS calcd for C₂₀H₁₇NO: 287.1310, Found: 287.1311.

(*E*)-(*S*)-*O*-[1-(2-Naphthyl)prop-2-enyl]benzaldehyde oxime (12Ad, R¹=Ph, R²=H, Ar=2-Naph).

A colorless oil. IR (CHCl₃) 2944, 1507, 1446 cm⁻¹. ¹H NMR (CDCl₃) δ 8.21 (1H, s), 7.90-7.77 (4H, m), 7.58-7.42 (5H, m), 7.35-7.26 (3H, m), 6.24 (1H, ddd, *J* = 17.0, 10.5, 6.0 Hz), 5.87 (1H, d, *J* =

6.0 Hz), 5.38 (1H, d, $J = 17.0$ Hz), 5.32 (1H, d, $J = 10.5$ Hz). ^{13}C NMR (CDCl_3) δ 149.3, 137.7, 137.6, 133.3, 133.1, 132.3, 129.8, 128.6, 128.2, 128.1, 127.7, 127.2, 126.4, 126.1, 126.0, 125.3, 117.6, 86.2. MS (EI^+) m/z : 287 (M^+ , < 0.1), 167 (100). HRMS calcd for $\text{C}_{20}\text{H}_{17}\text{NO}$: 287.1310, Found: 287.1308. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) t_r (S) = 9.9 min, t_r (R) = 16.3 min. A sample of 89%ee (S) by HPLC analysis gave $[\alpha]_{\text{D}}^{28} -76.3$ (c 1.00, CHCl_3).

(*E*)-*O*-[3-(2-Naphthyl)prop-2-enyl]benzaldehyde oxime (13Ad, $\text{R}^1=\text{Ph}$, $\text{R}^2=\text{H}$, $\text{Ar}=2\text{-Naph}$).

A colorless crystal. mp 70-73 °C (hexane). IR (CHCl_3) 2927, 1506, 1446 cm^{-1} . ^1H NMR (CDCl_3) δ 8.16 (1H, s), 7.83-7.75 (4H, m), 7.66-7.58 (3H, m), 7.49-7.34 (5H, m), 6.85 (1H, d, $J = 15.9$ Hz), 6.56 (1H, dt, $J = 15.9, 6.1$ Hz), 4.90 (2H, d, $J = 6.1$ Hz). ^{13}C NMR (CDCl_3) δ 149.0, 134.1, 133.6, 133.1, 132.2, 129.9, 128.7 (2C), 128.2, 128.0, 127.7, 127.1, 126.7, 126.3, 126.0, 125.5, 123.6, 75.0. MS (EI^+) m/z : 287 (M^+ , 5.5), 167 (100). HRMS calcd for $\text{C}_{20}\text{H}_{17}\text{NO}$: 287.1310, Found: 287.1312. Anal. Calcd for $\text{C}_{20}\text{H}_{17}\text{NO}$: C, 83.59; H, 5.96; N, 4.87. Found: C, 83.72; H, 5.97; N, 4.83.

(*E*)-(*S*)-*O*-[1-(4-Fluorophenyl)prop-2-enyl]benzaldehyde oxime (12Ae, $\text{R}^1=\text{Ph}$, $\text{R}^2=\text{H}$, $\text{Ar}=4\text{-F-C}_6\text{H}_4$).

A colorless oil. IR (CHCl_3) 2906, 1509, 1447, 1415 cm^{-1} . ^1H NMR (CDCl_3) δ 8.17 (1H, s), 7.58-7.53 (2H, m), 7.40-7.32 (5H, m), 7.08-7.02 (2H, m), 6.13 (1H, ddd, $J = 17.0, 10.6, 6.2$ Hz), 5.68 (1H, d, $J = 6.2$ Hz), 5.32 (1H, d, $J = 17.0$ Hz), 5.31 (1H, d, $J = 10.6$ Hz). ^{13}C NMR (CDCl_3) δ 162.4 ($J = 246$ Hz), 149.3, 137.4, 136.0, 132.2, 129.9, 129.1 ($J = 7.2$ Hz), 128.6, 127.1, 117.5, 115.3 ($J = 20.7$ Hz), 85.3. MS (EI^+) m/z : 255 (M^+ , 0.2), 135 (100). HRMS calcd for $\text{C}_{16}\text{H}_{14}\text{FNO}$: 255.1059, Found: 255.1065. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) t_r (S) = 8.6 min, t_r (R) = 12.8 min. A sample of 90%ee (S) by HPLC analysis gave $[\alpha]_{\text{D}}^{28} -53.7$ (c 1.04, CHCl_3).

(*E*)-*O*-[3-(4-Fluorophenyl)prop-2-enyl]benzaldehyde oxime (13Ae, $\text{R}^1=\text{Ph}$, $\text{R}^2=\text{H}$, $\text{Ar}=4\text{-F-C}_6\text{H}_4$).

A colorless crystal. mp 54-57 °C (hexane). IR (CHCl₃) 2926, 1509, 1447, 1414 cm⁻¹. ¹H NMR (CDCl₃) δ 8.14 (1H, s), 7.63-7.57 (2H, m), 7.42-7.34 (5H, m), 7.05-6.97 (2H, m), 6.65 (1H, d, *J* = 15.8 Hz), 6.34 (1H, dt, *J* = 15.8, 6.1 Hz), 4.82 (2H, d, *J* = 6.1 Hz). ¹³C NMR (CDCl₃) δ 162.5 (*J* = 247 Hz), 149.0, 132.8, 132.3, 132.2, 129.8, 128.7, 128.1 (*J* = 8.3 Hz), 127.1, 124.9, 115.4 (*J* = 21.7 Hz), 74.7. MS (EI⁺) *m/z*: 255 (M⁺, 0.5), 135 (100). HRMS calcd for C₁₆H₁₄FNO: 255.1059, Found: 255.1062. Anal. Calcd for C₁₆H₁₄FNO: C, 75.28; H, 5.53; N, 5.49; F, 7.44. Found: C, 75.33; H, 5.72; N, 5.45; F, 7.46.

(*E*)-(S)-O-[1-(4-Methylphenyl)prop-2-enyl]benzaldehyde oxime (12Af, R¹=Ph, R²=H, Ar=4-Me-C₆H₄).

A colorless oil. IR (CHCl₃) 2945, 1513, 1493, 1448, 1413 cm⁻¹. ¹H NMR (CDCl₃) δ 8.17 (1H, s), 7.58-7.52 (2H, m), 7.35-7.32 (3H, m), 7.29 (2H, d, *J* = 7.9 Hz), 7.18 (2H, d, *J* = 7.9 Hz), 6.16 (1H, ddd, *J* = 17.4, 10.5, 6.3 Hz), 5.67 (1H, d, *J* = 6.3 Hz), 5.33 (1H, d, *J* = 17.4 Hz), 5.28 (1H, d, *J* = 10.5 Hz), 2.34 (3H, s). ¹³C NMR (CDCl₃) δ 149.1, 137.8, 137.6, 137.2, 132.4, 129.8, 129.1, 128.6, 127.4, 127.1, 117.1, 86.0, 21.1. MS (EI⁺) *m/z*: 251 (M⁺, < 0.1), 131 (100). HRMS calcd for C₁₇H₁₇NO: 251.1310, Found: 251.1306. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*S*) = 8.0 min, *t_r* (*R*) = 11.0 min. A sample of 90% ee (*S*) by HPLC analysis gave [α]_D²⁸ -72.9 (*c* 1.01, CHCl₃).

(*E*)-O-[3-(4-Methylphenyl)prop-2-enyl]benzaldehyde oxime (13Af, R¹=Ph, R²=H, Ar=4-Me-C₆H₄).

A colorless crystal. mp 70-72 °C (hexane). IR (CHCl₃) 2925, 1512, 1493, 1448, 1414 cm⁻¹. ¹H NMR (CDCl₃) δ 8.14 (1H, s), 7.63-7.56 (2H, m), 7.40-7.34 (3H, m), 7.31 (2H, d, *J* = 7.9 Hz), 7.12 (2H, d, *J* = 7.9 Hz), 6.65 (1H, d, *J* = 15.9 Hz), 6.38 (1H, dt, *J* = 15.9, 6.4 Hz), 4.83 (2H, d, *J* = 6.4

(Hz), 2.34 (3H, s). ¹³C NMR (CDCl₃) 148.9, 137.7, 133.9, 133.5, 132.3, 129.8, 129.3, 128.7, 127.1, 126.5, 124.0, 75.0, 21.1. MS (EI⁺) m/z: 251 (M⁺, 1.4), 131 (100). HRMS calcd for C₁₇H₁₇NO: 251.1310, Found: 251.1314. Anal. Calcd for C₁₇H₁₇NO: C, 81.24; H, 6.82; N, 5.57. Found: C, 81.54; H, 6.92; N, 5.53.

(*E*)-(*S*)-*O*-(1-Phenylprop-2-enyl)-4-trifluoromethylbenzaldehyde oxime (12Ba), R¹=4-F₃C-C₆H₄, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2907, 1494, 1453, 1412 cm⁻¹. ¹H NMR (CDCl₃) δ 8.20 (1H, s), 7.67 (2H, d, *J* = 7.9 Hz), 7.58 (2H, d, *J* = 7.9 Hz), 7.44-7.30 (5H, m), 6.16 (1H, ddd, *J* = 17.1, 10.5, 6.3 Hz), 5.72 (1H, d, *J* = 6.3 Hz), 5.34 (1H, d, *J* = 17.1 Hz), 5.31 (1H, d, *J* = 10.5 Hz). ¹³C NMR (CDCl₃) δ 147.7, 139.9, 137.4, 135.7, 131.4 (*J* = 32.0 Hz), 128.5, 128.0, 127.4, 127.3, 125.6, 124.0 (*J* = 272 Hz), 117.6, 86.6. MS (EI⁺) m/z: 306 (M + H⁺, < 0.1), 117 (100). HRMS calcd for C₁₇H₁₅F₃NO (M + H⁺): 306.1104, Found: 306.1099. HPLC (Chiralcel OD-H, hexane/2-propanol=99/1, 0.5 ml/min 254 nm) t_r (*R*) = 12.1 min, t_r (*S*) = 13.4 min. A sample of 92% ee (*S*) by HPLC analysis gave [α]_D²⁷ -53.3 (*c* 0.98, CHCl₃).

(*E*)-*O*-(3-Phenylprop-2-enyl)-4-trifluoromethylbenzaldehyde oxime (13Ba), R¹=4-F₃C-C₆H₄, R²=H, Ar=Ph).

A colorless crystal. mp 90-93 °C (hexane). IR (CHCl₃) 2928, 1494, 1450, 1413 cm⁻¹. ¹H NMR (CDCl₃) δ 8.15 (1H, s), 7.71 (2H, d, *J* = 8.1 Hz), 7.60 (2H, d, *J* = 8.1 Hz), 7.42 (2H, d, *J* = 7.6 Hz), 7.36-7.23 (3H, m), 6.70 (1H, d, *J* = 15.9 Hz), 6.42 (1H, dt, *J* = 15.9, 6.4 Hz), 4.86 (2H, d, *J* = 6.4 Hz). ¹³C NMR (CDCl₃) δ 147.4, 136.5, 135.7, 133.8, 131.4 (*J* = 32.1 Hz), 128.6, 128.0, 127.2, 126.6, 125.6, 124.8, 124.0 (*J* = 272 Hz), 75.3. MS (EI⁺) m/z: 305 (M⁺, 1.1), 117 (100). HRMS calcd for C₁₇H₁₄F₃NO: 305.1027, Found: 305.1023. Anal. Calcd for C₁₇H₁₄F₃NO: C, 66.88; H, 4.62; N, 4.59; F, 18.67. Found: C, 67.16; H, 4.81; N, 4.59; F, 18.67.

(*E*)-(*S*)-*O*-(1-Phenylprop-2-enyl)-4-methoxybenzaldehyde oxime (12Ca), R¹=4-MeO-C₆H₄, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2936, 1513, 1459, 1418 cm⁻¹. ¹H NMR (CDCl₃) δ 8.13 (1H, s), 7.49 (2H, d, *J* = 8.6 Hz), 7.42-7.26 (5H, m), 6.86 (2H, d, *J* = 8.6 Hz), 6.15 (1H, ddd, *J* = 17.3, 10.5, 6.4 Hz), 5.67 (1H, d, *J* = 6.4 Hz), 5.32 (1H, d, *J* = 17.3 Hz), 5.28 (1H, d, *J* = 10.5 Hz), 3.80 (3H, s). ¹³C NMR (CDCl₃) δ 161.0, 148.9, 140.3, 137.8, 128.6, 128.4, 127.8, 127.4, 125.0, 117.2, 114.1, 85.9, 55.3. MS (EI⁺) *m/z*: 267 (M⁺, 6.4), 117 (100). HRMS calcd for C₁₇H₁₇NO₂: 267.1259, Found: 267.1261. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*S*) = 10.2 min, *t_r* (*R*) = 13.1 min. A sample of 93%ee (*S*) by HPLC analysis gave [α]_D²⁷ -64.4 (*c* 0.99, CHCl₃).

(*E*)-O-(3-Phenylprop-2-enyl)-4-methoxybenzaldehyde oxime (13Ca, R¹=4-MeO-C₆H₄, R²=H, Ar=Ph).

A colorless crystal. mp 72-75 °C (hexane). IR (CHCl₃) 2935, 1513, 1463, 1419 cm⁻¹. ¹H NMR (CDCl₃) δ 8.09 (1H, s), 7.53 (2H, d, *J* = 8.6 Hz), 7.40 (2H, d, *J* = 7.5 Hz), 7.31 (2H, d, *J* = 7.5 Hz), 7.23 (1H, m), 6.88 (2H, d, *J* = 8.6 Hz), 6.69 (1H, d, *J* = 15.9 Hz), 6.43 (1H, dt, *J* = 15.9, 6.2 Hz), 4.81 (2H, d, *J* = 6.2 Hz), 3.82 (3H, s). ¹³C NMR (CDCl₃) δ 161.0, 148.6, 136.7, 133.3, 128.5 (2C), 127.8, 126.6, 125.3, 124.9, 114.1, 74.7, 55.2. MS (EI⁺) *m/z*: 267 (M⁺, 10.3), 117 (100). HRMS calcd for C₁₇H₁₇NO₂: 267.1259, Found: 267.1260. Anal. Calcd for C₁₇H₁₇NO₂: C, 76.38; H, 6.41; N, 5.24. Found: C, 76.52; H, 6.39; N, 5.22.

Methyl (*E*)-(*S*)-2-[(1-Phenylprop-2-enyloxy)imino]ethanate (12Da, R¹=CO₂Me, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2955, 1739, 1494, 1442, 1415 cm⁻¹. ¹H NMR (CDCl₃) δ 7.60 (1H, s), 7.40-7.28 (5H, m), 6.12 (1H, ddd, *J* = 17.1, 10.7, 6.2 Hz), 5.81 (1H, d, *J* = 6.2 Hz), 5.34-5.28 (2H, m), 3.82 (3H, s). ¹³C NMR (CDCl₃) δ 162.4, 141.1, 138.7, 136.4, 128.6, 128.3, 127.4, 118.2, 87.9, 52.4. MS (EI⁺) *m/z*: 219 (M⁺, < 0.1), 117 (100). HRMS calcd for C₁₂H₁₃NO₃: 219.0895, Found: 219.0893. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*R*) = 13.9 min, *t_r* (*S*) = 15.4 min. A sample of 89%ee (*S*) by HPLC analysis gave [α]_D²⁶ -1.7 (*c* 1.05, CHCl₃).

Methyl (*E*)-2-[(3-Phenylprop-2-enyloxy)imino]ethanate (13Da, R¹=CO₂Me, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2954, 1727, 1494, 1442 cm⁻¹. ¹H NMR (CDCl₃) δ 7.52 (1H, s), 7.40 (2H, d, *J* = 7.3 Hz), 7.32 (2H, t, *J* = 7.3 Hz), 7.26 (1H, br t, *J* = 7.3 Hz), 6.67 (1H, d, *J* = 15.9 Hz), 6.36 (1H, dt, *J* = 15.9, 6.6 Hz), 4.91 (2H, d, *J* = 6.6 Hz), 3.86 (3H, s). ¹³C NMR (CDCl₃) δ 162.4, 140.8, 136.2, 134.7, 128.6, 128.1, 126.7, 123.4, 76.6, 52.4. MS (EI⁺) *m/z*: 219 (M⁺, 5.7), 117 (100). HRMS calcd for C₁₂H₁₃NO₃: 219.0895, Found: 219.0897.

(*E*)-(*S*)-*O*-(1-Phenylprop-2-enyl)-2-pyridinecarboxaldehyde oxime (12Ea, R¹=2-Pyridyl, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2941, 1493, 1471, 1454, 1435 cm⁻¹. ¹H NMR (CDCl₃) δ 8.59 (1H, d, *J* = 4.9 Hz), 8.28 (1H, s), 7.76 (1H, d, *J* = 7.9 Hz), 7.63 (1H, m), 7.45-7.18 (6H, m), 6.17 (1H, ddd, *J* = 17.1, 10.5, 6.2 Hz), 5.76 (1H, d, *J* = 6.2 Hz), 5.34 (1H, d, *J* = 17.1 Hz), 5.31 (1H, d, *J* = 10.5 Hz). ¹³C NMR (CDCl₃) δ 151.8, 150.0, 149.6, 139.9, 137.3, 136.3, 128.4, 127.9, 127.4, 123.9, 121.0, 117.5, 86.6. MS (EI⁺) *m/z*: 238 (M⁺, 0.6), 117 (100). HRMS calcd for C₁₅H₁₄N₂O: 238.1106, Found: 238.1111. HPLC (Chiralcel OD-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t*_r (*S*) = 10.8 min, *t*_r (*R*) = 12.4 min. A sample of 76%ee (*S*) by HPLC analysis gave [α]_D²⁶ -45.2 (*c* 0.98, CHCl₃).

(*E*)-*O*-(3-Phenylprop-2-enyl)-2-pyridinecarboxaldehyde oxime (13Ea, R¹=2-Pyridyl, R²=H, Ar=Ph).

A colorless oil. IR (CHCl₃) 2928, 1494, 1470, 1436 cm⁻¹. ¹H NMR (CDCl₃) δ 8.61 (1H, d, *J* = 4.0 Hz), 8.23 (1H, s), 7.80 (1H, d, *J* = 7.9 Hz), 7.67 (1H, m), 7.41 (2H, d, *J* = 7.6 Hz), 7.32 (2H, d, *J* = 7.6 Hz), 7.27-7.20 (2H, m), 6.69 (1H, d, *J* = 15.9 Hz), 6.42 (1H, dt, *J* = 15.9, 6.4 Hz), 4.89 (2H, d, *J* = 6.4 Hz). ¹³C NMR (CDCl₃) δ 151.7, 149.7, 149.6, 136.5, 136.4, 133.7, 128.6, 127.9, 126.6, 124.8, 124.0, 121.1, 75.4. MS (EI⁺) *m/z*: 238 (M⁺, 2.4), 117 (100). HRMS calcd for C₁₅H₁₄N₂O: 238.1106,

Found: 238.1114.

(S)-O-(1-Phenylprop-2-enyl)benzophenone oxime (12Fa, R¹=Ph, R²=Ph, Ar=Ph).

A colorless oil. IR (CHCl₃) 1493, 1447, 1414 cm⁻¹. ¹H NMR (CDCl₃) δ 7.48-7.21 (15H, m), 6.07 (1H, ddd, *J* = 16.7, 10.3, 6.0 Hz), 5.74 (1H, d, *J* = 6.0 Hz), 5.23 (1H, d, *J* = 16.7 Hz), 5.20 (1H, d, *J* = 10.3 Hz). ¹³C NMR (CDCl₃) δ 157.2, 140.6, 138.0, 136.6, 133.5, 129.4, 129.2, 128.7, 128.3, 128.2, 128.1, 128.0, 127.5, 127.1, 116.7, 86.3. MS (EI⁺) *m/z*: 313 (M⁺, 2.5), 117 (100). HRMS calcd for C₂₂H₁₉NO: 313.1467, Found: 313.1466. HPLC (Chiralcel OJ-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*R*) = 12.4 min, *t_r* (*S*) = 13.6 min. A sample of 94%ee (*S*) by HPLC analysis gave [α]_D²⁶ +8.3 (*c* 0.99, CHCl₃).

O-(3-Phenylprop-2-enyl)benzophenone oxime (13Fa, R¹=Ph, R²=Ph, Ar=Ph).

A colorless crystal. mp 66-68 °C (hexane). IR (CHCl₃) 2924, 1494, 1444 cm⁻¹. ¹H NMR (CDCl₃) δ 7.55-7.20 (15H, m), 6.61 (1H, d, *J* = 16.1 Hz), 6.41 (1H, dt, *J* = 16.1, 6.1 Hz), 4.86 (2H, d, *J* = 6.1 Hz). ¹³C NMR (CDCl₃) δ 157.0, 136.8, 136.6, 133.4, 132.9, 129.4, 129.3, 128.9, 128.6, 128.2, 128.1, 128.0, 127.7, 126.6, 125.8, 75.2. MS (EI⁺) *m/z*: 313 (M⁺, 3.7), 117 (100). HRMS calcd for C₂₂H₁₉NO: 313.1467, Found: 313.1469. Anal. Calcd for C₂₂H₁₉NO: C, 84.31; H, 6.11; N, 4.47. Found: C, 84.44; H, 6.16; N, 4.43.

(S)-O-(1-Phenylprop-2-enyl)cyclohexanone oxime (12Ga, R¹, R²=(CH₂)₅, Ar=Ph).

A colorless oil. IR (CHCl₃) 2938, 1493, 1450 cm⁻¹. ¹H NMR (CDCl₃) δ 7.39-7.22 (5H, m), 6.07 (1H, ddd, *J* = 17.1, 10.7, 6.1 Hz), 5.54 (1H, d, *J* = 6.1 Hz), 5.25 (1H, d, *J* = 17.1 Hz), 5.21 (1H, d, *J* = 10.7 Hz), 2.60-2.51 (2H, m), 2.23-2.13 (2H, m), 1.70-1.53 (6H, m). ¹³C NMR (CDCl₃) δ 161.1, 140.9, 138.5, 128.3, 127.5, 127.0, 116.3, 84.8, 32.1, 27.0, 25.8, 25.7, 25.6. MS (EI⁺) *m/z*: 229 (M⁺, 0.2), 117 (100). HRMS calcd for C₁₅H₁₉NO: 229.1467, Found: 229.1466. HPLC (Chiralcel OJ-H, hexane/2-propanol=90/10, 0.5 ml/min 254 nm) *t_r* (*R*) = 11.0 min, *t_r* (*S*) = 12.6 min. A sample of 73%ee (*S*) by HPLC analysis gave [α]_D²⁶ +41.6 (*c* 1.12, CHCl₃).

O-(3-Phenylprop-2-enyl)cyclohexanone oxime (13Ga, R¹, R²=(CH₂)₅, Ar=Ph).

A colorless oil. IR (CHCl₃) 2938, 1495, 1449 cm⁻¹. ¹H NMR (CDCl₃) δ 7.39 (2H, d, *J* = 7.6 Hz), 7.30 (2H, d, *J* = 7.6 Hz), 7.22 (1H, br t, *J* = 7.6 Hz), 6.62 (1H, d, *J* = 15.9 Hz), 6.37 (1H, dt, *J* = 15.9, 6.0 Hz), 4.68 (2H, d, *J* = 6.0 Hz), 2.50 (2H, br t, *J* = 6.1 Hz), 2.21 (2H, br t, *J* = 6.1 Hz), 1.73-1.52 (6H, m). ¹³C NMR (CDCl₃) δ 160.5, 136.9, 132.5, 128.5, 127.6, 126.5, 125.9, 73.8, 32.1, 26.9, 25.7, 25.6, 25.3. MS (EI⁺) *m/z*: 229 (M⁺, 2.1), 117 (100). HRMS calcd for C₁₅H₁₉NO: 229.1467, Found: 229.1467.

3. Selective Reduction of Oxime Ether (12Aa)

Alcohol 14

A suspension of oxime ether **12Aa** (119 mg, 0.50 mmol) and 20% Pd(OH)₂-C (30 mg) in MeOH (2.5 mL) was stirred under a hydrogen atmosphere at 20 °C for 3 h. After the reaction mixture was filtered, the filtrate was concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=5:1) afforded the product **14** (47.6 mg, 70%). A colorless oil. IR (CHCl₃) 3428, 2969, 1493, 1455, 1380 cm⁻¹. ¹H NMR (CDCl₃) δ 7.37-7.20 (5H, m), 4.55 (1H, t, *J* = 6.6 Hz), 2.12 (1H, br s), 1.84-1.67 (2H, m), 0.89 (3H, t, *J* = 7.5 Hz). ¹³C NMR (CDCl₃) δ 144.6, 128.4, 127.4, 126.0, 75.9, 31.7, 10.0. MS (EI⁺) *m/z*: 136 (M⁺, 9.0), 91 (100). HRMS calcd for C₉H₁₂O: 136.0888, Found: 136.0888. A sample of 95%ee gave [α]_D²⁶ -49.0 (*c* 1.01, hexane).

Oxime Ether 15

A suspension of oxime ether **12Aa** (71.2 mg, 0.30 mmol) and 10% Pd-C (21.4 mg) in AcOEt (3.0 mL) was stirred under a hydrogen atmosphere at 20 °C for 10 min. After the reaction mixture was filtered, the filtrate was concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane:AcOEt=15:1) afforded the product **15** (53.3 mg, 74%). A colorless oil. IR (CHCl₃) 2971, 1494, 1451 cm⁻¹. ¹H NMR (CDCl₃) δ 8.15 (1H, s), 7.56-7.51 (2H, m), 7.35-7.25 (8H, m), 5.11 (1H, t, *J* = 6.7 Hz), 2.05 (1H, m), 1.86 (1H, m), 0.95 (3H, t, *J* = 7.5 Hz). ¹³C NMR (CDCl₃) δ 148.6,

142.1, 132.5, 129.6, 128.6, 128.2, 127.4, 127.0, 126.9, 86.9, 28.9, 10.0. MS (EI^+) m/z : 239 (M^+ , 2.5), 119 (100). HRMS calcd for $\text{C}_{16}\text{H}_{17}\text{NO}$: 239.1310, Found: 239.1314. A sample of 95%ee gave $[\alpha]_{\text{D}}^{26} -83.2$ (c 1.20, CHCl_3).

Alcohol **16**

To a suspension of oxime ether **12Aa** (47.5 mg, 0.20 mmol) in $\text{AcOH-H}_2\text{O}$ (1:1 v/v, 1.2 mL) was added Zn powder (523 mg, 8.0 mmol) under argon atmosphere at 20 °C. After being stirred at the same temperature for 2 h, the reaction mixture was diluted with Et_2O . After the reaction mixture was filtered and then extracted with Et_2O . The organic phase was dried over MgSO_4 and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane: AcOEt =8:1) afforded the product **16** (22.6 mg, 84%). A colorless oil. IR (CHCl_3) 3427, 1493, 1452, 1431 cm^{-1} . ^1H NMR (CDCl_3) δ 7.39-7.25 (5H, m), 6.05 (1H, ddd, J = 17.1, 10.6, 6.0 Hz), 5.35 (1H, d, J = 17.1 Hz), 5.23-5.15 (2H, m). ^{13}C NMR (CDCl_3) δ 142.6, 140.2, 128.6, 127.8, 126.3, 115.1, 75.3. MS (EI^+) m/z : 134 (M^+ , 64), 133 (100). HRMS calcd for $\text{C}_9\text{H}_{10}\text{O}$: 134.0732, Found: 134.0731. A sample of 95%ee gave $[\alpha]_{\text{D}}^{23} -8.2$ (c 1.23, benzene).

Hydroxylamine **17**

To a solution of oxime ether **12Aa** (119 mg, 0.50 mmol) in Et_2O (1.0 mL) was added borane-pyridine complex (0.25 mL, 2.5 mmol) under argon atmosphere at 0 °C. And then, 2 M HCl in Et_2O (2.0 mL) added to the reaction mixture over a period of 10 min under argon atmosphere at 0 °C. After being stirred at 20 °C for 3 h, the reaction mixture was diluted with saturated NaHCO_3 . The reaction mixture was extracted with CHCl_3 . The organic phase was dried over MgSO_4 and concentrated at reduced pressure. Purification of the residue by preparative TLC (hexane: AcOEt =5:1) afforded the product **17** (96.4 mg, 81%). A colorless oil. IR (CHCl_3) 3258, 1495, 1452, 1420 cm^{-1} . ^1H NMR (CDCl_3) δ 7.40-7.23 (10H, m), 5.97 (1H, ddd, J = 17.1, 10.7, 6.7 Hz), 5.63 (1H, br s), 5.23-5.16 (2H, m), 5.03 (1H, d, J = 6.7 Hz), 4.08 (1H, d, J = 12.8 Hz), 4.04 (1H, d, J = 12.8 Hz). ^{13}C NMR (CDCl_3) δ 140.6, 138.0, 137.5, 129.0, 128.3 (2C), 127.7, 127.4, 127.2, 116.9, 85.9, 56.5. MS (EI^+) m/z : 239 (M^+ , 0.3), 117 (100). HRMS calcd for $\text{C}_{16}\text{H}_{17}\text{NO}$: 239.1310, Found: 239.1309. A sample of 95%ee gave $[\alpha]_{\text{D}}^{26} -10.8$ (c 0.99, CHCl_3).

S35

S37

S41

S45

S47

No.	Chemical Shift (ppm)	Integration	Type	Integration
1	8.17	8.15	d	1
2	8.12	8.05	d	1
3	7.82	7.79	d	1
4	7.75	7.72	d	1
5	7.60	7.56	m	3
6	7.45	7.38	m	4
7	7.35	7.32	m	3
8	6.44	6.38	d	1
9	6.44	6.38	d	1
10	6.44	6.38	d	1
11	4.91	4.91	d	2

No.	Chemical Shift (ppm)	Integration	Type	Integration
1	8.17	8.15	d	1
2	8.12	8.05	d	1
3	7.82	7.79	d	1
4	7.75	7.72	d	1
5	7.60	7.56	m	3
6	7.45	7.38	m	4
7	7.35	7.32	m	3
8	6.44	6.38	d	1
9	6.44	6.38	d	1
10	6.44	6.38	d	1
11	4.91	4.91	d	2

S65

S85