Supporting Information

Zn-Promoted Regio- and Sequence-Selective One-Pot Joining Reaction of Three Components, Alkyl Iodides, α,β -Unsaturated Esters (or Nitriles) and Acylating Agents

Yoshimasa Yamamoto, Seiji Nakano, Hirofumi Maekawa, and Ikuzo Nishiguchi*

Department of Chemistry, Nagaoka University of Technology, 1603-1, Kamitomioka-cho, Nagaoka, Niigata 940-2188, Japan.

nishiiku@vos.nagaokaut.ac.jp

General methods.

Benzyl methacrylate (**1a**), methacrylonitrile (**1e**), alkyl iodides (**2**) and acid anhydrides (**6**) were purchased from Aldrich Chemical Co. Ltd, Tokyo Kasei Kogyo Co. Ltd or Nacalai Tesque Ltd, and were used without further purification. The compounds, 2-benzyloxycarbonyl-1-butene (**1b**) and 2-benzyloxycarbonyl-1-pentene (**1c**) were prepared by the reported method^[a]. Benzyl acrylate (**1d**) was prepared by the treatment of acrylic acid and benzyl bromide in the presence of potassium carbonate in acetonitrile at the refluxing temperature for 2 hours. 3-Iodopropionitrile (**7a**) was prepared by the iodination of 3-hydroxypropionitrile in the presence of sodium iodide and trimethylsilyl chloride in acetonitrile at the refluxing temperature for 2 hours. 4-Iodobutyronitrile (**7b**) was prepared by the iodination of 4-bromobutyronitrile in the presence of sodium iodide in acetone at the refluxing temperature for 2 hours. DMF, acetonitrile, propionitrile and butyronitrile were dried on CaH_2 and then was distilled.

^1H - and ^{13}C -NMR spectra were measured by a JEOL JNM-EX-400 (400 and 100MHz) spectrometer or a JEOL JNM-GX-270 (270 and 67.5MHz) spectrometer, respectively, with TMS as an internal standard. IR spectra were measured by a JASCO IR-810 spectrometer or JASCO FT/IR-400 Plus spectrometer. Mass spectra were obtained using a Shimadzu GC-MS QP-2000A or a JEOL JMS-600H. Gas chromatographic analysis was conducted through a Shimadzu GC-14B with FID using a capillary column packed with DB-17. The apparatus of elemental analysis was Yanaco CHN CORDER MT-6.

Typical procedure for Zn-promoted three components joining reaction of α,β -unsaturated esters (or nitriles), alkyl iodides and nitriles as the acylating agents is as follows: Into a acetonitrile (20 ml) solution containing commercially available Zn plate (60 mmol) was added dropwise a acetonitrile solution (40 ml) of α,β -unsaturated esters (or nitriles) (**1**, 10 mmol) and alkyl iodides (**2**, 100 mmol) at 20-25°C with magnetic

stirring under nitrogen atmosphere during a period of 45 min. in a water bath. After about 5 h., the color of mixture changed to dark gray. When the color of mixture did not change, 3mmol of trimethylsilyl chloride was added in order to activation of zinc. The mixture was stirred at 20-25°C for additional about 10 h. After the reaction, the reaction mixture was poured into 300 ml of 1% HCl solution and was extracted with three 100 ml portions of diethyl ether. The combined ethereal solution are washed with three 50 ml portions of saturated NaHCO₃ solution, two 50 ml portions of 0.1 N Na₂S₂O₃ solution and three 50 ml portions of saturated NaCl solution, and then dried over anhydrous MgSO₄. After removal of MgSO₄ by filtration and evaporation, the isolation by column chromatography gave the corresponding α,α -dialkylketoesters (**3**).

All the new compounds among the products were identified by spectroscopic and elemental analysis, and others were characterized by comparison of their chromatographic and spectroscopic behaviors with those of the authentic samples, as shown below.

Benzyl 2-ethyl-2-methyl-3-oxobutanoate (3a)

¹H-NMR(400MHz, CDCl₃) δ 0.81(t, J =7.6Hz, 3H), 1.33(s, 3H), 1.82(dq, J =14.0Hz, J =7.6Hz, 1H), 1.96(dq, J =14.0Hz, J =7.6Hz, 1H), 2.06(s, 3H), 5.16(s, 2H) and 7.30-7.36(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 8.54, 18.17, 26.10, 27.66, 60.02, 66.80, 128.08, 128.20, 128.39, 135.23, 172.67 and 205.22 ppm. IR(neat): 3070, 3030, 2970, 2940, 2880, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 234(M⁺). Anal.Calcd.for C₁₄H₁₈O₃: C, 71.77; H, 7.74. Found: C, 71.94; H, 8.03.

Benzyl 2-acetyl-2-methylpentanoate (3b)

¹H-NMR(400MHz, CDCl₃) δ 0.90(t, J =7.0Hz, 3H), 1.12-1.19(m, 2H), 1.34(s, 3H), 1.70-1.77(m, 1H), 1.84-1.92(m, 1H), 2.06(s, 3H), 5.16(s, 2H) and 7.26-7.33(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 14.42, 17.51, 18.72, 26.05, 36.93, 59.73, 66.79, 128.07, 128.20, 128.39, 135.23, 172.61 and 205.20 ppm. IR(neat): 3070, 3030, 2960, 2870, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 248(M⁺). Anal.Calcd.for C₁₅H₂₀O₃: C, 72.55; H, 8.12. Found: C, 72.29; H, 8.15.

Benzyl 2-acetyl-2-methylhexanoate (3c)

¹H-NMR(400MHz, CDCl₃) δ 0.85(t, J =7.2Hz, 3H), 1.05-1.14(m, 2H), 1.29(sext, J =7.2Hz, 2H), 1.33(s, 3H), 1.71-1.79(m, 1H), 1.86-1.93(m, 1H), 2.06(s, 3H), 5.15(d, J =12.4Hz, 1H), 5.18(d, J =12.4Hz, 1H) and 7.29-7.37(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 13.84, 18.73, 23.03, 26.08, 26.28, 34.48, 59.68, 66.81, 128.15, 128.23, 128.40, 135.25, 172.66 and 205.28 ppm. IR(neat): 3070, 3030, 2960, 2870, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 262(M⁺). Anal.Calcd.for C₁₆H₂₂O₃: C, 73.25; H, 8.45. Found: C, 73.06; H, 8.62.

Benzyl 2-acetyl-2, 4-dimethylpentanoate (3d)

¹H-NMR(400MHz, CDCl₃) δ 0.83(d, J =6.8Hz, 3H), 0.88(d, J =6.8Hz, 3H), 1.37(s, 3H), 1.54-1.68(m, 1H), 1.75(dd, J =14.4Hz, J =5.6Hz, 1H), 1.91(dd, J =14.4Hz, J =7.2Hz, 1H), 2.05(s, 3H), 5.13(d, J =12.0Hz, 1H), 5.18(d, J =12.0Hz, 1H) and 7.26-7.37(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 18.88, 23.42, 24.25, 24.49, 26.04, 42.85, 59.62, 66.98, 128.27, 128.29, 128.44, 135.11, 173.02 and 205.39 ppm. IR(neat): 3030, 2960, 2870, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 262(M⁺). Anal.Calcd.for C₁₆H₂₂O₃: C, 73.25; H, 8.45. Found: C, 72.96; H, 8.42.

Benzyl 2-acetyl-2-methylheptanoate (3e)

¹H-NMR(400MHz, CDCl₃) δ 0.85(t, J =6.8Hz, 3H), 1.07-1.27(m, 6H), 1.33(s, 3H), 1.70-1.78(m, 1H), 1.85-1.93(m, 1H), 2.06(s, 3H), 5.14(d, J =12.2Hz, 1H), 5.18(d, J =12.2Hz, 1H) and 7.27-7.36(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 13.95, 18.78, 22.37, 23.83, 26.10, 32.11, 34.77, 59.77, 66.85, 128.17, 128.25, 128.44, 135.31, 172.69

and 205.29 ppm. IR(neat): 3090, 3070, 3030, 2960, 2870, 1740, 1710 and 1500 cm^{-1} . EI-MS: m/z 276(M^+). Anal.Calcd.for $\text{C}_{17}\text{H}_{24}\text{O}_3$: C, 73.88; H, 8.75. Found: C, 73.79; H, 8.80.

Benzyl 2-(cyclohexylmethyl)-2-methyl-3-oxobutanoate (3f)

^1H -NMR(400MHz, CDCl_3) δ 0.84-1.25(m, 6H), 1.35(s, 3H), 1.50-1.62(m, 5H), 1.69(dd, $J=14.4\text{Hz}$, $J=5.6\text{Hz}$, 1H), 1.88(dd, $J=14.4\text{Hz}$, $J=6.8\text{Hz}$, 1H), 5.13(d, $J=12.2\text{Hz}$, 1H), 5.16(d, $J=12.2\text{Hz}$, 1H) and 7.30-7.37(m, 5H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 19.03, 26.00, 26.05, 26.24, 26.28, 33.82, 34.03, 34.71, 41.64, 59.41, 66.90, 128.24, 128.29, 128.39, 135.11, 172.98 and 205.35 ppm. IR(neat): 3066, 3030, 2920, 2850, 1740, 1710 and 1500 cm^{-1} . EI-MS: m/z 302(M^+). Anal.Calcd.for $\text{C}_{19}\text{H}_{26}\text{O}_3$: C, 75.46; H, 8.67. Found: C, 75.20; H, 8.63.

Benzyl 2-acetyl-2-ethylpentanoate (3g)

^1H -NMR(400MHz, CDCl_3) δ 0.74(t, $J=7.4\text{Hz}$, 3H), 0.89(t, $J=7.4\text{Hz}$, 3H), 1.07(sext, $J=7.4\text{Hz}$, 2H), 1.74-1.99(m, 4H), 2.03(s, 3H), 5.14(d, $J=12.6\text{Hz}$, 1H), 5.17(d, $J=12.6\text{Hz}$, 1H) and 7.29-7.36(m, 5H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 8.21, 14.46, 17.13, 24.13, 26.67, 32.87, 63.92, 66.75, 128.20, 128.25, 128.41, 135.26, 172.22 and 204.92 ppm. IR(neat): 3070, 3030, 2960, 2890, 1740, 1710 and 1500 cm^{-1} . EI-MS: m/z 262(M^+). Anal.Calcd.for $\text{C}_{16}\text{H}_{22}\text{O}_3$: C, 73.25; H, 8.45. Found: C, 73.23; H, 8.67.

Benzyl 2-acetyl-2-propylpentanoate (3h)

^1H -NMR(270MHz, CDCl_3) δ 0.88(t, $J=7.1\text{Hz}$, 6H), 1.01-1.14(m, 4H), 1.73-1.93(m, 4H), 2.03(s, 3H), 5.15(s, 2H) and 7.30-7.38(m, 5H) ppm. ^{13}C -NMR(67.5MHz, CDCl_3) δ 14.49, 17.24, 26.68, 33.57, 63.65, 66.75, 128.18, 128.22, 128.38, 135.26, 172.22 and 204.85 ppm. IR(neat): 3070, 3030, 2960, 2870, 1740, 1710 and 1500 cm^{-1} . EI-MS: m/z 276(M^+). Anal.Calcd.for $\text{C}_{17}\text{H}_{24}\text{O}_3$: C, 73.88; H, 8.75. Found: C, 73.59; H, 8.77.

Benzyl 2-acetyl-2-pentanoate (3i)^[b]

^1H -NMR(400MHz, CDCl_3) δ 0.91(t, $J=7.4\text{Hz}$, 3H), 1.27-1.33(m, 2H), 1.80-1.88(m, 2H), 2.17(s, 3H), 3.47(dd, $J=8.0\text{Hz}$, $J=7.2\text{Hz}$, 1H), 5.14(d, $J=12.4\text{Hz}$, 1H), 5.18(d, $J=12.4\text{Hz}$, 1H) and 7.31-7.37(m, 5H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 13.77, 20.62, 28.73, 30.21, 59.56, 66.92, 128.13, 128.27, 128.45, 135.23, 169.54 and 202.81 ppm. IR(neat): 3070, 3030, 2960, 2870, 1740, 1710 and 1500 cm^{-1} . EI-MS: m/z 234(M^+).

2-Acetyl-2-methylpentanenitrile (3j)^[c]

^1H -NMR(400MHz, CDCl_3) δ 0.97(t, $J=7.2\text{Hz}$, 3H), 1.35-1.67(m, 6H), 1.85-1.92(m, 1H) and 2.41(s, 3H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 13.81, 18.57, 22.11, 26.92, 39.00, 49.13, 121.11 and 202.07 ppm. IR(neat): 2970, 2880, 2240, 1730 and 1460 cm^{-1} . EI-MS: m/z 139(M^+). Anal.Calcd.for $\text{C}_{14}\text{H}_{17}\text{N}_5\text{O}_4$ (2,4-dinitrophenylhydrazine): C, 52.66; H, 5.73; N, 21.93. Found: C, 52.54; H, 5.31; N, 21.64.

2-Acetyl-2-methylhexanenitrile (3k)^[c]

^1H -NMR(400MHz, CDCl_3) δ 0.93(t, $J=7.0\text{Hz}$, 3H), 1.31-1.69 (m, 8H), 1.86-1.94(m, 1H) and 2.42(s, 3H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 13.66, 22.14, 22.48, 26.99, 27.25, 36.78, 49.15, 121.18 and 202.14 ppm. IR(neat): 2960, 2930, 2870, 2240, 1730 and 1460 cm^{-1} . EI-MS: m/z 153(M^+).

2-Acetyl-2,4-dimethylpentanenitrile (3l)^[c]

^1H -NMR(400MHz, CDCl_3) δ 0.92(d, $J=6.6\text{Hz}$, 3H), 1.02(d, $J=6.6\text{Hz}$, 3H), 1.49(s, 3H), 1.59(dd, $J=5.0\text{Hz}$, $J=13.4\text{Hz}$, 1H), 1.79-1.93(m, 2H) and 2.44(s, 3H) ppm. ^{13}C -NMR(100MHz, CDCl_3) δ 22.84, 23.43, 23.67, 25.95, 26.91, 44.91, 48.28, 121.40 and 202.23 ppm. IR(neat): 2960, 2880, 2240, 1730 and 1460 cm^{-1} . EI-MS: m/z 153(M^+).

Typical procedure for Zn-promoted three components joining reaction of α,β -unsaturated esters, alkyl iodides and acid anhydrides as the acylating agents is as follows: Into a DMF (20 ml) solution containing

commercially available Zn plate (60 mmol) was added dropwise a DMF solution (40 ml) of α,β -unsaturated esters (**1**, 10 mmol), alkyl iodides (**2**, 50 mmol) and acid anhydrides (**6**, 50mmol) at 20-25°C with magnetic stirring under nitrogen atmosphere during a period of 45min. in a water bath. After the addition, the mixture was stirred at 20-25°C for additional about 15 h. After the reaction, the reaction mixture was poured into 300 ml of saturated NaHCO₃ solution. Subsequent work up was carried out according to the procedure previously described.

Benzyl 2-methyl-3-oxo-2-propylpentanoate (3m)

¹H-NMR(400MHz, CDCl₃) δ 0.89(t, J =7.2Hz, 3H), 0.97(t, J =7.2Hz, 3H), 1.11(m, 2H), 1.34(s, 3H), 1.71-1.79(m, 1H), 1.84-1.91(m, 1H), 2.34(dq, J =17.4Hz, J =7.2Hz, 1H), 2.39(dq, J =17.4Hz, J =7.2Hz, 1H), 5.13(d, J =12.8Hz, 1H), 5.16(d, J =12.8Hz, 1H) and 7.28-7.36(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 8.09, 14.48, 17.60, 18.93, 31.58, 37.11, 59.49, 66.80, 128.12, 128.25, 128.44, 135.32, 172.92 and 208.12 ppm. IR(neat): 3070, 3030, 2960, 2890, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 262(M⁺). Anal.Calcd.for C₁₆H₂₂O₃: C, 73.25; H, 8.45. Found: C, 73.11; H, 8.56.

Benzyl 2-methyl-3-oxo-2-propylhexanoate (3n)

¹H-NMR(400MHz, CDCl₃) δ 0.79(t, J =7.2Hz, 3H), 0.89(t, J =7.2Hz, 3H), 1.13-1.21(m, 2H), 1.33(s, 3H), 1.51(sext, J =7.2Hz, 2H), 1.69-1.77(m, 1H), 1.84-1.91(m, 1H), 2.27(dt, J =17.8Hz, J =7.2Hz, 1H), 2.32(dt, J =17.8Hz, J =7.2Hz, 1H), 5.13(d, J =12.8Hz, 1H), 5.16(d, J =12.8Hz, 1H) and 7.27-2.32(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 13.57, 14.46, 17.15, 17.55, 18.76, 36.94, 40.19, 59.62, 66.77, 128.13, 128.23, 128.40, 135.30, 172.83 and 207.27 ppm. IR(neat): 3060, 3030, 2960, 2870, 1740, 1710 and 1500 cm⁻¹. EI-MS: m/z 276(M⁺). Anal.Calcd.for C₁₇H₂₄O₃: C, 73.88; H, 8.75. Found: C, 73.71; H, 8.54.

2-Benzyloxycarbonyl-2-methylcyclopentanone (3o)

¹H-NMR(400MHz, CDCl₃) δ 1.33(s, 3H), 1.83-2.07(m, 3H), 2.26-2.55(m, 3H), 5.11(d, J =12.4Hz, 1H), 5.16(d, J =12.4Hz, 1H) and 7.29-7.37(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 19.41, 19.56, 36.10, 37.61, 55.90, 66.83, 127.64, 128.06, 128.41, 135.50, 171.96 and 215.33 ppm. IR(neat): 3060, 3030, 2970, 2890, 1750, 1730 and 1500 cm⁻¹. EI-MS: m/z 232(M⁺). Anal.Calcd.for C₁₄H₁₆O₃: C, 73.39; H, 6.94. Found: C, 73.43; H, 6.65.

2-Benzyloxycarbonyl-2-methylcyclohexanone (3p)

¹H-NMR(400MHz, CDCl₃) δ 1.30(s, 3H), 1.43-1.50(m, 1H), 1.60-1.73(m, 3H), 1.95-1.99(m, 1H), 2.39-2.43(m, 2H), 2.49-2.54(m, 1H), 5.14(d, J =12.4Hz, 1H), 5.20(d, J =12.4Hz, 1H) and 7.29-7.37(m, 5H) ppm. ¹³C-NMR(100MHz, CDCl₃) δ 21.17, 22.51, 27.42, 38.16, 40.56, 57.10, 66.76, 127.90, 128.17, 128.42, 135.29, 172.64 and 207.74 ppm. IR(neat): 3070, 3030, 2940, 2870, 1730, 1710 and 1500 cm⁻¹. EI-MS: m/z 246(M⁺). Anal.Calcd.for C₁₅H₁₈O₃: C, 73.15; H, 7.37. Found: C, 72.96; H, 7.24.

Benzyl 2-ethyl-2-methylpentanoate (4)

¹H-NMR(270MHz, CDCl₃) δ 0.80(t, J =7.3Hz, 3H), 0.86(t, J =7.3Hz, 3H), 1.13(s, 3H), 1.14-1.74(m, 6H), 5.11(s, 2H) and 7.27-7.36(m, 5H) ppm. ¹³C-NMR(67.5MHz, CDCl₃) δ 8.92, 14.61, 17.89, 20.64, 31.96, 41.37, 46.39, 65.81, 127.74, 128.00, 128.26, 136.31 and 176.97 ppm. IR(neat): 3090, 3070, 3030, 2960, 2870, 1730 and 1500 cm⁻¹. EI-MS: m/z 234(M⁺). Anal.Calcd.for C₁₅H₂₂O₂: C, 76.88; H, 9.46. Found: C, 76.65; H, 9.41.

Benzyl 2-Methylpentanoate (5)^[d]

¹H-NMR(270MHz, CDCl₃) δ 0.88(t, J =7.2Hz, 3H), 1.16(d, J =6.8Hz, 3H), 1.24-1.73(m, 4H), 2.50(sext, J =6.8Hz, 1H), 5.10(s, 2H) and 7.28-7.34(m, 5H) ppm. ¹³C-NMR(67.5MHz, CDCl₃) δ 13.89, 16.96, 20.33, 35.82, 39.24, 65.79, 127.77, 127.84, 128.25, 136.08 and 176.26 ppm. IR(neat): 3090, 3070, 3030, 2960, 2940, 1740 and 1500 cm⁻¹. EI-MS: m/z 206(M⁺).

References and Notes

[a] Ueno, Y.; Setoi, H.; Okawara, M. *Tetrahedron Lett.* **1978**, 39, 3753.

2-Benzylloxycarbonyl-1-butene (1b)

¹H-NMR(270MHz, CDCl₃) δ 1.08(t, *J*=7.4Hz, 3H), 2.35(q, *J*=7.4Hz, 2H), 5.18(s, 2H), 5.53(s, 1H), 6.18(s, 1H) and 7.28-7.39(m, 5H) ppm. ¹³C-NMR(67.5MHz, CDCl₃) δ 12.63, 24.78, 66.19, 123.68, 127.82, 127.92, 128.34, 135.98, 141.97 and 166.83 ppm. IR(neat): 3090, 3060, 3030, 2970, 2940, 1720, 1630 and 1500 cm⁻¹. EI-MS: *m/z* 190(M⁺).

2-Benzylloxycarbonyl-1-pentene (1c)

¹H-NMR(270MHz, CDCl₃) δ 0.92(t, *J*=7.4Hz, 3H), 1.50(sext, *J*=7.4Hz, 2H), 2.30(t, *J*=7.4Hz, 2H), 5.18(s, 2H), 5.53(s, 1H), 6.19(s, 1H) and 7.30-7.36(m, 5H) ppm. ¹³C-NMR(67.5MHz, CDCl₃) δ 13.66, 21.54, 33.88, 66.24, 124.77, 127.83, 127.95, 128.38, 136.04, 140.42 and 166.93 ppm. IR(neat): 3070, 3030, 2960, 2930, 1720, 1630 and 1500 cm⁻¹. EI-MS: *m/z* 204(M⁺).

[c] Nakamura, K.; Miyai, T.; Nagar, A.; Oka, S.; Ohno, A. *Bull. Chem. Soc. Jap.* **1989**, 62, 1179.

[d] Lutz, A. H.; Schnider, O. *Chimia* **1958**, 12, 291.

[e] Takimoto, S.; Inanaga, J.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jap.* **1981**, 54, 1470.

Benzyl alph-propylacrylate in the presence of EtI and Zn

Benzyl alph-propylacrylate in the presence of EtI and Zn

C:\WINNMR98\COMMON\DEFAULT.ALS
2-Acetyl-2-methylhexanenitrile

Z-Acetyl-2-methylhexanenitrile

2-Acetyl-2,4-dimethylpentanenitrile

2-Acetyl-2,4-dimethylpentanenitrile

1H NMR

13C NMR