π -Allylic C1-Substitution in Water with Nitromethane Using Amphiphilic Resin-Supported Palladium Complexes

Yasuhiro Uozumi and Toshimasa Suzuka.*

Institute for Molecular Science (IMS) and CREST, Higashiyama 5-1, Myodaiji,
Okazaki 444-8787, Japan

uo@ims.ac.jp

SUPPORTING INFORMATION

Page S2: General methods and materials

Page S3: Palladium-Catalyzed Allylic Substitution

Page S3-S5: Spectroscopic data for compounds (2)(2')(5)(7)(9)(11)(13)(15)(17)

Page S6: Asymmetric Palladium-Catalyzed Allylic Substitution

Page S6-S7: Spectroscopic for compound (11)(13)(15)(20)

Page S8: Conversion to carboxylic acid (21)

Page S8-S9: Spectra for compound (21) and Reference

Page S10-S24: NMR charts (^1H and ^{13}C)

General. All manipulations were carried out under a nitrogen atmosphere. Nitrogen gas was dried by passage through P_2O_5 . NMR spectra were recorded on a JEOL JNM LA-500 spectrometer (500 MHz for 1H , 125 MHz for ^{13}C , and 376 MHz for ^{19}F NMR). Chemical shifts are reported in δ ppm referenced to an internal tetramethylsilane standard for 1H NMR, chloroform-*d* (δ 77.0) for ^{13}C , and trifluoroacetic acid (δ -79.0) for ^{19}F . GC-MS analysis was performed on an Agilent 6890 (GC system) and an Agilent 5973 (Mass detector). IR analysis was performed on a JASCO FTIR-460. HPLC analysis was performed on a JASCO PU-1580 liquid chromatograph system. Optical rotations were measured on a JASCO P-1020 polarimeter. The agitation of the reaction mixture containing resin-supported catalysts was performed on a wrist-action shaker (Burrel Scientific, Inc.) and Peti-Szyer (Shimadzu Scientific Research Inc).

Materials. The PS-PEG resin-supported Palladium Complex **3** was prepared according to the reported procedures.^{S1} The supported P, N-Chelating Chiral Palladium Complex **18** was prepared according to the reported procedures.^{S2}

Palladium-Catalyzed Allylic Substitution of Cinnamyl Carbonate (1) with Nitromethane (Schemes 1, 2 and 3). A typical procedure is given for the reaction with nitromethane and methyl cinnamyl carbonate (2) (Scheme 1). To a solution of 3 (64.0 mg, 0.02 mmol), lithium carbonate (116 mg, 1.60 mmol) and methyl cinnamyl carbonate (2) (76.8 mg, 0.40 mmol) in H₂O (0.40 mL) was added nitromethane (73.2 mg, 1.2 mmol) and the mixture was stirred at 60 °C for 12 h. The reaction mixture was filtered and the resin beads were rinsed three times with THF and MeOH. To the combined filtrate and washings were added aqueous NH₄Cl and the mixture was extracted with methyl tert-butyl ether (MTBE). The combined extract was washed with aqueous sodium chloride and dried over anhydrous magnesium sulfate. After removal of the solvent, the residual oil was chromatographed on silica gel (hexane/ethyl acetate/Et₃N = 9/1/0.5) to give 65.1 mg (92% yield) of a mixture of 4-nitro-1-phenylbutene (2) and 4-nitro-3-phenylbutene (2') in a ratio of 89:11: **4-Nitro-1-phenylbutene (2):** ¹H NMR (CDCl₃) δ 7.34-7.22 (m, 5 H), 6.52 (d, *J* = 15.5 Hz, 1 H), 6.11 (dt, *J* = 15.5, 7.0 Hz, 1H), 4.50 (t, *J* = 7.0 Hz, 2 H), 2.89 (dt, *J* = 7.0, 7.0 Hz, 2 H): ¹³C NMR (CDCl₃) δ 136.5, 134.0, 128.6, 127.8, 126.3, 122.9, 75.0, 30.7. MS (EI): *m/z* (rel%) 177 (3, M+), 130 (bp), 91 (33). CAS registry number: 76024-91-4. **4-Nitro-3-phenylbutene (2'):** ¹H NMR (CDCl₃) δ 7.40-7.15 (m, 5H), 5.98 (ddd, *J* = 17.5, 10.5, 8.0 Hz, 1H), 5.22 (d, *J* = 10.5 Hz, 1 H), 5.17 (d, *J* = 17.5 Hz, 1 H), 4.65 (m, 2 H), 4.20 (td, *J* = 8.0, 8.0 Hz, 1 H): ¹³C NMR (CDCl₃) δ 135.7, 133.4, 129.0, 127.8, 124.9, 117.5, 79.4, 74.5. MS (EI): *m/z* (rel%) 177 (0.8, M+), 130 (bp), 91 (52). CAS registry number: 207393-02-0. **4-Nitro-1-(*p*-tolyl)butene (5):** ¹H NMR (CDCl₃) δ

7.20 (d, $J = 8.2$ 1 H), 7.11 (d, $J = 8.2$ Hz, 1 H), 6.48 (d, $J = 15.6$ Hz, 1H), 6.04 (dt, $J = 15.6, 7.3$ Hz, 2 H), 4.46 (t, $J = 6.8$ Hz, 2 H), 2.88 (m, 2H), 2.32 (s, 3H): ^{13}C NMR (CDCl_3) δ 136.5, 134.0, 128.6, 127.8, 126.3, 122.9, 75.0, 30.7. MS (EI): m/z (rel%) 191 (9, M^+), 129 (bp), 105 (45). EI-HRMS m/z 191.094 (M^+ calcd for 191.094 $\text{C}_{11}\text{H}_{13}\text{NO}_2$). IR (ATR): (cm^{-1}) ν 3012, 2919, 1546, 1512. **4-Nitro-3-(*p*-tolyl)butene (5')**: MS (EI): m/z (rel%) 191 (0.8, M^+), 129 (bp), 91 (19). EI-HRMS m/e 191.095 (M^+ calcd for 191.094 $\text{C}_{11}\text{H}_{13}\text{NO}_2$). **4-Nitro-1-(*p*-anisyl)butene (7)**: ^1H NMR (CDCl_3) δ 7.26 (d, $J = 9.1$ Hz, 1 H), 6.84 (d, $J = 9.1$ Hz, 1 H), 6.46 (d, $J = 15.8$ Hz, 1H), 5.96 (dt, $J = 15.8, 7.3$ Hz, 1 H), 4.48 (t, $J = 6.7$ Hz, 2 H), 3.80 (s, 3H), 2.88 (td, $J = 6.7, 7.3$ Hz, 2H): ^{13}C NMR (CDCl_3) δ 159.3, 133.4, 129.2, 127.4, 120.5, 114.0, 75.1, 55.2, 30.8. MS (EI): m/z (rel%) 207 (53, M^+), 160 (bp), 121 (73), 91 (54). EI-HRMS m/z 207.088 (M^+ calcd for 207.089 $\text{C}_{11}\text{H}_{13}\text{NO}_3$). IR (ATR): (cm^{-1}) ν 2917, 1550, 1510. **4-Nitro-3-(*p*-anisyl)butene (7')**: ^1H NMR (CDCl_3) δ 7.13 (d, $J = 8.2$ Hz, 1H), 6.88 (d, $J = 8.2$ Hz, 1H), 5.98-5.92 (m, 1H), 5.17 (dd, $J = 17.0, 10.5$ Hz, 2 H), 4.65-4.56 (m, 2H), 4.2 (m, 1 H): ^{13}C NMR (CDCl_3) δ 159.0, 136.0, 129.2, 128.5, 117.3, 113.9, 79.5, 55.2, 46.9. MS (EI): m/z (rel%) 207 (7.8, M^+), 160 (bp), 91 (49). EI-HRMS m/z 207.089 (M^+ calcd for 207.089 $\text{C}_{11}\text{H}_{13}\text{NO}_3$). **1,3-Diphenyl-2-propenylnitromethane (9)**: ^1H NMR (CDCl_3) δ 7.39-7.21 (m, 10 H), 6.51 (d, $J = 15.8$ Hz, 1 H), 6.30 (dd, $J = 15.8, 7.3$ Hz, 1 H), 4.78-4.70 (m, 2 H), 4.37 (dd, $J = 15.8, 7.3$ Hz, 1 H): ^{13}C NMR (CDCl_3) δ 138.3, 136.2, 132.9, 129.1, 128.5, 127.9, 127.8, 127.5, 126.9, 126.4, 79.6, 47.2. MS (EI): m/z (rel%) 253 (1, M^+), 206 (52), 115 (46), 91 (bp). CAS registry number: 174707-03-0. **Cyclopentenyl nitromethane (11)**: ^1H NMR (CDCl_3) δ 5.92 (m, 1H),

5.62 (m, 1H), 4.41 (dd, $J = 11.5, 6.0$ Hz, 1 H), 4.27 (dd, $J = 11.5, 8.0$ Hz, 1 H), 2.41 (m, 2H), 2.15 (dddd, $J = 5.5, 5.5, 8.5, 22$ Hz, 1 H), 1.61 (m, 2H): ^{13}C NMR (CDCl_3) δ 134.5, 129.2, 79.5, 44.4, 31.6, 26.9. MS (EI): m/z (rel%) 127 (0.1, M^+), 97 (27), 83 (86), 67 (bp). EI-MS m/z 127.062 (M^+ calcd for 127.063 $\text{C}_6\text{H}_9\text{NO}_2$). CAS registry number: 305805-53-2. **Cyclohexenylnitromethane (13):** ^1H NMR (CDCl_3) δ 5.87 (m, 1H), 5.49 (dd, $J = 9.5, 1.5$ Hz, 1 H), 4.32 (dd, $J = 11.5, 6.5$ Hz, 1 H), 4.27 (dd, $J = 11.5, 8.0$ Hz, 1 H), 2.97 (m, 1 H), 2.03 (m, 2H), 1.83 (m, 1H), 1.68 (m, 1 H), 1.65 (m, 1 H), 1.39 (m, 1 H): ^{13}C NMR (CDCl_3) δ 131.1, 125.0, 80.0, 34.4, 25.9, 24.8, 20.0. MS (EI): m/z (rel%) 94 (87), 79 (bp), 67 (26). FAB-HRMS m/z 141.079 (M^+ calcd for 141.078 $\text{C}_7\text{H}_{11}\text{NO}_2$). CAS registry number: 85097-24-1.

Cycloheptenylnitromethane (15): ^1H NMR (CDCl_3) δ 5.91 (m, 1H), 5.49 (dd, $J = 11.0, 2.5$ Hz, 1H), 4.38 (dd, $J = 12.0, 7.5$ Hz, 1 H), 4.34 (dd, $J = 12.0, 7.5$ Hz, 1 H), 3.12 (m, 1 H), 2.20 (m, 2 H), 1.97 (m, 1 H), 1.67 (m, 3H), 1.40 (m, 2H): ^{13}C NMR (CDCl_3) δ 134.4, 130.5, 80.3, 38.7, 30.3, 29.3, 28.5, 26.3. MS (EI): m/z (rel%) 108 (74), 93 (bp), 79 (83), 67 (69). FAB-HRMS m/z 155.094 (M^+ calcd for 155.094 $\text{C}_8\text{H}_{13}\text{NO}_2$). CAS registry number: 305805-55-4. **cis-5-Methoxycarbonyl-3-nitromethylcyclohexene (17):** ^1H NMR (CDCl_3) δ 5.88 (m, 1 H), 5.50 (d, $J = 10.0$ Hz, 1 H), 4.35 (dd, $J = 18.5, 6.5$ Hz, 1 H), 4.28 (dd, $J = 12.0, 8.0$ Hz, 1 H), 3.70 (br s, 3 H), 3.09 (br, 1 H), 2.67 (m, 1 H), 2.32 (m, 3 H), 1.51 (ddd, $J = 12.2, 12.2, 12.2$ Hz, 1 H): ^{13}C NMR (CDCl_3) δ 175.0, 128.9, 125.0, 79.9, 51.8, 38.7, 35.0, 29.1, 27.5. MS (EI): m/z (rel%) 168 ($\text{M}^+ - 31$ (^-OMe) (7)), 93 (bp). CAS registry number: 305805-56-5.

Palladium-Catalyzed Asymmetric Allylic Substitution of Cyclic Allylic Carbonates (10, 12, or 14) with Nitromethane (Schemes 4 and 5). A typical procedure is given for the reaction of methyl cycloheptenyl carbonate (**14**) with nitromethane. To a mixture of **18** (68.0 mg, 0.02 mmol Pd), lithium carbonate (116 mg, 1.60 mmol) and methyl cycloheptenyl carbonate (**14**) (76.8 mg, 0.40 mmol) in H₂O (0.40 mL) was added nitromethane (73.2 mg, 1.2 mmol) and the mixture was stirred at 60 °C for 12 h. The reaction mixture was filtered and the resin beads were rinsed three times with THF and MeOH. To the combined filtrate and washings was added aqueous NH₄Cl and the mixture was extracted with MTBE. The combined extracts were washed with aqueous sodium chloride and dried over anhydrous magnesium sulfate. The solvent was evaporated and the residue was chromatographed on silica gel ((hexane/ethyl acetate/Et₃N = 9/1/0.5) to give 56.4 mg (91% yield) of a cycloheptenylnitromethane (**15**): ¹H NMR (CDCl₃) δ 5.91 (m, 1H), 5.49 (dd, *J* = 11.0, 2.5 Hz, 1H), 4.38 (dd, *J* = 12.0, 7.5 Hz, 1 H), 4.34 (dd, *J* = 12.0, 7.5 Hz, 1 H), 3.12 (m, 1 H), 2.20 (m, 2 H), 1.97 (m, 1 H), 1.67 (m, 3H), 1.40 (m, 2H): ¹³C NMR (CDCl₃) δ 134.4, 130.5, 80.3, 38.7, 30.3, 29.3, 28.5, 26.3. MS (EI): *m/z* (rel%) 108 (74), 93 (bp), 79 (83), 67 (69). FAB-MS *m/e* 155.094 (*M*⁺ calcd for 155.094 C₈H₁₃NO₂). [α]_D²⁰ +3.4 (*c* 0.5, CHCl₃) for (*S*)-**15** of 98% ee. (lit.^{S3} [α]_D²⁰ +3.2 (*c* 0.5, CHCl₃) for (*S*)-**15** of 96% ee). The enantiomeric excess was determined by HPLC analysis using a chiral stationary phase column (Chiralcel AD-H (500mm), eluent; n-hexane, flow rate; 0.5 mL/min, retention times; major isomer 70 min and minor 67) to be 98% ee.

Cyclopentenynitromethane (11): [α]_D²⁴ -105.0 (*c* 1.0, CHCl₃) for (*S*)-**11** of 80% ee.

The enantiomeric excess was determined by HPLC analysis using a chiral stationary phase column (Chiralcel AD-H (500mm), eluent; *n*-hexane:*i*-PrOH = 500:1, flow rate; 0.5 mL/min, retention times; major isomer 55 min and minor 54) to be 80% ee. CAS registry number: 305805-53-2. **Cyclohexenyl nitromethane (13):** $[\alpha]_{\text{D}}^{25} -51.0$ (*c* 1.0, CHCl₃) for (*S*)-**13** of 70% ee. The enantiomeric excess was determined by HPLC analysis using a chiral stationary phase column (Chiralcel AD-H (500mm), eluent; *n*-hexane:*i*-PrOH = 200:1, flow rate; 0.5 mL/min, retention times; major isomer 23 min and minor 22) to be 70% ee. CAS registry number: 85097-24-1. ***tert*-Butyl (5*R*)-5-nitromethyl-1,2,5,6-tetrahydropyridinecarboxylate (20):** ¹H NMR (CDCl₃) δ 5.87 (br, d, *J* = 9.7 Hz, 1H), 5.71 (br d, *J* = 7.9 Hz, 1H), 4.36 (dd, *J* = 12.8, 8.5 Hz, 1H), 4.26 (dd, *J* = 12.8, 5.6 Hz, 1H), 4.10 (br m, 1 H), 3.75-3.72 (m, 2 H), 3.36 (br d, *J* = 11.5 Hz, 1 H), 3.04 (br m, 1 H), 1.46 (s, 9 H); ¹³C NMR (CDCl₃) δ 154.8, 128.6, 123.7, 80.4, 43.5, 42.7, 34.3, 28.3, 28.2. MS (EI): *m/z* (rel%) 242 (0.1, M⁺), 169 (18), 94 (40), 57 (bp). IR (ATR): (cm⁻¹) ν 2931, 2977, 1691, 1551, 1420. FAB-HRMS *m/z* 242.126 (M⁺ calcd for 242.126 C₁₁H₁₈N₂O₄). $[\alpha]_{\text{D}}^{20} -37.1$ (*c* 1.00, CHCl₃). The enantiomeric excess was determined by the Mosher's method for *tert*-butyl 5-aminomethyl-1,2,3,4,5,6-hexahydropyridinecarboxylate derived from 5-nitromethyl-1,2,5,6-tetrahydropyridinecarboxylate (¹⁹F NMR (CDCl₃) major isomer δ 72.1 and minor δ 72.3) to be 97% ee.

Reaction of the *tert*-Butyl (5*R*)-5-nitromethyl-1,2,5,6-tetrahydropyridinecarboxylate (20) (97% ee) with Iodobutane Giving (3*S*)-*N*-*tert*-butoxycarbonyl-1,2,3,6-tetrahydropyridin-3-carboxylic acid (21). To a solution of *tert*-Butyl (5*R*)-5-nitromethyl-1,2,5,6-tetrahydropyridinecarboxylate (**20**) (97% ee) (76 mg, 0.3 mmol), KOH (67 mg, 1.2 mmol), and Tenta Gel (48 mg, 0.015 mmol) in H₂O (1.0 mL) was added iodobutane (110 mg, 0.8 mmol) and the mixture was stirred at 100 °C for 3 h. The reaction mixture was filtered and the resin beads were rinsed three times with H₂O. To the washings (H₂O) were added aqueous citric acid and the mixture was extracted with MTBE. The combined extracts were washed with aqueous sodium chloride and dried over anhydrous magnesium sulfate. The solvent was evaporated and the residue was chromatographed on silica gel (AcOEt) to give 34.1 mg (51% yield) of (3*S*)-*N*-*tert*-butoxycarbonyl-1,2,3,6-tetrahydropyridin-3-carboxylic acid (**21**): ¹H NMR (CDCl₃, 60 °C) δ 8.19 (br s, 1 H), 5.92 (d, *J* = 10.3 Hz, 1 H), 5.85 (d, *J* = 10.3 Hz, 1 H), 3.99-3.83 (m, 3 H), 3.62-3.58 (m, 1 H), 3.26 (m, 1 H), 1.46 (s, 9 H); ¹³C NMR (CDCl₃, 60 °C) δ 175.4, 154.9, 127.1, 122.8, 80.2, 43.2, 42.1, 40.7, 28.4. MS (EI): *m/z* (rel%) 170 (11, M⁺-57(-C₄H₉)), 126 (28), 62 (57), 57 (bp). [α]_D²⁰ -110 (c 1.0, MeOH) for (*R*)-**21** of 97% ee. (lit.^{S4} [α]_D²⁰ -112 (c 3.00, MeOH) for (*R*)-**21** of 99% ee). CAS registry number: 251948-99-9.

References

S1 (a) Uozumi, Y.; Danjo, H.; Hayashi, T. *Tetrahedron Lett.* **1997**, 38, 3557. (b)

Danjo, H.; Tanaka, D.; Hayashi, T.; Uozumi, Y. *Tetrahedron* **1999**, 55, 14341.

S2 Uozumi, Y.; Shibatomi, K. *J. Am. Chem. Soc.* **2001**, 123, 2919

S3 Trost, B. M. Surivet, J. P. *Angew. Chem. Int. Ed.* **2000**, 39, 3122.

S4 Altenbach, H. J.; Blanda, G. *Tetrahedron Asymmetry* **1998**, 9, 1519.

Supporting DATA (New Compound)

tert-Butyl 5-nitromethyl-1,2,5,6-tetrahydropyridinecarboxylate (**20**)

C:\Documents and Settings\ALPHA\Desktop\Uozumi\Gy\Uozumi\06042001 N boc H chart 1.als

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\06042001 N boc C.als

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\06042001 N boc H chart 1.als

4-Nitro-1-(*p*-tolyl)butene (5) + (5')

4-Nitro-1-(*p*-anisyl)butene (7)

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\060808 MeO-cinnamyl carbonate H pure Lals

```

MENUF non.th5atfTH5ATFG2
OBNUC 1H
OBFRQ 500.00 MHz
OBFIN 162410.00 Hz
PW1 6.40 usec
DEADT 56.80 usec
PREDL 0.20000 msec
INIWT 10.0000 msec
POINT 16384
SAMPO 16384
TIMES 32
DUMMY 1
FREQU 10000.00 Hz
FILTR 5000 Hz
DELAY 40.00 usec
ACQTM 1.6384 sec
PD 2.0000 sec
ADBIT 16
RGAIN 18
BF 0.23 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD SINGL
EXPCM Single pulse
IRNUC 1H
IRFRQ 500.00 MHz
IRFIN 162410.00 Hz
IRRPW 50 usec
IRATN 511
DFILE C:\Documents and Settings\ALI
SHMFL TH5ATFG2
LKFIN 70334.0 Hz
LKLEV 180
LGAIN 26
LKPHS 221
LKSG 1537
CSPED 11 Hz
FILDC
FILDF

```

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\060808 MeO-cinnamyl carbonate C pure Lals

```

MENUF bcm.h5xf2TH5ATFG2
OBNUC 13C
OBFRQ 125.65 MHz
OBFIN 127958.00 Hz
PW1 10.00 usec
DEADT 17.06 usec
PREDL 0.20000 msec
INIWT 10.0000 msec
POINT 32768
SAMPO 32768
TIMES 10000
DUMMY 1
FREQU 27173.91 Hz
FILTR 13600 Hz
DELAY 14.71 usec
ACQTM 1.2059 sec
PD 1.5000 sec
ADBIT 16
RGAIN 28
BF 0.23 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD SINGL
EXPCM Single pulse
IRNUC 1H
IRFRQ 500.00 MHz
IRFIN 162410.00 Hz
IRRPW 50 usec
IRATN 511
DFILE C:\Documents and Settings\ALI
SHMFL TH5ATFG2
LKFIN 70334.0 Hz
LKLEV 180
LGAIN 26
LKPHS 221
LKSG 1787
CSPED 10 Hz
FILDC
FILDF

```

4-Nitro-3-(*p*-anisyl)butene (7') + (7)

Known Compound

4-Nitro-1-phenylbutene (2)

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\06030701.als

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\06030702.als

C:\WINALPHA\COMMON\DEFAULT.ALS

1,3-Diphenyl-2-propenylnitromethane (9)

C:\Documents and Settings\ALPHA\Desktop\UozumiG\suzuka\06030602.als

C:\WINALPHA\COMMON\DEFAULT.ALS

Cyclopentenyl nitromethane (11)

Cyclohexenyl nitromethane (13)

C:\WINALPHA\COMMON\DEFAULT.ALS

```

MENUF non_th5atfTH5ATFG2
OBNUC 1H
OBFRO 500.00 MHz
OBFIN 162410.00 Hz
PW1 6.40 usec
DEADT 56.80 usec
PREDL 0.20000 msec
INIWT 10.0000 msec
POINT 16384
SAMPO 16384
TIMES 64
DUMMY 1
FREQU 10000.00 Hz
FILTR 5000 Hz
DELAY 40.00 usec
ACQTM 1.6384 sec
PD 2.0000 sec
ADBIT 16
RGAIN 19
BF 0.24 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD SINGL
EXPCM Single pulse
IRNUC 1H
IRFRQ 500.00 MHz
IRFIN 162410.00 Hz
IRRPW 50 usec
IRATN 511
DFILE C:\WINALPHA\COMMON\DEI
SHMFL TH5ATFG2
LKFIN 70334.0 Hz
LKLEV 180
LGAIN 23
LKPHS 200
LKSG 889
OSPED 11 Hz
FILDC
FILDF
  
```

C:\WINALPHA\COMMON\DEFAULT.ALS

```

MENUF bcm_th5
OBNUC 13C
OBFRO 125.65 MHz
OBFIN 127958.00 Hz
PW1 4.50 usec
DEADT 19.81 usec
PREDL 0.20000 msec
INIWT 10.0000 msec
POINT 32768
SAMPO 32768
TIMES 1000
DUMMY 1
FREQU 27173.91 Hz
FILTR 13600 Hz
DELAY 14.71 usec
ACQTM 1.2059 sec
PD 1.5000 sec
ADBIT 16
RGAIN 26
BF 0.24 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD SINGL
EXPCM Single pulse
IRNUC 1H
IRFRQ 500.00 MHz
IRFIN 162410.00 Hz
IRRPW 50 usec
IRATN 511
DFILE C:\WINALPHA\COMMON\DEI
SHMFL TH5ATFG2
LKFIN 70334.0 Hz
LKLEV 180
LGAIN 23
LKPHS 200
LKSG 1066
OSPED 11 Hz
FILDC
FILDF
  
```

Cycloheptenylnitromethane (15)

cis-5-Methoxycarbonyl-3-nitromethylcyclohexene (17)

C:\WINALPHA\COMMON\DEFAULT.ALS

C:\WINALPHA\COMMON\DEFAULT.ALS

(3S)-N-tert-butoxycarbonyl-1,2,3,6-tetrahydropyridin-3-carboxylic acid (21)

C:\WINALPHA\COMMON\DEFAULT.ALS

C:\WINALPHA\COMMON\DEFAULT.ALS