α -Amidation of Cyclic Ethers Catalyzed by Simple Copper Salt and A Mild and Efficient Preparation Method for α,ω -Aminoalcohols

Ling He^{a,b}, Jing Yu^b, Ji Zhang^a, Xiao-Qi Yu^{a*}

a. Department of Chemistry, Key Laboratory of Green Chemistry and Technology (Ministry of Education), Sichuan University, 29 Wangjiang Road, Chengdu, Sichuan 610064, P. R. China

b. Key Laboratory of Drug-Targeting of Education Ministry of China, West China School of Pharmacy, Sichuan University, No.17, Section3, South Renmin Road, Chengdu, Sichuan 610041, P. R. China;

Supporting Information

1. General Experimental Section:

All catalyses were performed under nitrogen and argon atmosphere. Silica gel F254 plates were used for TLC and spots were examined under UV light at 254 nm and developed by an iodine vapor. Flash chromatography was performed on silica gel H. Solvents were purified according to standard procedures. ¹H and ¹³C NMR spectra were recorded on Bruker AC-E 200 MHz, Varian Mercury 400 MHz and Bruker Avance 600 MHz spectrometer, using CDCl₃ as the solvent. The mass spectra (ESI / HRMS) were recorded on a Bruker Daltonics Data analysis 3.2 mass spectrometer.

2. General Procedure for the Amidation of Cyclic Ethers:

A reaction flask was charged with substrate, iodosobenzene diacetate, p-tolulensulfonamide, catalyst (10% mol) and dichloromethane, and then the mixture was stirred at 40 °C under nitrogen atmosphere for about 4 h. After cooling to room temperature, the resulting mixture was filtered; the filtrate was concentrated in vacuo. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate or dichloromethane/acetone as eluent to afford product of amidation.

3. General Procedure for Cleavage of products of amidation:

A mixture of tosylamide 2, sodium borohydride and tetrahydrofuran was stirred at room temperature under nitrogen atmosphere for about 2~4 h. To the resulting mixture, 20 ml of saturated NH₄Cl were added and extracted with ethyl acetate (20 ml $\times$ 3), the organic layer was dried with Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate or dichloromethane/acetone as eluent to give ω -amino-alcohols.

4. Characterization of compounds:

Tetrahydrofuran-2-yl-tolylsulfonamide

$^1\text{H-NMR}$ (CDCl_3 , 300MHz) δ (ppm)

1.73~1.84 (m, 3H), 1.86~1.89 (m, 1H), 2.44 (s, 3H), 3.68~3.73 (m, 2H), 5.29~5.39 (m, 1H), 7.28 (d, $J = 8.0$ Hz, 2H), 7.79 (d, $J = 8.0$ Hz, 2H).

$^{13}\text{C-NMR}$ (CDCl_3 , 75MHz) δ (ppm)

21.5, 23.9, 32.5, 67.1, 84.9, 126.9, 129.5, 138.5, 143.3.

MS (ESI-) m/z (%) for $\text{C}_{11}\text{H}_{13}\text{NO}_3\text{S}$ (M-H)

Calcd. 240.0. Found 240.0.

Cyclohexane-2-yl-tolylsulfonamide

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.33~1.38 (m, 2H), 1.45~1.54 (m, 1H), 1.78~1.84 (m, 2H), 2.40 (s, 3H), 3.38~3.40 (m, 1H), 3.68 (d, 1H), 4.76 (td, $J_1=2.0$, $J_2=9.2$, 1H), 5.40 (d, $J=9.2$ Hz, 1H), 7.26 (d, $J=8.0$ Hz, 2H), 7.78 (d, $J=8.0$ Hz, 2H).

$^{13}\text{C-NMR}$ (CDCl_3 , 75MHz) δ (ppm)

21.5, 22.3, 24.7, 31.8, 66.2, 82.0, 127.1, 129.4, 138.8, 143.2.

HRMS (ESI+) m/z (%) for $\text{C}_{12}\text{H}_{17}\text{NO}_3\text{S Na}$ (M+Na)

Calcd. 278.0821. Found 278.0826.

1,4-dioxane-2-yl-tolylsulfonamide

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

2.41 (s, 3H), 3.40~3.44 (m, 1H), 3.51~3.60 (m, 3H), 3.77~3.81 (dd, $J=2.4$ Hz and 12.6 Hz, 1H), 4.96 (m, 1H), 6.17 (d, $J=9.6$ Hz, 1H), 7.28 (d, $J=8$ Hz, 2H), 7.80 (d, $J=8$ Hz, 2H).

Cyclohexene-2-yl-tolylsulfonamide

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

2.43 (s, 3H), 2.57~2.64 (dt, 1H), 2.81~2.89 (m, 1H), 3.59~3.71 (m, 2H), 5.43 (d, $J=8.4\text{Hz}$, 1H), 6.10 (d, $J=8.8\text{Hz}$, 1H), 7.11 (d, $J=5.2\text{Hz}$, 1H), 7.19~7.22 (m, 2H), 7.31 (d, $J=8.4\text{Hz}$, 2H), 7.86 (d, $J=8.4\text{Hz}$, 2H)

HRMS (ESI+) m/z (%) for $\text{C}_{16}\text{H}_{17}\text{NO}_3\text{SNa}$ (M+Na)

Calcd. 326.0821. Found 326.0824.

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

2.42 (s, 3H), 3.07 (t, 2H), 4.57 (t, 2H), 7.22-7.35 (m, 2H), 7.31 (d, $J=8.4\text{Hz}$, 2H), 7.52 (td, $J=1.2$ and 0.8 Hz , 1H), 7.92 (d, $J=8.4\text{Hz}$, 2H), 8.08 (d, $J=8.0\text{Hz}$, 1H).

$^{13}\text{C-NMR-Dept}$ (CDCl_3 , 75MHz) δ (ppm)

21.6, 27.2 (-), 68.8 (-), 127.3, 127.6, 127.8, 129.1, 130.1, 134.2.

$^{13}\text{C-NMR}$ (CDCl_3 , 75MHz) δ (ppm)

21.5, 27.2, 68.6, 125.2, 127.3, 127.6, 127.8, 129.1, 130.1, 134.1, 137.9, 139.2, 142.9, 162.8.

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.54~1.67 (m, 4H), 2.43 (s, 3H), 2.97 (q, 2H), 3.63 (t, 2H), 4.86 (t, 1H), 7.31 (d, $J=8.4\text{ Hz}$, 2H), 7.74 (d, $J=8.4\text{ Hz}$, 2H),

$^{13}\text{C-NMR}$ (CDCl_3 , 100MHz) δ (ppm)

21.5, 26.2, 29.5, 42.9, 62.2, 127.1, 129.7, 143.3, 146.5.

MS (ESI+) m/z (%) for $\text{C}_{11}\text{H}_{18}\text{NO}_3\text{S}$ (M+H)

Calcd. 244.33. Found 244.4 (100%).

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.32~1.40 (m, 2H), 1.48~1.55 (m, 4H), 1.77 (br, 2H), 2.43 (s, 3H), 2.93 (q, 2H), 3.59 (t, 2H), 4.82 (br, 1H), 7.29 (d, $J=8.4\text{ Hz}$, 2H), 7.74 (d, $J=8.4\text{ Hz}$, 2H).

$^{13}\text{C-NMR}$ (CDCl_3 , 100MHz) δ (ppm)

21.5, 22.6, 29.1, 31.8, 42.9, 62.3, 127.0, 129.6, 136.9, 143.3.

HRMS (ESI+) m/z (%) for C₁₂H₁₉NO₃SNa (M+Na) Calcd. 280.0978. Found 280.0981.

¹H-NMR (CDCl₃, 600MHz) δ (ppm) 2.42 (s, 3H), 3.13 (t, 2H), 3.49 (m, 4H), 3.69 (t, 2H), 5.52 (br, 1H), 7.29 (d, *J* = 8.1 Hz, 2H), 7.76(d, *J* = 8.1 Hz, 2H).

¹³C-NMR (CDCl₃, 150MHz) δ (ppm) 21.5, 42.9, 61.6, 69.3, 72.2, 127.1, 129.7, 137.0, 143.4.

HRMS (ESI+) m/z (%) for C₁₁H₁₇NO₄SNa (M+Na) Calcd. 282.0784. Found 282.0793.

¹H-NMR (CDCl₃, 600MHz) δ (ppm) 1.26~1.33 (m, 4H), 1.43~1.54 (m, 4H), 2.43 (s, 3H), 2.92 (t, 2H), 3.59 (t, 2H), 7.29 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 8.4 Hz, 2H).

¹³C-NMR (CDCl₃, 150MHz) δ (ppm) 21.5, 25.1, 26.1, 29.4, 32.3, 43.0, 62.6, 127.1, 129.7, 136.9, 143.3.

HRMS (ESI+) m/z (%) for C₁₃H₂₁NO₃S (M+Na) Calcd. 294.1134. Found 294.1130.

2-Hydroxymethyl-1-phenylethyltolylsulfonamide

¹H-NMR (CDCl₃, 600MHz) δ (ppm) 2.44 (s, 3H), 2.71 (t, 2H), 3.88 (t, 2H), 4.07 (s, 2H), 7.12 (d, *J* = 4.0 Hz, 2H), 7.18 (d, *J* = 7.6 Hz, 1H), 7.25 (m, 1H), 7.31 (d, *J* = 8.0, 2H), 7.78 (d, *J* = 8.0, 2H).

¹³C-NMR (CDCl₃, 150MHz) δ (ppm) 21.5, 34.3, 45.3, 63.7, 126.6, 127.3, 128.9, 129.8, 129.9, 134.4, 142.0.

MS (ESI-) m/z (%) for C₁₆H₁₈NO₃S (M-H) Calcd. 304.1. Found 304.2 (100%).

N-Hydroxyethyl-tolylsulfonamide

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

2.43 (s, 3H), 3.09 (q, 2H), 3.69 (t, 2H), 5.20 (br, 1H), 7.31 (d, $J = 8.0$ Hz, 2H), 7.76 (d, $J = 8.0$ Hz, 2H).

$^{13}\text{C-NMR}$ (CDCl_3 , 100MHz) δ (ppm)

21.5, 45.2, 61.3, 127.1, 129.8, 143.6, 144.6.

Tosylamine-18-crown-6 (mixture)

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

2.40 (ns, 3nH), 3.25~3.85 (m, nH), 4.72 (d, $J=8.0$, nH), 6.97 (m, nH), 7.21~7.82 (nd, $J=8.0 \sim 8.4$ Hz, nH).

MS (ESI-) m/z (%) (M-H)

Calcd. $n=1$. 432.5; $n=2$, 601.7; $n=3$, 770.9; $n=4$, 939.2

Found. 432.6; 601.6; 770.7; 939.5

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.71~1.88 (m, 4H), 1.97~2.15 (m, 4H), 2.24 ~2.28 (m, 4H), 2.31 (s, 3H), 2.22 (s, 3H), 4.11 ~4.24 (m, 4H), 5.41~5.54 (m, 4H), 7.08 (d, $J=8.4$ Hz, 2H), 7.18 (m, $J=8.4$, 2H), 7.43~7.49 (m, 4H), 7.57~7.62 (m, 2H), 7.69 (d, $J=8.4$ Hz, 2H), 7.79 (d, $J=8.4$ Hz, 2H), 7.98(t, 4H).

$^{13}\text{C-NMR-DEPT}$ (CDCl_3 , 100MHz) δ (ppm)

21.3, 25.9(-), 26.3 (-), 32.5 (-), 32.8 (-), 66.2 (-), 66.4 (-), 75.8, 77.1, 85.5, 85.7, 126.4, 127.1, 127.2, 128.4, 128.5, 129.3, 129.5, 129.6, 133.1,

MS (ESI-) m/z (%) for $\text{C}_{19}\text{H}_{20}\text{NO}_5\text{S}$ (M-H)

Calcd. 374.1. Found 374.1 (100%).

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.62~2.25 (m), 2.12(s), 2.17 (s), 2.42 (s), 2.43 (s), 3.84~4.32(m), 5.03~5.12 (m, 2H), 5.32~5.43 (m, 2H), 5.59~5.63 (m, 2H), 7.25 (dd, $J=8.4\text{Hz}$, 8.0Hz, 4H), 7.80 (dd, $J=8.4\text{Hz}$, 8.0Hz, 4H).

MS (ESI-) m/z (%) for $\text{C}_{14}\text{H}_{18}\text{NO}_5\text{S}$ (M-H)

Calcd. 312.0. Found 311.9 (45%).

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ (ppm)

1.56~2.02 (m), 2.42 (s), 2.61~1.71~4.17 (m), 3.33~4.17 (m), 5.35~5.61 (m), 5.95 (d, $J=9.2\text{Hz}$), 6.12 (d, $J=8.8\text{Hz}$), 7.31 (m), 7.78 (m).

MS (ESI+) m/z (%) for $\text{C}_{12}\text{H}_{17}\text{ClNO}_3\text{S}$ (M+H)

Calcd. 290.3. Found 290.2 (100%).

MS (ESI-) m/z (%) for $\text{C}_{12}\text{H}_{15}\text{ClNO}_3\text{S}$ (M-H)

Calcd. 288.0. Found 288.0 (100%).

MS (ESI-) m/z (%) for $\text{C}_{13}\text{H}_{19}\text{N}_2\text{O}_3\text{S}$ (M-H)

Calcd. 283.1. Found 283.2 (88%).

HL-14 H1 CDC13 2004-11-8
Pulse Sequence: s2pu1

HL-17 H1 CDC13 2005-3-30

Pulse Sequence: s2pu1

243.09

HL-18-C-6 H1 CDC13 2006-12-20

Pulse Sequence: s2pu1

■ -Q1: 0.033 to 0.602 min from Sample 2 (Q1) of qual-061207-18-C-6-10.wiff (Turbo Spray), Centroided

Max. 2.8e6 cps

HL-21-2 H1 CDC13 2006-12-20

Pulse Sequence: s2pu1

HL-21(12.5) H1 CDCl3 2006-12-13
Pulse Sequence: s2pul

■ -Q1: 0.067 to 0.635 min from Sample 2 (Q1) of qual-061207-HL-21.wiff (Turbo Spray)

Max. 6.4e4 cps.

HL-35 H1 CDC13 2007-1-31

Pulse Sequence: szpu1

HL-54 H1 CDC13 2006-6-8
Pulse Sequence: s2pu1

MW = 313

HL-68-A H1 CDC13 2006-4-27
Pulse Sequence: s2pu1

Bruker Avance 600 probe: 13C-1H DUL TE: 300K sample: HL70 solvent: CDCL3 spectrum: 1H

Bruker Avance 600 probe: 13C-1H DUL TE: 300K sample: HL70 solvent: CDCL3 spectrum: 13C

Mass Spectrum Molecular Formular Report

Analysis Info

Analysis Name D:\Bruker\data\msdata\HL-70\1.d
Method 1pass_pos_low.tofpar
Sample Name HL-70
Comment ESI source

Acquisition Date 7/20/2006 3:37:00 PM

Operator operator name
Instrument BioTOF Q

Acquisition Parameter

Capillary End Plate -4500 V
EndP -4000 V

Capillary Exit 120 V
Collision energy 0 eV

detbias 2 V
Number of Averages 100

Sum Formula	Sigma	m/z	Err [ppm]	Mean Err [ppm]	rdB	N Rule	e ⁻
C 12 H 13 N 5 Na 1 S 1	0.03	282.0784	-3.08	-3.78	8.50	ok	even

20050302

Current Data Parameters
 NAME 0302
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date 20050302
 Time 13.49
 INSTRUM av300
 PROBHD 5 mm PABBO BB-
 PULPROG zg30
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 5995.204 Hz
 FIDRES 0.182959 Hz
 AQ 2.7329011 sec
 RG 101.6
 DW 83.400 usec
 DE 6.00 usec
 TE 292.8 K
 D1 2.50000000 sec
 TD0 1

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.70 usec
 PL1 -5.00 dB
 SFO1 300.1324010 MHz

F2 - Processing parameters
 SI 16384
 SF 300.1300025 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.40

13C

apt

129.6991
129.4068
127.1382
126.4539

82.0289

66.1795

31.7621
24.6642
22.3360
21.5590

Current Data Parameters
NAME 0302
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050302
Time 14.32
INSTRUM av300
PROBHD 5 mm PABBO BB-
PULPROG dept135
TD 65536
SOLVENT CDCl3
NS 325
DS 16
SWH 18832.393 Hz
FIDRES 0.287360 Hz
AQ 1.7400308 sec
RG 23170.5
DW 26.550 usec
DE 6.90 usec
TE 294.2 K
CNST2 140.0000000
d2 0.00357143 sec
d12 0.00002000 sec
D1 4.00000000 sec
DELTA 0.00000764 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
p2 12.00 usec
P1 6.00 usec
PL1 -6.00 dB
SFO1 75.4752960 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
p4 15.40 usec
P3 7.70 usec
PCPD2 80.00 usec
PL12 16.00 dB
PL2 -5.00 dB
SFO2 300.1314410 MHz

F2 - Processing parameters
SI 32768
SF 75.4677490 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.40

Pulse Sequence: s2pu1

Mass Spectrum Molecular Formular Report

Analysis Info

Analysis Name D:\Bruker\data\msdata\HL-72-1\1.d
 Method 1pass_pos_low.tofpar
 Sample Name HL-72-1
 Comment ESI source

Acquisition Date 7/20/2006 3:51:38 PM

Operator operator name
 Instrument BioTOF Q

Acquisition Parameter

Capillary End Plate -4500 V
 EndP -4000 V

Capillary Exit 120 V
 Collision energy 0 eV

detbias 2 V
 Number of 100
 Averages

Sum	Formula	Sigma	m/z	Err [ppm]	Mean Err [ppm]	rdb	N Rule	e ⁻
C 12 H 17 N 1 Na 1 O 3 S 1		0.04	278.0821	-1.52	-4.80	4.50	ok	even

HL-72-2 H1 CDC13 2006-5-26
Pulse Sequence: s2pu1

Mass Spectrum Molecular Formular Report

Analysis Info

Analysis Name D:\Bruker\data\msdata\HL-72-2\1.d
Method 1pass_pos_low.tofpar
Sample Name HL-72-2
Comment ESI source

Acquisition Date 7/20/2006 3:48:58 PM

Operator operator name
Instrument BioTOF Q

Acquisition Parameter

Capillary End Plate -4500 V
EndP -4000 V

Capillary Exit 120 V
Collision energy 0 eV

detbias 2 V
Number of Averages 100

Sum	Formula	Sigma	m/z	Err (ppm)	Mean Err (ppm)	rdB	N Rule	e ⁻
C 12 H 19 N 1 Na 1 O 3 S 1		0.02	280.0978	-1.30	-0.50	3.50	ok	even

$C_{12}H_{19}O_3NS$

HL-75 H1 CDC13 2006-6-6
Pulse Sequence: s2pu1

Mass Spectrum Molecular Formular Report

Analysis Info

Analysis Name D:\Bruker\data\msdata\HL-75\1.d
 Method 1pass_pos_low.tofpar
 Sample Name HL-75
 Comment ESI source

Acquisition Date 7/20/2006 3:34:35 PM

Operator operator name
 Instrument BioTOF Q

Acquisition Parameter

Capillary End Plate	-4500 V	Capillary Exit	120 V	detbias	2 V
EndP	-4000 V	Collision energy	0 eV	Number of Averages	100

Sum Formula	Sigma	m/z	Err [ppm]	Mean Err [ppm]	rdb	N Rule	e ⁻
C 13 H 21 N 1 Na 1 O 3 S 1	0.01	294.1134	1.36	-0.86	3.50	ok	even

HL-76-1-CDCL3-H1-2006-7-20

Pulse Sequence: s2pu1

MW = 284

HL-80-1-2-CDCL3-H1-2006-7-19

Pulse Sequence: s2pu1

HL-80-1 H1 CDC13 2006-6-20

Pulse Sequence: s2pu1

Mass Spectrum Molecular Formular Report

Analysis Info

Analysis Name D:\Bruker\data\msdata\HL-80-1R\1.d
 Method 1pass_pos_low.tofpar
 Sample Name HL-80-1R
 Comment ESI source

Acquisition Date 7/20/2006 3:59:28 PM

Operator operator name
 Instrument BioTOF Q

Acquisition Parameter

Capillary End Plate	-4500 V	Capillary Exit	120 V	detbias	2 V
EndP	-4000 V	Collision energy	0 eV	Number of Averages	100

Sum	Formula	Sigma	m/z	Err [ppm]	Mean Err [ppm]	rdB	N Rule	e ⁻
C 16 H 17 N 1 Na 1 O 3 S 1		0.02	326.0821	-0.86	-1.79	8.50	ok	even

HL-80-3-CDCL3-DEPT-2006-7-17

HL-84-3-CDCL3-C13-2006-10-18
Pulse Sequence: s2pu1

HL-84-3 H1 CDC13 2006-10-11
Pulse Sequence: s2pu1