SUPPORTING INFORMATION

FOR

[Rh(CO)₂Cl]₂-Catalyzed Domino Reactions Involving Allylic Substitution and Subsequent Carbocyclization Reactions

Brandon L. Ashfeld, Kenneth A. Miller, Anna J. Smith, Kristy Tran

and Stephen F. Martin^{*}

Department of Chemistry and Biochemistry, The University of Texas at Austin, Austin, TX

78712 USA

Experimental Procedures

Solvents and reagents were reagent-grade and used without purification unless otherwise noted. Tetrahydrofuran (THF) and diethyl ether (Et₂O) were passed through a column of neutral alumina and stored under argon. Acetonitrile was passed through two columns of molecular sieves and stored under argon. Toluene was passed through a column of Q5 reactant and stored under argon. All reactions were performed in flame-dried glassware under argon unless otherwise indicated. ¹H nuclear magnetic resonance (NMR) spectra were obtained at either 500, 400 or 300 MHz as solutions in CDCl₃. ¹³C NMR were obtained at either 125, 100 or 75 MHz as solutions in CDCl₃. Chemical shifts are reported in parts per million (ppm, δ), and referenced to TMS, and coupling constants are reported in Hertz (Hz). Spectral splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; comp, complex; and br, broad. Infrared (IR) spectra were obtained using a Perkin-Elmer FTIR 1600 spectrophotometer using sodium chloride plates and reported as wave numbers. Low resolution chemical ionization mass spectra were obtained with a Finnigan TSQ-70 instrument. High resolution measurements were made with a VG Analytical ZAB2-E instrument. Analytical thin layer chromatography was performed using Merck 250 micron 60F-254 silica gel plates. The plates were visualized with UV light, ninhydrin, phosphomolybdic acid, *p*-anisaldehyde, and potassium permanganate. Flash column chromatography was performed according to Still's procedure (Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923) using ICN Silitech 32-63 D 60A silica gel.

Procedure for the [Rh(CO)₂Cl]₂-Catalyzed Allylic Alkylation of **12.** Allylic carbonate **12** (1.0 mmol) was added to a solution of [Rh(CO)₂Cl]₂ (10 mol%) in THF (**12a** and **12b**) or DMF (**12c**) (5 mL), and the solution was stirred for 10 min at room temperature for **12a** and **12b**, and at -20 °C for **12c**. In a separate flask malonate **11** (1.5 mmol) was added to a slurry of NaH

(60% w/w in mineral oil, 1.4 mmol) in THF (5 mL) at room temperature, and the mixture was stirred for 20 min. The resulting sodium enolate was added *via* syringe to the solution of **12** and $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ at room temperature for **12a** and **12b** and $-20\text{ }^\circ\text{C}$ for **12c**. The mixture was then sealed in a screw cap vial under an atmosphere of argon, and stirring was continued until the starting carbonate was consumed (as indicated by TLC analysis). The reaction was then filtered through a short plug of silica gel eluting with Et_2O (50 mL), and the filtrate was concentrated under reduced pressure. The crude residue was purified by flash chromatography, eluting with pentane/ Et_2O (5:1 or 10:1) to furnish **13a-c**.

Procedure for the $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ -Catalyzed Tandem Allylic Alkylation/Pauson-Khand Annulation of 14. Malonate **15-17** (0.26 mmol) was added to a suspension of NaH (60% dispersion in mineral oil, 0.22 mmol) in dry, degassed THF (1 mL) and the reaction stirred for 15 min at room temperature. The solvent was removed under reduced pressure and the resulting yellow oil was azeotroped with toluene (2 x 3 mL). The residue was dissolved in THF (1 mL) and added *via* syringe to a solution of **14** (0.2 mmol) and $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ (10 mol%) in THF (1 mL). The reaction was stirred at room temperature for 1 h then heated to reflux and stirring continued for 12-24 h. The solvent was removed under reduced pressure and the residue was purified by flash chromatography eluting with hexanes/ EtOAc (3:1) to furnish the known products **18-20**.

Procedure for the $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ -Catalyzed Tandem Allylic Alkylation/Cycloisomerization of 21. Allylic trifluoroacetate **21** (0.67 mmol) was added to a solution of $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ (10 mol%) in degassed MeCN (0.75 mL), and the solution was stirred for 10 min at $-40\text{ }^\circ\text{C}$. In a separate flask, malonate **11** (0.27 mmol) was added to a slurry of NaH (60% dispersion in mineral oil, 0.41 mmol) in MeCN (2.0 mL) at room temperature, and the mixture

was stirred for 30 min. The resulting solution of sodium enolate was then added *via* syringe to the solution of **21** and $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ at $-40\text{ }^\circ\text{C}$. The reaction mixture was stirred for 6 h at $-40\text{ }^\circ\text{C}$, then sealed in a screw cap vial and heated to $110\text{ }^\circ\text{C}$ until the intermediate enyne **13c** was consumed (as indicated by GLC analysis). The reaction mixture was diluted with water (2 mL) and ether (2 mL), and the layers were separated. The aqueous phase was extracted with ether (5 x 2 mL), and the combined organic fractions were dried (MgSO_4) and concentrated under reduced pressure. The crude residue was purified by flash chromatography eluting with pentane/ Et_2O (5:1) to furnish **22**.

Procedure for the $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ -Catalyzed Tandem Allylic Alkylation/[5+2] Cycloaddition of Allylic Trifluoroacetates **23, **25** and **27**.** The allylic trifluoroacetate (0.1 mmol) was added to a solution of $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ (5 mol%) in degassed MeCN (0.5 mL), and the solution was stirred for 10 min at room temperature. In a separate flask, malonate **11** (1.5 mmol) was added to a slurry of NaH (60% w/w in mineral oil, 1.4 mmol) in degassed MeCN (0.5 mL) at room temperature, and the mixture was stirred for 20 min. The resulting sodium enolate was added *via* syringe to the solution of the trifluoroacetate and $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ at room temperature. The mixture was then sealed in a screw cap vial under an atmosphere of argon, and stirring was continued at room temperature until the starting material was consumed (as indicated by TLC analysis). The reaction was then heated to $80\text{ }^\circ\text{C}$ (e.g. bath temperature) and stirring continued until intermediate enyne was consumed (as indicated by TLC analysis). The reaction was then filtered through a short plug of silica gel eluting with Et_2O (50 mL), and the filtrate was concentrated under reduced pressure. The crude residue was purified by flash chromatography, eluting with pentane/ Et_2O (5:1 or 10:1) to furnish the products **24**, **26** or **28**.

b1a VI 108
expl s2put1

SAMPLE DEC. & VT
date Aug 15 2003 dfreq 499.349
solvent CDCl3 dn H1
file exp dpr 30
ACQUISITION dof 0
sfrq 499.350 dm nm
tn H1 dmm C
at 4.865 dmf 200
np 89888 dres 1.0
sw 9237.9 dres 1.0
fb 5000 homo n
bs 32 temp 27.0
tpwr 54 PROCESSING
pw 2.0 lb 0.10
d1 2.000 wfile
tof 1123.4 proc ft
nt 32 fn not used
ct 32 math f
atlock n
gain 40 werr
il n wexp
in n wbs
dp n wnt
hs y wft
hs nm
DISPLAY
sp -249.8
wp 5243.0
vs 16
sc 0
wc 250
h2mm 20.97
is 706.95
rfl 1000.3
rfp 0
th 2
ins 1.000
ai cdc ph

13b

b1a-VII-179

Pulse Sequence: s2pul
Solvent: CDCl₃
Ambient temperature
Mercury-400BB "nmr6"

Relax. delay 2.000 sec
Pulse 47.8 degrees
Acq. time 2.856 sec
Width 5602.2 Hz
8 repetitions
OBSERVE H1, 400.2669761 MHz
DATA PROCESSING
Line broadening 0.1 Hz
Ft size 32768
Total time 0 min, 41 sec

STANDARD 1H OBSERVE

Pulse Sequence: szpul
Solvent: CDCl3
Ambient temperature
UNITYplus-300 "nmr2"

Relax. delay 1.000 sec
Pulse 15.0 degrees
Acq. time 3.813 sec
Width 4196.4 Hz
40 repetitions
OBSERVE H1, 300.1390308 MHz
DATA PROCESSING
Line broadening 0.1 Hz
FT size 32768
Total time 8 min, 2 sec

21

STANDARD 1H OBSERVE

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 File: AJS-1-cycloadduct
 UNITYplus-300 "nmr2"
 Relax. delay 1.000 sec
 Pulse 15.0 degrees
 Acq. time 3.813 sec
 Width 4196.4 Hz
 72 repetitions
 OBSERVE H1, 300.1390303 MHz
 DATA PROCESSING
 FT size 32768
 Total time 20 min, 5 sec

b1a-VII-122

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
Mercury-400BB "nmr6"

Relax. delay 2.000 sec
Pulse 15.9 degrees
Acq. time 2.856 sec
Width 5602.2 Hz
8 repetitions
OBSERVE H1, 400.2669771 MHz
DATA PROCESSING
Line broadening 0.1 Hz
FT size 32768
Total time 0 min, 41 sec

23

b1a-VII-176

Pulse Sequence: s2pu1
Solvent: CDCl3
Ambient temperature
Mercury-400BB "nmr6"

Relax. delay 2.000 sec
Pulse 47.8 degrees
Acq. time 2.856 sec
Width 5602.2 Hz
8 repetitions
OBSERVE H1, 400.2669768 MHz
DATA PROCESSING
Line broadening 0.1 Hz
FT size 32768
Total time 0 min, 0 sec

25

b1a-VII-138

Pulse Sequence: s2pu1

Solvent: CDCl3

Ambient temperature

Mercury-400BB "nmr6"

Relax. delay 2.000 sec

Pulse 47.8 degrees

Acq. time 2.856 sec

Width 5602.2 Hz

8 repetitions

OBSERVE H1, 400.2669771 MHz

DATA PROCESSING

Line broadening 0.1 Hz

FT size 32768

Total time 0 min, 0 sec

27