N^G -hydroxyguanidines from Primary Amines

Nathaniel I. Martin,* Joshua J. Woodward, and Michael A. Marletta

*Department of Chemistry, University of California Berkeley,
204 Lewis Hall, Berkeley, California 94720*

nimartin@berkeley.edu

Supplemental Material (part 1 of 3)

Table of Contents

Part	Page	
1 of 3	S2-3	General Procedures
	S3-13	Experimental Procedures and Data for Compounds 7-12
2 of 3	S14-19	^1H and ^{13}C NMR Data for Selected Compounds in Table 1
3 of 3	S20-23	^1H and ^{13}C NMR Data for Compounds 12a, 12b
	S24	References

General Procedures

Reagents, solvents and solutions. All reagents employed were of American Chemical Society (ACS) grade or finer and were used without further purification unless otherwise stated. For anhydrous reactions, solvents were dried according to Perrin *et al.*¹ and Vogel.² Removal of solvent was performed under reduced pressure using a Büchi rotary evaporator, followed by evacuation (< 0.1 mm Hg) to constant sample weight. Deionized water was obtained from a Milli-Q reagent water system (Millipore Co., Milford, MA).

Purification Techniques. All reactions and fractions from column chromatography were monitored by thin layer chromatography (TLC) using plates with a UV fluorescent indicator (normal SiO₂, Merck 60 F₂₅₄; reversed-phase, Merck RP-8 and RP-18 F_{254S}). One or more of the following methods were used for visualization: UV absorption by fluorescence quenching; iodine staining; phosphomolybdic acid:ceric sulfate:sulfuric acid:H₂O (10 g:1.25 g:12 mL:238 mL) spray; and 50% sulfuric acid spray. Flash chromatography was performed according to the method of Still *et al.*³ using Merck type 60, 230-400 mesh silica gel.

Instrumentation for Compound Characterization. Melting points are uncorrected and were determined on a Büchi oil immersion apparatus using open capillary tubes. Low resolution FAB mass spectra (FAB-MS) were recorded on a VG 70-SE (VG Analytical) mass spectrometer and high resolution FAB mass spectra were obtained using a ZAB2-EQ instrument. Microanalyses were obtained using a Perkin Elmer 2400 Series II combustion analyzer. Nuclear magnetic resonance (NMR) spectra were obtained on Bruker AVQ-400 and AV-300 MHz spectrometers. ¹H NMR chemical shifts are reported in parts per million (ppm) downfield relative to tetramethylsilane (TMS) using the residual proton resonance of solvents as reference: CDCl₃ δ 7.24 and D₂O δ 4.72.

^{13}C NMR chemical shifts are reported relative to CDCl_3 δ 77.0. ^{19}F NMR chemical shifts are reported relative to CFCl_3 δ 0.0. ^1H NMR data are reported in the following order: multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; qn, quintet and m, multiplet), number of protons, coupling constant (J) in Hertz (Hz) and assignment. When appropriate, the multiplicity is preceded by br, indicating that the signal was broad. All literature compounds had ^1H NMR, and mass spectra consistent with the assigned structures.

Experimental Procedures and Data for Compounds 7-11

Compound 8

The O-THP protected hydroxylamine was prepared by a modified literature procedure.⁴ Briefly: To a vigorously stirred solution of N-hydroxyphthalimide (10.0 g, 61.4 mmol) in CH_2Cl_2 (70 mL) and dioxane (80 mL) was added dihydropyran (6.16 mL, 67.6 mmol) and p-toluenesulfonic acid (200 mg). After 2 hours the reaction was quenched by slow addition of saturated NaHCO_3 (100 mL). After separation, the organic layer was washed with brine and dried over Na_2SO_4 . Solvent removal under vacuum yielded a white solid which was next dissolved in dry benzene (30 mL), treated with methylhydrazine (2.82 mL, 53.0 mmol) and heated to 80°C for 15 minutes. The reaction mixture was then cooled on ice and filtered through celite (with benzene washes) to remove the precipitate. The benzene solution was then concentrated under vacuum to an approximate volume of 10 mL and the product purified by vacuum distillation (b.p. = 70°C at 10 mm Hg). 4.68 g (75%) of pure $\text{H}_2\text{N-OTHP}$ was thus obtained as a clear, colorless oil which solidified upon storage at 4°C (full agreement with published spectral characterization).

Compound 9

Using the approach of Jirgensons and coworkers,⁵ CbzCl (6.6 mL, 46 mmol) and dibenzo-18-crown-6 (0.84 g, 2.4 mmol) were added to a

suspension of KNCS (5.4 g, 56 mmol) in benzene (80 mL) and the mixture refluxed at 85°C for 3 hours. The solution was then cooled on ice and treated with H₂N-OTHP (2.93 g, 25 mmol) followed by warming to room temperature. After stirring for 1 hour the mixture was filtered through celite to remove salts and the filtrate evaporated to yield a white solid. The product was further treated with warm hexanes and after filtration and drying under vacuum 5.00 g (35% yield) of product was obtained. mp 103-105 °C; ¹H NMR (CDCl₃, 400 MHz) δ 11.83 (br s, 1H), 8.19 (br s, 1H), 7.42-7.32 (m, 5H), 5.24 (m, 1H), 5.18 (s, 2H), 3.95 (m, 1H), 3.73 (m, 1H), 2.00-1.55 (m, 6 H); ¹³C NMR (CDCl₃, 100 MHz) δ 176.2, 152.1, 134.3, 128.9, 128.8, 128.4, 102.2, 68.6, 62.6, 27.6, 24.9, 18.2; HRMS (EI, M+H) Calcd for C₁₄H₁₉N₂O₄S 311.1073, found 311.1066; Anal. Calcd for C₁₄H₁₈N₂O₄S: C, 54.18; H, 5.85; N, 9.03. Found: C, 54.40; H, 6.14; N, 8.86.

Compound 10

Benzyl amine (110 µL, 1.0 mmol) was added to a solution of thiourea **8** (310 mg, 1.0 mmol) in DMF (10.0 mL) followed by addition of NEt₃ (140 µL, 1.0 mmol) and HgCl₂ (472 mg, 1.0 mmol) and the mixture left to stir overnight.

The following morning the reaction mixture is partitioned between brine (20 mL) and ethyl acetate (30 mL). After separation, the aqueous layer is further extracted with ethyl acetate (30 mL) and the combined organic extracts washed with 1M NaHCO₃ (2 x 30 mL), filtered through celite and dried over Na₂SO₄. Following solvent removal, the product was isolated via flash chromatography (SiO₂, 1:3 ethyl acetate/hexane, R_f = 0.45) to yield 226 mg (75%) of a white solid. mp 108-110 °C; ¹H NMR (CDCl₃, 400 MHz) δ 9.95 (br s, 1H), 8.49 (br s, 1H), 7.40-7.32 (m, 10H), 5.17 (s, 2H), 4.87 (d, *J* = 5.3 Hz, 2H); ¹³C NMR (CDCl₃, 100 MHz) δ 179.2, 152.5, 136.2, 134.5, 128.8, 128.7, 128.3, 127.94, 127.90, 68.18, 49.60; HRMS (EI, M+H) Calcd for C₁₆H₁₇N₂O₂S 301.1007, found 301.1011; Anal. Calcd for C₁₆H₁₆N₂O₂S: C, 63.98; H, 5.37; N, 9.33. Found: C, 63.60; H, 5.19; N, 9.20.

Compound 11

Compound 10 (150 mg, 0.50 mmol) was dissolved in CH_2Cl_2 (10 mL) and treated with EDCI (100mg, 0.5 mmol), NEt_3 (70 μL , 0.5 mmol), and $\text{H}_2\text{N-OTHP}$ (70 μL , 0.55 mmol). TLC analysis (3:7 ethyl acetate/hexane) after 30 minutes revealed partial consumption of the starting material with formation of a slightly more polar product. Additional half equivalents of EDCI, NEt_3 , and $\text{H}_2\text{N-OTHP}$, were then added at 30-minute intervals until complete consumption of starting material was indicated by TLC. The reaction mixture was next treated with H_2O (5 mL) and after separation the organic layer dried over Na_2SO_4 . After solvent removal under vacuum, the product was isolated via flash chromatography (SiO_2 , 2:3 ethyl acetate/hexane, $R_f = 0.40$) to yield 188 mg (98%) of a colorless oil. ^1H NMR (CDCl_3 , 400 MHz) δ 7.97 (br s, 1H), 7.44-7.28 (m, 10H), 6.65 (br s, 1H), 5.15 (m, 2H), 4.99 (m, 1H), 4.33 (m, 2H), 3.94 (m, 1H), 3.60 (m, 1H), 1.90-1.80 (m, 2H), 1.70-1.50 (m, 4H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 152.8, 148.8, 138.3, 135.0, 128.7, 128.6, 128.5, 127.8, 127.3, 101.4, 67.7, 64.1, 45.2, 29.4, 25.1, 20.8; HRMS (EI, $\text{M}+\text{H}$) Calcd for $\text{C}_{21}\text{H}_{26}\text{N}_3\text{O}_4$ 384.1921, found 384.1923; See ^1H and ^{13}C NMR (pages S12-13) for purity assessment.

Compound 7

The improved preparation of this known compound⁶ involved treating KNCS (4.5 g, 46.3 mmol), stirred vigorously in C_2Cl_4 (100 mL), with 18-crown-6 (5%, 500 mg, 1.75 mmol) and benzyl chloroformate (5.0 mL, 35.0 mmol). The flask was fitted with a condenser and the mixture warmed to 85°C for the night. The following morning TLC analysis (1:4 CH_2Cl_2 /hexane) showed complete consumption of the benzyl chloroformate ($R_f = 0.45$) and formation of the more polar product ($R_f = 0.30$) along with trace amounts of the benzyl thiocyanate side product ($R_f = 0.10$). The mixture was then diluted with 100 mL hexane and cooled on ice before filtering through celite to remove salts (washed with 1:1 CH_2Cl_2 /hexane). After solvent

removal under vacuum 7.48 g of a bright yellow oil was obtained. NMR analysis of this material showed it to be the desired product CbzNCS in major proportion (83% estimated yield by ^1H NMR) along with trace amounts ($> 5\%$) of benzyl thiocyanate, benzyl isothiocyanate, and 18-crown-6. These trace impurities were found not to interfere in the subsequent use CbzNCS and are easily removed during the later chromatographic purification of the protected N^G -hydroxyguanidine products. The crude CbzNCS was dissolved in CH_2Cl_2 to a concentration of 0.5 M providing a conveniently used stock which when stored at 4°C showed no degradation after many weeks. Spectral data for the CbzNCS prepared in this way matched that of the previously described preparation.⁶

General Procedure and Data for Protected N^G -hydroxyguanidines in Table 1

General Procedure. To a solution of primary amine (0.5 mmol) in CH_2Cl_2 (10 mL) was added CbzNCS (0.5 mmol, 1.0 mL of a 0.5 M solution in CH_2Cl_2). TLC analysis (1:2 ethyl acetate/hexane, ninhydrin visualization) generally showed rapid consumption of the amine and formation of the less polar thiourea (R_f typically 0.50-0.60 using the afore mentioned solvent system). Once the amine was completely consumed the intermediate thiourea was treated directed with EDCI (100mg, 0.5 mmol), NEt_3 (70 μL , 0.5 mmol), and H_2N -OTHP (70 μL , 0.55 mmol). TLC analysis after 30 minutes showed the formation of product as a more polar species (R_f typically 0.35-0.45). To drive the reaction to completion additional half-equivalents of EDCI, NEt_3 , and H_2N -OTHP, were then added at 30-minute intervals until complete consumption of starting material was indicated by TLC. The reaction mixture was next treated with H_2O (5 mL) and after separation, the organic layer dried over Na_2SO_4 . After solvent removal under vacuum, the products were isolated via flash chromatography (SiO_2 , ethyl acetate/hexane).

Compound S1 (Table 1, Entry 1)

For complete spectral characterization see **compound 11** (page S5).

Compound S2 (Table 1, Entry 2)

Yield 95%; White solid; mp 52-54°C; ^1H NMR (CDCl_3 , 400 MHz) δ 7.93

(br s, 1H), 7.43-7.34 (m, 5H), 6.34 (br s, 1H), 5.15 (m, 2H), 4.95 (m, 1H), 3.90 (m, 2H), 3.57 (m, 1H), 2.02-1.91 (m, 2H), 1.87-1.74 (m, 2H), 1.72-

1.39 (m, 10H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 152.9, 148.8, 135.1, 128.7, 128.4, 101.5, 67.5, 64.1, 52.2, 33.1, 29.4, 25.2, 23.7, 23.6, 20.9; HRMS (EI, M+H) Calcd for $\text{C}_{19}\text{H}_{28}\text{N}_3\text{O}_4$ 362.2077, found 362.2080; Anal. Calcd for $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_4$: C, 63.14; H, 7.53; N, 11.63. Found: C, 62.93; H, 7.86; N, 11.29.

Compound S3 (Table 1, Entry 3)

Yield 96%; Colorless oil; ^1H NMR (CDCl_3 , 400 MHz) δ 7.95 (br s, 1H),

7.42-7.35 (m, 5H), 6.37 (br s, 1H), 5.14 (m, 2H), 4.95 (m, 1H), 3.89 (m, 1H), 3.56 (m, 1H), 1.88-1.75 (m, 2H), 1.68-1.51 (m, 4H), 1.37 (s, 9H); ^{13}C NMR

(CDCl_3 , 100 MHz) δ 153.0, 147.3, 135.2, 128.7, 128.6, 128.4, 101.4, 67.4, 64.0, 51.0, 29.4, 28.8, 25.2, 20.8; HRMS (EI, M+H) Calcd for $\text{C}_{18}\text{H}_{28}\text{N}_3\text{O}_4$ 350.2083, found 350.2080; Anal. Calcd for $\text{C}_{18}\text{H}_{27}\text{N}_3\text{O}_4$: C, 61.87; H, 7.79; N, 12.03. Found: C, 61.49; H, 7.74; N, 11.78.

Compound S4 (Table 1, Entry 4)

Yield 99%; White solid; mp 86-88°C; ^1H NMR (CDCl_3 , 400 MHz) δ

8.78 (br s, 1H), 8.10 (br s, 1H), 7.45-7.37 (m, 5H), 7.11 (br s, 2H), 6.63 (br s, 1H), 5.22 (m, 2H), 5.13 (m, 1H), 3.96 (m, 1H), 3.64 (m, 1H), 2.29

(s, 6H), 1.91-1.81 (m, 2H), 1.74-1.54 (m, 4H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 153.1, 145.7, 138.6, 138.3, 134.9, 128.8, 128.7, 128.6, 123.8, 116.9, 101.6, 68.0, 64.1, 29.3, 25.2, 21.2, 20.7; HRMS (EI, $\text{M}+\text{Li}$) Calcd for $\text{LiC}_{22}\text{H}_{27}\text{N}_3\text{O}_4$ 404.2171, found 404.2162; Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{N}_3\text{O}_4$: C, 66.48; H, 6.85; N, 10.57. Found: C, 66.18; H, 7.10; N, 10.22.

Compound S5 (Table 1, Entry 5)

Yield 97%; White solid; mp 104-106°C; ^1H NMR (CDCl_3 , 400 MHz)

δ 8.94 (br s, 1H), 8.06 (br s, 1H), 7.45-7.34 (m, 9H), 5.22 (m, 2H), 5.09 (m, 1H), 3.92 (m, 1H), 3.62 (m, 1H), 1.81-1.75 (m, 2H), 1.74-1.50

(m, 4H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 153.1, 145.1, 137.8, 134.7, 131.6, 128.9, 128.8, 128.6, 120.6, 114.1, 101.5, 68.1, 64.0, 29.2, 25.1, 20.6; HRMS (EI, $\text{M}+\text{H}$) Calcd for $\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_4\text{Br}$ 448.0873, found 448.0872; Anal. Calcd for $\text{C}_{20}\text{H}_{22}\text{N}_3\text{O}_4\text{Br}$: C, 53.58; H, 4.95; N, 9.37. Found: C, 53.58; H, 5.06; N, 9.27.

Compound S6 (Table 1, Entry 6)

Yield 98%; Colorless oil; 1:1 mixture of diastereomers; ^1H NMR

(CDCl_3 , 400 MHz) δ 7.82 (br m, 1H), 7.44-7.34 (m, 5H), 7.33-7.23 (m, 3H), 7.22-7.15 (m, 2H), 6.89 (br m, 1H), 5.17 (m, 2H), 4.91 (m,

1H), 4.54 (m, 1H), 3.86 (m, 1H), 3.68 (s, 3H), 3.56 (m, 1H), 3.19 (m, 1H), 3.10 (m, 1H), 1.85-1.71 (m, 2H), 1.66-1.50 (m, 4H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 172.6, 172.4, 152.5, 147.6, 136.4, 136.3, 135.0, 129.3, 128.7, 128.6, 128.43, 128.38, 101.4, 101.2, 67.8, 63.9, 63.7, 55.0, 54.9, 52.0, 37.8, 37.6, 29.2, 25.1, 20.6, 20.5; HRMS (EI, $\text{M}+\text{H}$) Calcd for $\text{C}_{24}\text{H}_{30}\text{N}_3\text{O}_4$ 456.2129, found 456.2135; See ^1H and ^{13}C NMR (page S14-15) for purity assessment.

Compound S7 (Table 1, Entry 7)

Yield 94%; Colorless oil; 1:1 mixture of diastereomers; ^1H NMR (CDCl_3 , 400 MHz) δ 7.89 (br m, 1H), 7.43-7.33 (m, 5H), 6.33 (br m, 1H), 5.14 (m, 3H), 4.92 (m, 1H), 4.15 (m, 1H), 3.89 (m, 1H), 3.56 (m, 1H), 3.12 (m, 2H), 1.88-1.74 (m, 3H), 1.71-1.50 (m, 7H), 1.46 (s, 9H), 1.44 (s, 9H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 171.8, 155.3, 152.8, 148.9, 135.0, 128.7, 128.5, 126.9, 101.3, 81.8, 79.5, 67.6, 65.3, 64.1, 55.7, 46.1, 40.5, 30.3, 30.2, 28.3, 28.0, 25.1, 24.9, 24.8, 20.9; HRMS (EI, M+H) Calcd for $\text{C}_{28}\text{H}_{45}\text{N}_4\text{O}_8$ 565.3244, found 565.3237; See ^1H and ^{13}C NMR (pages S16-17 for purity assessment).

Compound S8 (Table 1, Entry 8)

Yield 96%; Colorless oil; 1:1 mixture of diastereomers; ^1H NMR (CDCl_3 , 400 MHz) δ 7.85 (br m, 1H), 7.43-7.33 (m, 5H), 6.75 (br m, 1H), 6.01 and 5.75 (br m, 1H), 5.16 (m, 2H), 4.99 (m, 1H), 4.42 (m, 1H), 3.88 (m, 1H), 3.74 (m, 1H), 3.65-3.41 (m, 2H), 2.50-2.29 (m, 2H), 1.86-1.73 (m, 2H), 1.64-1.50 (m, 4H), 1.49-1.40 (m, 18H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 170.5, 170.4, 155.4, 155.3, 152.7, 148.4, 148.0, 134.8, 128.8, 128.7, 128.6, 101.1, 101.0, 82.0, 81.9, 79.7, 79.5, 68.0, 63.9, 63.7, 49.8, 45.2 (m), 35.1 (m), 29.14, 29.10, 28.4, 28.3, 27.84, 27.81, 25.1, 25.0, 20.8, 20.5; ^{19}F NMR (CDCl_3 , 376 MHz) δ -93.9 to -98.4 (m), -101.4 to -102.8 (m); HRMS (EI, M+H) Calcd for $\text{C}_{28}\text{H}_{43}\text{N}_4\text{O}_8\text{F}_2$ 601.3036, found 601.3049; See ^1H and ^{13}C NMR (page S18-19) for purity assessment.

Deprotection and Characterization of 12a,b

N^G-hydroxy-L-arginine (12a)

Following the method of Kiso and coworkers,⁷ the protected *N*^G-hydroxy-L-arginine (226 mg, 0.4 mmol) was treated with trifluoroacetic acid (15.0 mL, 202 mmol) and thioanisole (2.4 mL, 20 mmol) at room temperature. During the course of the reaction

the mixture became a bright green color, which faded to dark after 5 hours at which time the mixture was concentrated under vacuum. To remove nonpolar impurities the residue was partitioned between ether (15 mL) and H₂O (15 mL) and after separation, the ether layer further extracted with an additional 5 mL H₂O. The aqueous layers were combined, concentrated under vacuum (final volume ~ 5.0 mL), and purified using a Varian C18 megabond elut column (60 mL volume containing 10 g C18 silica). Prior to applying the crude product, the column was pre-equilibrated with methanol (50 mL) followed by 0.1% TFA in H₂O (50 mL). After loading the column, product was eluted with 0.1% TFA in H₂O and collected in 10 mL fractions. The first 3 fractions (positive ninhydrin test) were pooled, concentrated, and lyophilized to give the product as a clear, glassy foam (168 mg, quantitative yield). Spectral data in agreement with published values.⁸ ¹H NMR (D₂O, 400 MHz) δ 3.97 (t, *J* = 6.3 Hz, 1H), 3.14 (t, *J* = 6.8 Hz, 2H), 1.95-1.76 (m, 2H), 1.72-1.50 (m, 2H); ¹³C NMR (D₂O, 100 MHz) δ 171.5, 158.6, 52.3, 40.1, 26.8, 23.6; HRMS (EI, M+H) Calcd for C₆H₁₅N₄O₃ 191.1139, found 191.1144; See ¹H and ¹³C NMR (page S20-21) for purity assessment.

4,4-difluoro-*N*^G-hydroxy-L-arginine (12b)

Using deprotection conditions identical to those described for 12a the product was obtained in quantitative yield as a clear, glassy

foam. ^1H NMR (D_2O , 400 MHz) δ 4.32 (dd, J = 3.8, 7.9 Hz, 1H), 4.66 (t, J = 14.9 Hz, 2H), 2.80-2.43 (m, 2H); ^{13}C NMR (D_2O , 100 MHz) δ 170.4, 158.9, 121.5 (t, J = 244.3 Hz), 47.5, 44.9 (t, J = 27.5 Hz), 33.5 (t, J = 22.7 Hz); ^{19}F NMR (D_2O , 376 MHz) δ -103.8 (app. q, J = 246.9 Hz); HRMS (EI, M+H) Calcd for $\text{C}_6\text{H}_{13}\text{N}_4\text{O}_3\text{F}_2$ 227.0952, found 227.0956; See ^1H and ^{13}C NMR (page S22-23) for purity assessment.

NMR Data for Selected Compounds

Compound 11: ¹H NMR

Compound 11: ^{13}C NMR