Supporting Information

meso-Porphyrinyl-Substituted Expanded Porphyrins

Y. Inokuma and A. Osuka

General Procedure. All reagents and solvents were of the commercial reagent grade and were used without further purification except where noted. Dry toluene and dry CH₂Cl₂ were obtained by distilling over CaH₂. ¹H NMR spectra were recorded on a JEOL delta-600 spectrometer, and chemical shifts were reported as the delta scale in ppm relative to CHCl₃ (δ = 7.260). The spectroscopic grade CH₂Cl₂ and CHCl₃ were used as solvents for all spectroscopic studies. UV/visible/NIR absorption spectra were recorded on a Shimadzu UV-3100 spectrometer. Mass spectra were recorded on a JEOL HX-110 spectrometer, using positive-FAB ionization method with accelerating voltage 10 kV and a 3-nitrobenzylalcohol matrix, or on a Shimadzu/KRATOS KOMPACT MALDI 4 spectrometer, using positive-MALDI ionization method with/without CA matrix. Preparative separations were performed by silica gel flash column chromatography (Merck Kieselgel 60H Art. 7736), silica gel gravity column chromatography (Wako gel C-400) and size exclusion gel permeation chromatography (Bio-Rad Bio-Beads S-X1, packed with toluene or CHCl₃ in a 4 × 100 gravity flow column: flow rate, 3.8 mLmin⁻¹). Recycling preparative GPC-HPLC was carried out on JAI LC-908 using preparative JAIGEL-2.5H, 2.5H, and 3H columns. Redox potentials were measured by the cyclic voltammetry method on an ALS electrochemical analyzer model 660.

5-(4-(5,5-Dimethyl-1,3-dioxan-2-yl)-2,3,5,6-tetrafluorophenyl)-2,8,12,18-tetraethyl-3,7,13,17-tetramethyl-15-phenylporphyrin (3c) and 5-(4-formyl-2,3,5,6-tetrafluorophenyl)-2,8,12,18-tetraethyl-3,7,13,17-tetramethyl-15-phenylporphyrin (3a)

Benzaldehyde (0.110 mL, 1.09 mmol) and 2,3,5,6-tetrafluoro-4-(5,5-dimethyl-1,3-dioxan-2-yl)benzaldehyde (**1a**, 317 mg, 1.09 mmol) were dissolved in dry acetonitrile (15 mL). A solution of dry

trichloroacetic acid (21.0 mg, 0.129 mmol) in dry acetonitrile (2 mL) was added and the mixture was stirred under argon atmosphere in the dark. Then a solution of bis(3-ethyl-4-methyl-2-pyrrolyl)methane (**2**, 500 mg, 2.17 mmol) in dry acetonitrile (3 mL) was added and resulting the mixture was stirred for 5.5 h. A suspension of *p*-chloranil (790 mg, 3.21 mmol) in 15 mL of dry THF was added and the mixture was stirred overnight. The reaction mixture was passed through a short alumina column and the solvent was removed by a rotary evaporator. A fraction that contained porphyrin products was separated by silica gel column chromatography with CH₂Cl₂ as an eluent and was further separated by GPC (gel permeation chromatography) with THF as an eluent. The second fraction was collected and further purified by silica gel column chromatography with a 1:1 mixture of AcOEt and hexane, giving **3c** (146 mg, 16.5%). The acetal porphyrin **3c** (100 mg) was dissolved in a mixture of TFA (5 mL), 10% HCl (5 mL), and one drop of concentrated H₂SO₄. The resulting mixture was refluxed for 3 h moderately. The reaction solution was cooled to room temperature, and then neutralized with saturated NaHCO₃ aqueous solution. Organic phase was collected, washed with brine and dried over Na₂SO₄. Removal of the solvent gave porphyrin **3a** quantitatively.

3c: ¹H-NMR (600 MHz, CDCl₃, 298 K) δ 10.25 (s, 2H, *meso*), 8.04 (d, 2H, phenyl), 7.79 (t, 1H, phenyl), 7.73 (t, 2H, phenyl), 6.15 (s, 1H, benzyl), 4.02 (q, 8H, β-CH₂CH₃), 2.72 (s, 6H, β-CH₃), 2.46 (s, 6H, β-CH₃), 1.78 (m, 12H, β-CH₂CH₃), 1.42 (s, 3H, CH₃), 0.92 (s, 3H, CH₃), -2.39 (br, 1H, inner NH), and -2.43 (br, 1H, inner NH); ¹³C-NMR (150 MHz, CDCl₃, 298 K) δ 145.7, 145.2, 144.8, 142.1, 141.6, 141.3, 136.5, 133.8, 132.9, 128.5, 127.8, 119.7, 98.1, 97.2, 95.5, 78.3, 30.7, 23.0, 22.0, 20.1, 20.0, 17.7, 17.6, 14.4, and 13.6; ¹⁹F-NMR (565 MHz, CDCl₃, 298 K) δ -137.5 (dd, *J* = 13.8 Hz, 10.4 Hz, 2F) and -143.1 (dd, *J* = 13.8 Hz, 10.3 Hz, 2F); m.p. >300 °C (decomp.); FAB-MS (3-nitrobenzyl alcohol) *m/z* = 816.4 (Calcd for C₅₀H₅₂N₄F₄O₂ = 816.4).

3a: ¹H NMR (600 MHz, CDCl₃, 298 K) δ = 10.69 (s, 1H, -CHO), 10.26 (s, 2H, *meso*), 8.06 (d, *J* = 6.0 Hz, 2H, phenyl), 7.82 (t, *J* = 15.6 Hz, 1H, phenyl), 7.76 (t, *J* = 15.0 Hz, 2H, phenyl), 4.04 (m, 8H, β-CH₂CH₃), 2.76 (s, 6H, β-CH₃), 2.48 (s, 6H, β-CH₃), 1.85 (m, 12H, β-CH₂CH₃), and -2.39 (br, 2H, inner NH); ¹³C-NMR (150 MHz, CDCl₃, 298 K) δ 182.9, 146.1, 145.4, 145.0, 144.1, 141.8, 141.3, 136.8, 133.0, 132.8, 128.6, 127.9, 120.1, 115.7, 97.5, 96.8, 20.1, 20.0, 17.7, 17.6, 14.4, and 13.7; ¹⁹F-NMR (565 MHz,

CDCl₃, 298 K) δ -135.7 (dd, J = 13.8 Hz, 9.5 Hz, 2F), and -145.8 (dd, J = 13.8 Hz, 9.5 Hz, 2F); m.p. >300 °C (decomp.); FAB-MS (3-nitrobenzyl alcohol) m/z = 730.3 (Calcd for C₄₅H₄₂N₄F₄O = 730.3).

5-(4-formyl-2,3,5,6-tetrafluorophenyl)-2,8,12,18-tetraethyl-3,7,13,17-tetramethyl-15-(3,5-dioctyloxyphenyl)porphyrin (3b)

Porphyrin **3b** was prepared in 29% yield according to the same procedure used for **3a** with replacement of benzaldehyde by 3,5-dioctyloxybenzaldehyde.

3b: ¹H NMR (600 MHz, CDCl₃, 298 K) δ 10.69 (s, 1H, -CHO), 10.27 (s, 2H, *meso*), 7.21 (s, 2H, phenyl), 6.92 (s, 1H, phenyl), 4.11 (dq, 4H, -OCH₂CH₂-), 4.02 (q, 8H, β -CH₂CH₃), 2.76 (s, 6H, β -CH₃), 2.69 (s, 6H, β -CH₃), 1.81 (m, 12H, β -CH₂CH₃), 1.5–1.2 (m, 30H, octyloxy) and -2.38 (br, 2H, inner NH); ¹³C-NMR (150 MHz, CDCl₃, 298 K) δ 183.0, 159.9, 146.1, 145.2, 145.0, 144.1, 143.4, 141.8, 141.3, 136.9, 133.0, 120.1, 112.5, 102.2, 97.5, 31.9, 29.5, 29.3, 26.1, 22.7, 20.0, 17.8, 17.7, 14.3, 14.2 and 13.7; ¹⁹F-NMR (565 MHz, CDCl₃, 298 K) δ -135.7 (dd, J = 13.4 Hz, 9.4 Hz, 2F), and -145.7 (dd, J = 13.8 Hz, 9.5 Hz, 2F); FAB-MS (3-nitrobenzyl alcohol) m/z = 986.6 (Calcd for C₆₁H₇₄N₄F₄O₃ = 986.6).

***meso*-Porphyrinyl-substituted expanded porphyrins (4-6, heptaphyrin 7', and octaphyrin 8')**

The porphyrin **3a** (89.2 mg, 122 μ mol) was dissolved in distilled CH₂Cl₂ (200 mL). To this solution were added 0.847 mL of 144 mM pyrrole in distilled CH₂Cl₂ and 0.420 mL of 873 mM BF₃·OEt₂ in distilled CH₂Cl₂ and the resulting mixture was stirred under argon atmosphere for 2 h in the dark. Then, DDQ (64 mg) was added and the reaction mixture was stirred overnight. The reaction mixture was washed with saturated NaHCO₃ solution and brine, and the organic extract was then dried over anhydrous Na₂SO₄ and evaporated. The dark residual mixture was first separated by silica gel column chromatography (CH₂Cl₂ : hexane = 7:3). First fraction contained unreacted starting porphyrin **3a**, second fraction contained many expanded porphyrin products including **6**, **7'**, **8'**, and higher homologues, and third fraction contained pentakis(porphyrinyl) N-fused pentaphyrin **5-ox** (7.78 mg, 8.2%). The second fraction was further separated by GPC column chromatography (solvent: CHCl₃), to give tetrakis(porphyrinyl)porphyrin **4** (1.69 mg, 1.78%), hexakis(porphyrinyl)hexaphyrin **6-red** (10.84 mg, 11.4%), heptakis(porphyrinyl)heptaphyrin **7'** (2.93 mg, 3.09%), and

octakis(porphyrinyl)octaphyrin **8'** (1.55 mg, 1.63%). As judged from the color and eluting behavior, the hexaphyrin **6** existed as its oxidized form **6-ox** in this step, and was reduced to **6-red** during separation over recycling GPC-HPLC with CHCl₃ containing a small amount of ethanol.

***meso*-tetratakis(porphyrinyl) substituted porphyrin (4)**

¹H NMR (400 MHz, CDCl₃ and TFA, 298 K): δ 10.37 (s, 8H, porphyrinyl-*meso*), 9.53 (br d, 8H, core porphyrin-β-H), 8.35 (br d, *J* = 6.8 Hz, 8H, phenyl), 7.97 (br, 12H, phenyl), 3.97 (br q, 16H, β-CH₂CH₃), 3.77 (br q, 16H, β-CH₂CH₃), 3.19 (br s, 24H, β-CH₃), 2.28 (br s, 24H, β-CH₃), 1.59 (br t, 24H, β-CH₂CH₃), 1.42 (br t, 24H, β-CH₂CH₃), -1.49, and -1.62 (br, 20H, porphyrin-inner NH (protonated)); ¹³C-NMR (150 MHz, CDCl₃ and TFA, 298 K) δ 147.8 (br), 146.3, 146.1, 143.8, 143.7, 142.2, 142.0, 141.5, 141.0, 138.0, 136.8, 135.4, 133.1, 131.3, 130.1, 129.5, 98.6, 20.0, 15.7, 15.6, 13.0, and 12.6; ¹⁹F-NMR (565 MHz, CDCl₃ and TFA, 298 K) -135.9 (br, 8F), and -137.2 (br, 8F); m.p. >300 °C (decomp.); MALDI TOF-Mass *m/z* = 3111 (calcd for C₁₉₆H₁₇₄N₂₀F₁₆ = 3111). UV-vis (CH₂Cl₂): λ_{max} (ε [mol⁻¹dm³cm⁻¹]) = 411 (832000), 423 (1010000), 508 (111000), 540 (40000), 575 (40000), 626 (16000)

***meso*-Pentakis(porphyrinyl) substituted [22]N-fused pentaphyrin (5-ox)**

¹H NMR (600 MHz, CDCl₃, 298 K): δ 10.48 (s, 1H, porphyrinyl-*meso*), 10.46 (s, 1H, porphyrinyl-*meso*), 10.44 (s, 1H, porphyrinyl-*meso*), 10.43 (s, 2H, porphyrinyl-*meso*), 10.33 (s, 2H, porphyrinyl-*meso*), 9.68 (s, 1H, pentaphyrin outer-β-H), 9.60 (s, 1H, porphyrinyl-*meso*), 9.52 (s, 1H, porphyrinyl-*meso*), 9.36 (s, 1H, porphyrinyl-*meso*), 9.56 (d, *J* = 6.2 Hz, 1H, pentaphyrin outer-β-H), 9.14 (d, *J* = 4.1 Hz, 1H, pentaphyrin outer-β-H), 9.11 (d, *J* = 6.2 Hz, 1H, pentaphyrin outer-β-H), 9.04 (d, *J* = 4.1 Hz, 1H, pentaphyrin outer-β-H), 8.99 (d, *J* = 4.8 Hz, 1H, pentaphyrin outer-β-H), 8.21, 8.14 (m, 10H, phenyl), 8.08, 7.86, 7.81 (m, 15H, phenyl), 4.38 (m, 12H, β-CH₂CH₃), 4.13 (m, 12H, β-CH₂CH₃), 4.05 (m, 8H, β-CH₂CH₃), 3.91 (m, 8H, β-CH₂CH₃), 3.64 (br s, 3H, β-CH₃), 3.58 (br s, 3H, β-CH₃), 3.52 (br s, 3H, β-CH₃), 3.51 (br s, 3H, β-CH₃), 3.48 (br s, 3H, β-CH₃), 3.47 (br s, 3H, β-CH₃), 3.42 (d, *J* = 4.1 Hz, 1H, pentaphyrin inner-β-H), 3.32 (br s, 6H, β-CH₃), 3.12 (br s, 3H, β-CH₃), 3.01 (br s, 3H, β-CH₃), 2.88 (d,

$J = 4.1$ Hz, 1H, pentaphyrin inner- β -H), 2.60 (br s, 3H, β -CH₃), 2.56 (br s, 3H, β -CH₃), 2.54 (br s, 3H, β -CH₃), 2.50 (br s, 3H, β -CH₃), 2.45 (br s, 3H, β -CH₃), 2.40 (br s, 3H, β -CH₃), 2.36 (br s, pentaphyrin-
inner NH), 2.17 (br s, 3H, β -CH₃), 2.02 (br s, 3H, β -CH₃), 2.08 (m, 12H, β -CH₂CH₃), 1.87 (m, 12H, β -
CH₂CH₃), 1.80 (t, 6H, β -CH₂CH₃), 1.64 (m, 6H, β -CH₂CH₃), 1.19 (t, 3H, β -CH₂CH₃), 0.34 (t, 3H, β -
CH₂CH₃), 0.30 (t, 3H, β -CH₂CH₃), -0.03 (t, 3H, β -CH₂CH₃), -0.48 (s, 1H, pentaphyrin inner- β -H),
-1.97, -2.10, -2.15, -2.21, and -2.32 (br s, 10H, porphyrin-inner NH); ¹⁹F-NMR (565 MHz, CDCl₃, 298
K) δ -135.3 (br, 2F), -136.0 (br, 1F), -136.1 (br, 2F), -136.9 (br, 3F), -137.5 (br, 1F), -137.7 (br, 1F),
-137.9 (br, 3F), -138.2 (br, 1F), -138.7 (br, 1F), -138.9 (br, 1F), -139.1 (br, 2F), -139.4 (br, 1F), and
-140.2 (br, 1F); m.p. >300 °C (decomp.); MALDI TOF-Mass m/z = 3885 (calcd for C₂₄₅H₂₁₅N₂₅F₂₀ =
3887). UV-vis (CH₂Cl₂): λ_{max} (ϵ [mol⁻¹dm³cm⁻¹]) = 408 (723000), 507 (121000), 531 (62000), 574
(51000), 626 (21000)

***meso*-Pentakis(porphyrinyl) substituted [24]N-fused pentaphyrin (5-red)**

meso-Pentakis(porphyrinyl) substituted [22]N-fused pentaphyrin **5-ox** was dissolved in a 10:1 mixture
of CH₂Cl₂ and MeOH. 50 equivalent of NaBH₄ was added to the solution and the resulting solution
was stirred for 3 h. The solution was washed with water, and the organic phase was separated, dried
over anhydrous Na₂SO₄. Removal of the solvent gave **5-red** quantitatively.

5-red: ¹H NMR (600 MHz, CDCl₃, 298 K): δ 12.2 (br, 1H, pentaphyrin NH), 11.70, 10.28, 10.38, 10.37,
10.36, 10.32, 10.26, 9.56, 9.52, 9.39 (s, 1H each, porphyrinyl-*meso*), 10.02, 9.46, 8.20–7.59, 7.10–6.95
(m, 28H, pentaphyrin β -H and porphyrin phenyl), 8.77 (br, 1H, pentaphyrin NH), 7.55 (br, 1H,
pentaphyrin NH), 8.31 (s, 1H, pentaphyrin β -H), 7.28 (d, $J = 3.7$ Hz, 1H, pentaphyrin b-H), 6.40 (d, J
= 4.1 Hz, 1H, pentaphyrin β -H), 6.27 (d, $J = 5.5$ Hz, 1H, pentaphyrin β -H), 6.17 (d, $J = 4.6$ Hz, 1H,
pentaphyrin β -H), 5.66 (d, $J = 4.1$ Hz, 1H, pentaphyrin β -H), 4.25–4.00, 3.62–3.35 (m, 40H, β -
CH₂CH₃), 3.18–2.88, 2.58–2.18 (m, 60H, β -CH₃), 1.99–1.75, 0.43–0.13 (m, 60H, β -CH₂CH₃), and
-2.10–2.60 (br, porphyrin inner-NH); ¹⁹F-NMR (565 MHz, CDCl₃, 298 K) δ -135.3 (br, 2F), -135.5 (br,
2F), -135.7 (br, 1F), -135.9 (br, 2F), -136.7 (br, 2F), -136.9 (br, 1F), -138.1 (br, 1F), -138.5 (br, 2F),

–138.7 (br, 2F), –139.0 (br, 1F), –139.3 (br, 1F), –140.4 (br, 2F), and –140.8 (br, 1F); m.p. >300 °C (decomp.); MALDI TOF-MS m/z = 3888 (calcd for $C_{245}H_{217}N_{25}F_{20}$ = 3889). $\lambda_{\max}$ (ϵ [$\text{mol}^{-1}\text{dm}^3\text{cm}^{-1}$]) = 408 (766000), 507 (128000), 535 (60000) 573 (46000), 619 (12000)

***meso*-Hexakis(porphyrinyl) [28]hexaphyrin (6-red)**

6-red: ^1H NMR (600 MHz, CDCl_3 , 298 K): δ 10.41 (s, 4H, porphyrinyl-*meso*), 10.3–9.80 (br, 8H, porphyrinyl-*meso*), 9.75 (br, 2H, hexaphyrin NH), 8.52, 8.40 (br, 8H, hexaphyrin β -H), 8.15–7.53 (m, 30H, phenyl), 5.28 (br, 2H, hexaphyrin NH), 4.27, 4.11 (br, 48H, β - CH_2CH_3), 3.35–2.40 (br, 72H, β - CH_3), 2.10–1.30 (br, 48H, β - CH_2CH_3), and –2.15, –2.25, –2.35, –2.44 (br, 12H, porphyrinyl inner-NH); ^{19}F -NMR (565 MHz, CDCl_3 , 298 K) δ –135.7 (br, 8F), –136.8 (br, 4F), –137.9 (br, 4F), and –138.6 (br, 8F); m.p. >300 °C (decomp.); MALDI TOF-Mass m/z = 4667 (calcd for $C_{294}H_{262}N_{30}F_{24}$ = 4668). $\lambda_{\max}$ (ϵ [$\text{mol}^{-1}\text{dm}^3\text{cm}^{-1}$]) = 408 (1370000), 507 (148000), 541 (89000), 603 (188000)

***meso*-Hexakis(porphyrinyl) [26]hexaphyrin (6-ox)**

The *meso*-hexakis(porphyrinyl)-substituted [28]hexaphyrin **6-red** was dissolved in CH_2Cl_2 . To this solution, DDQ (5 equiv.) was added and the mixture was stirred for 15 minutes. The resulting solution was passed through a short alumina column, and the solvent was evaporated. Recrystallization from a mixture of CH_2Cl_2 and MeOH gave **6-ox** quantitatively.

6-ox: ^1H NMR (600 MHz, CDCl_3 , 298 K): δ 10.43 (br, 12H, porphyrinyl-*meso*), 9.61 (br, 4H, hexaphyrin outer- β -H), 9.04 (br, 4H, hexaphyrin outer- β -CH), 8.14, 8.01 (br d, 12H, phenyl), 7.86, 7.82, 7.77 (br t, 18H, phenyl), 4.33, 4.12, 3.48, 3.03 (br, 48H, β - CH_2CH_3), 2.54, 2.42 (br, 72H, β - CH_3), 2.06, 1.86 (br, 48H, β - CH_2CH_3), 0.56 (br, 4H, hexaphyrin inner- β -H), and –2.13, –2.19, –2.24, –2.30 (br, 14H, porphyrin and hexaphyrin inner-NH); ^{19}F -NMR (565 MHz, CDCl_3 , 298 K) δ –135.9 (br, 4F), –137.2 (br, 8F), –137.6 (br, 4F), and –138.7 (br, 8F); m.p. >300 °C (decomp.); MALDI TOF-Mass m/z = 4665 (calcd for $C_{294}H_{260}N_{30}F_{24}$ = 4666). $\lambda_{\max}$ (ϵ [$\text{mol}^{-1}\text{dm}^3\text{cm}^{-1}$]) = 408 (1110000), 597 (112000), 541

(105000), 580 (192000)

Fig. S1 Chromatogram of 4, 5-ox, 6-red, 7, and 8

column : JAIGEL 2H, 2.5H, 3H ; solvent : THF (no stabilizer) ; flow rate = 1.2 ml/min

Figure S2. MALDI TOF-Mass spectra of 4, 5, 6, 7' and 8'.

tetrakis(porphyrinyl)porphyrin 4

calculated for : $C_{196}H_{174}N_{20}F_{16} = 3111$

observed : 3111

pentakis(porphyrinyl)N-fused pentaphyrin(ox.) **5-ox**

calculated for : $C_{245}H_{215}N_{25}F_{20} = 3887$

observed : 3885

hexakis(porphyrinyl)hexaphyrin(ox.) **6-ox**

calculated for : $C_{294}H_{260}N_{30}F_{24} = 4666$

observed : 4667

heptakis(porphyrinyl)heptaphyrin **7'**

calculated for : $C_{343}H_{305}N_{35}F_{28} = 5445$

observed : 5442

octakis(porphyrinyl)octaphyrin 8'

calculated for : $C_{392}H_{348}N_{40}F_{32} = 6223$

observed : 6222

Fig. S3 ^{19}F NMR spectra of 4, 5-ox, 6-ox

4 (in CDCl_3 and TFA, at 298 K)

5-ox (in CDCl_3 at 298 K)

6-ox (in CDCl_3 , at 298 K)

Fig. S4 ^{13}C NMR spectra of 4 (in CDCl_3 and TFA)