***Trans*-Hydroalumination/Alkylation: One-Pot Synthesis of Trisubstituted Allylic Alcohols**

Neil F. Langille and Timothy F. Jamison*
*Department of Chemistry, Massachusetts Institute of Technology
Cambridge, MA 02139*

Supporting Information

General Methods:

All reactions were performed in oven-dried glassware, under an oxygen-free argon atmosphere with rigid exclusion of moisture from reagents and solvents. Vitride (Red-Al[®] sodium bis(2-methoxyethoxy)aluminum hydride, ≥ 65 wt. % in toluene) and diisobutylaluminum hydride (reagent grade) were purchased from Aldrich Chemical Co., and used without further purification. Iodomethane and iodoethane were purified by washing with aqueous sodium thiosulfate, water, and then aqueous sodium carbonate, followed by distillation and storage over copper powder. Methyllithium and *n*-butyllithium were titrated prior to use (menthol, Et₂O, 0 °C, 1,10-phenanthroline indicator). 2,6-Lutidine was distilled over calcium hydride and stored over solid potassium hydroxide. Chlorotrimethylsilane and dichloromethane were distilled from calcium hydride. Tetrahydrofuran and ethyl ether were distilled from a purple solution of sodium benzophenone ketyl. All other reagents were used as supplied by the chemical manufacturers.

Analytical thin layer chromatography (TLC) was performed on silica gel 60 F₂₅₄ aluminum plates precoated with a fluorescent indicator. Visualization of the developed chromatogram was accomplished with UV light (254 nm) and stained with either iodine-silica or vanillin. Data are reported as follows: R_f (solvent mixture, UV activity, color (stain)). Liquid chromatography was performed using a forced air-flow (flash chromatography^{S1}) on silica gel (Silicycle, 230-400 mesh), using the indicated solvent (underline denotes compound elution with indicated

(S1) Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923-2925.

solvent mixture(s)²). ¹H and ¹³C NMR spectra were recorded, unless otherwise noted, in CDCl₃ on a Varian Inova 500 MHz spectrometer. Chemical shifts are reported in parts per million using the solvent internal standard (chloroform, 7.27 and 77.23 ppm, respectively). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, app = apparent, b = broad), coupling constant (in Hz). Infrared spectra were recorded on a Perkin-Elmer 2000 FT-IR. High resolution mass spectra were obtained on a Bruker Daltonics APEXII 3 Tesla Fourier Transform Mass Spectrometer by Dr. Li Li of the MIT Department of Chemistry Instrumentation Facility.

Standard Procedures:

Copper iodide triethylphosphite. Following Nishizawa's procedure,^{S3} copper (I) iodide (5.00 g, 26.3 mmol) was added, slowly in portions over 10 min, to a rt solution of triethylphosphite (4.50 mL, 26.3 mmol, 100 mol %) in benzene (26.3 mL, 1.00 M), maintaining the reaction temperature at approximately 25 °C. After the addition was complete, the reaction mixture was stirred, protected from light, for 1 h. The crude mixture was filtered through Celite and concentrated *in vacuo*. The product was recrystallized from cyclohexane to afford 5.40 g (57.7%) desired product (**SI1**), displaying spectral properties in accord with literature precedent. The white crystalline material proved stable to storage at rt, protected from light, in a dessicator for several months.^{S4} ¹H NMR (500 MHz, CDCl₃) δ 4.12 (6H, q, *J* = 7.0), 1.30 (9H, t, *J* = 7.0).^{S5} ¹³C NMR (125 MHz, CDCl₃) δ 61.1, 16.8.

(S2) Example: (silica; 2, 5, 10% EtOAc/hex) denotes silica as chromatography sorbent, 2% EtOAc/hex to elute impurities, desired compound elution at 5% and 10% EtOAc/hex.

(S3) Nishizawa, Y. *Bull. Chem. Soc. Jpn.* **1961**, *34*, 1170-1178.

(S4) Decomposition of CuI·P(OEt)₃ could be first noted by decreased solubility in THF.

(S5) Compare to P(OEt)₃: ¹H NMR (500 MHz, CDCl₃) δ 3.88 (6H, q, *J* = 7.0), 1.28 (9H, t, *J* = 7.0).

General procedure for hydroalumination/alkylation with CuI•P(OEt)₃:

A solution of propargyl alcohol (1.0 mmol) in Et₂O (2.0 mL, 0.50 M) was cooled to 0 °C, and Vitride (≥65 wt. % in toluene, 0.69 mL, 227 mol %) was added dropwise. The cold bath was removed and the reaction stirred for 6 h. The flask was cooled to 0 °C and MeLi (1.6 M in Et₂O, 0.78 mL, 125 mol %) was added dropwise. The cold bath was removed and, after stirring for 90 min, the reaction was cooled to –78 °C (CO₂/acetone). A solution of CuI•P(OEt)₃ (357 mg, 100 mol %) in THF (5.0 mL, 0.20 M) was added by cannula (exothermic!), followed by addition of neat electrophile (300 mol %). The reaction was stirred for 20 h, allowing the cold bath to warm slowly to rt. The crude reaction mixture was poured onto sat. aq. potassium sodium tartrate (50 mL), rinsed with 40 mL EtOAc, and the mixture was stirred for 30 min. The layers were separated, and the aqueous layer was extracted 2× into 40 mL EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification by flash chromatography was generally required to afford pure allylic alcohol products

General procedure for hydroalumination/alkylation with CuCl:

A solution of propargyl alcohol (1.0 mmol) in Et₂O (2.0 mL, 0.50 M) was cooled to 0 °C, and Vitride (≥65 wt. % in toluene, 0.69 mL, 227 mol %) was added dropwise. The cold bath was removed and the reaction stirred for 6 h. The flask was cooled to 0 °C and MeLi (1.6 M in Et₂O, 0.78 mL, 125 mol %) was added dropwise. The cold bath was removed and, after stirring for 90 min, the reaction was cooled to –78 °C (CO₂/acetone). THF (5.0 mL, 0.20 M) was added by syringe, followed by addition of neat electrophile (300 mol %) and solid CuCl (99 mg, 100 mol %). The reaction was stirred for 20 h, allowing the cold bath to warm slowly to rt. The crude reaction mixture was poured onto sat. aq. potassium sodium tartrate (50 mL), rinsed with 40 mL EtOAc, and the mixture was stirred for 30 min. The layers were separated, and the aqueous layer was extracted 2× into 40 mL EtOAc. The combined organic layers were washed with

brine, dried over MgSO_4 , filtered, and concentrated *in vacuo*. Purification by flash chromatography was generally required to afford pure allylic alcohol products.

Experimental Data and Procedures for New Compounds:

5-(Triisopropylsilyloxy)pent-2-yn-1-ol (**1**).

To a 0 °C solution of 3-butyn-1-ol (2.00 mL, 26.4 mmol) in CH_2Cl_2 (66.0 mL, 0.400 M) were added sequentially: 2,6-lutidine (6.95 mL, 63.4 mmol, 240 mol %) and TIPS-OTf (8.55 mL, 31.7 mmol, 120 mol %). The reaction was stirred at 0 °C for 4 h, then quenched by addition of 50 mL of water. The layers were separated, and the aqueous layer was extracted 3× into 50 mL CH_2Cl_2 . The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated *in vacuo*. Purification by flash chromatography (silica; 0, 2% EtOAc/hex) provided the desired TIPS-protected alcohol as a colorless oil (5.80 g, 97.0%). R_f 0.90 (5% EtOAc/hex), no UV, brown (I_2 stain).

Following Journet and Cai's procedure,^{S6} the TIPS ether was dissolved in 64.0 mL THF (0.400 M) and the solution was cooled to −40 °C. *n*-Butyllithium (2.4 M in hexanes, 13.9 mL, 33.3 mmol, 130 mol %) was added dropwise, and the mixture was stirred for 5 min. *N,N*-Dimethylformamide (4.36 mL, 56.3 mmol, 220 mol %) was added in one portion. After 1-2 min, the cold bath was removed and the reaction was allowed to stir 30 min. Separately, the following were combined in a beaker, cooled to 0 °C, and stirred vigorously: 11.6 g KHPO_4 , 104 mL H_2O , and 96.0 mL MTBE. After 30 min, the reaction mixture was carefully poured onto the aq. KHPO_4 /MTBE mix, and stirred for 5 min. The layers were separated, and the aqueous layers were extracted 3× into 100 mL MTBE. The combined organic layers were dried over MgSO_4 , filtered, concentrated *in vacuo*,

(S6) Journet, M.; Cai, D.; DiMichele, L. M.; Larsen, R. D. *Tetrahedron Lett.* **1998**, 39, 6427-6428.

and used crude in the next step. R_f 0.80 (20% EtOAc/hex), UV active, purple (vanillin).

The crude aldehyde was dissolved in 128 mL MeOH (0.200 M) and cooled to 0 °C. Solid NaBH₄ (1.94 g, 200 mol %) was added carefully, in portions over 5 min. After 15 min at 0 °C, the reaction was carefully quenched with 50% aq. NH₄Cl to a pH of 5. The organic layer was extracted 3× into EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography (silica; 5, 10, 20% EtOAc/hex) provided the title compound **1** as a colorless oil (6.43 g, 97.9% yield over 2 steps). R_f 0.20 (20% EtOAc/hex), no UV, green (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 4.25 (2H, t, J = 2.1), 3.81 (2H, t, J = 7.3), 2.48 (2H, tt, J = 2.1, 7.3), 1.51 (1H, bs), 1.06-1.07 (21H, m). ¹³C NMR (125 MHz, CDCl₃) δ 83.5, 79.7, 62.3, 51.5, 23.4, 18.1, 12.1. IR (thin film NaCl) 3442, 2968, 2240, 1210, 1050 cm⁻¹. HRMS ESI (m/z): [M+Na]⁺ calcd for C₁₄H₂₈O₂SiNa, 279.1756; found, 279.1750.

(*E*)-3-(2-(Triisopropylsilyloxy)ethyl)hexa-2,5-dien-1-ol (2a). Reaction was conducted with alkynol **1** (256 mg, 1.00 mmol) and allyl bromide (0.26 mL, 3.00 mmol, 300 mol %), using standard CuI•P(OEt)₃ or CuCl procedure. Purification by flash chromatography (silica; 5, 10% EtOAc/hex) provided the title compound (86% and 97% yield, respectively). R_f 0.80 (30% EtOAc/hex), faint UV, blue (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 5.77 (1H, ddt, J = 6.2, 10.0, 17.1), 5.55 (1H, t, J = 6.9), 5.06 (1H, ddd, J = 1.8, 3.5, 17.1), 5.03 (1H, ddd, J = 1.5, 3.3, 10.0), 4.17 (2H, d, J = 6.9), 3.77 (2H, t, J = 7.1), 2.87 (2H, d, J = 6.2), 2.30 (2H, t, J = 7.1), 1.36 (1H, bs), 1.05-1.06 (21H, m). ¹³C NMR (125 MHz, CDCl₃) δ 138.5, 136.1, 126.7, 115.8, 62.7, 59.2, 40.5, 35.5, 18.2, 12.1.

IR (thin film NaCl) 3468, 3081, 2928, 1642, 1218, 1030 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{34}\text{O}_2\text{SiNa}$, 321.2226; found, 321.2222.

(*E*)-6-Methyl-3-(2-(triisopropylsilyloxy)ethyl)hepta-2,5-dien-1-ol (2b).

Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and 3,3-dimethylallyl bromide (0.17 mL, 1.50 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 2, 5% EtOAc/hex) provided the title compound (81% and 82% yield, respectively). R_f 0.30 (20% EtOAc/hex), faint UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.46 (1H, t, $J = 6.9$), 5.02 (1H, t, $J = 7.1$), 4.17 (2H, d, $J = 6.9$), 3.76 (2H, t, $J = 7.2$), 2.79 (2H, d, $J = 7.1$), 2.27 (2H, t, $J = 7.2$), 1.69 (3H, s), 1.66 (3H, s), 1.18 (1H, bs), 1.06 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 143.4, 138.1, 127.3, 111.6, 62.7, 59.3, 40.4, 39.3, 22.6, 18.2, 12.1. IR (thin film NaCl) 3460, 3060, 2941, 1380, 1365, 1212, 1030 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{19}\text{H}_{38}\text{O}_2\text{SiNa}$, 349.2539; found, 349.2537.

(*E*)-5-Methyl-3-(2-(triisopropylsilyloxy)ethyl)hexa-2,5-dien-1-ol (2c). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and 3-bromo-2-methylpropene (0.15 mL, 1.50 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 4, 6% EtOAc/hex) provided the title compound (84% and 94% yield, respectively). R_f 0.30 (20% EtOAc/hex), faint UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.62 (1H, t, $J = 6.5$), 4.79 (1H, d, $J = 0.7$), 4.72 (1H, d, $J = 0.7$), 4.15 (2H, t, $J = 6.5$), 3.77 (2H,

t, $J = 7.0$), 2.81 (2H, s), 2.27 (2H, t, $J = 7.0$), 1.70 (3H, s), 1.05-1.06 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 143.4, 138.1, 127.3, 111.6, 62.7, 59.3, 40.4, 39.3, 22.6, 18.2, 12.1. IR (thin film NaCl) 3462, 3078, 2929, 1220, 1035 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{18}\text{H}_{36}\text{O}_2\text{SiNa}$, 335.2382; found, 335.2386.

(*E*)-3-Benzyl-5-(triisopropylsilyloxy)pent-2-en-1-ol (2d). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and benzyl bromide (0.18 mL, 1.50 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 4, 6% EtOAc/hex) provided the title compound (62% and 60% yield, respectively). R_f 0.30 (20% EtOAc/hex), faint UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 7.19-7.31 (5H, m), 5.66 (1H, t, $J = 7.0$), 4.29 (2H, m), 3.72 (2H, t, $J = 6.9$), 3.48 (2H, s), 2.23 (2H, t, $J = 6.9$), 1.03-1.04 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 138.3, 129.5, 129.2, 126.6, 126.5, 62.5, 59.1, 40.2, 38.6, 18.4, 12.0. IR (thin film NaCl) 3446, 3030, 2942, 1210, 1035, 850 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{21}\text{H}_{36}\text{O}_2\text{SiNa}$, 371.2382; found, 371.2381.

(*E*)-3-ethyl-5-(triisopropylsilyloxy)pent-2-en-1-ol (2e). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and iodoethane (0.12 mL, 1.50 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. ^1H NMR analysis of the crude reaction mixture revealed a mixture of desired product **2e** and proton quench (**SI2**, see scheme above). Careful purification by flash chromatography (silica; 5, 8% EtOAc/hex) provided the title compound (31% and

14% yield from $\text{CuI} \cdot \text{P}(\text{OEt})_3$ and CuCl , respectively). R_f 0.25 (20% EtOAc/hex), faint UV, blue/purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.89 (1H, t, $J = 5.9$), 4.18 (2H, t, $J = 5.9$), 3.81 (2H, m), 2.76 (2H, m), 1.82 (2H, q, $J = 7.9$), 1.05-1.12 (22H, m), 0.94 (3H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 137.8, 125.8, 62.3, 59.4, 40.0, 25.0, 18.3, 12.1, 11.9. IR (thin film NaCl) 3442, 3061, 2932, 1215, 1052 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{34}\text{O}_2\text{SiNa}$, 309.2226; found, 309.2223.

(*E*)-5-(triisopropylsilyloxy)pent-2-en-1-ol (SI-2). ^1H NMR (500 MHz, CDCl_3) δ 5.73 (2H, m), 4.16 (2H, d, $J = 7.0$), 3.73 (2H, m), 2.31 (2H, m), 1.07 (21H, m).

(*E*)-3-Methyl-5-(triisopropylsilyloxy)pent-2-en-1-ol (2f). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and iodomethane (94 μL , 1.50 mmol, 300 mol %), using standard $\text{CuI} \cdot \text{P}(\text{OEt})_3$ or CuCl procedure. ^1H NMR analysis of the crude reaction mixture revealed a mixture of desired product **2f** and proton quench (**SI2**, see scheme above). Careful purification by flash chromatography (silica; 5, 8% EtOAc/hex) provided the title compound (44% and 31% yield from $\text{CuI} \cdot \text{P}(\text{OEt})_3$ and CuCl , respectively). R_f 0.25 (20% EtOAc/hex), faint UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.93 (1H, t, $J = 5.9$), 4.19 (2H, t, $J = 5.9$), 3.82 (2H, t, $J = 6.4$), 2.74 (2H, t, $J = 6.4$), 1.59 (3H, s), 1.05 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 135.9, 127.1, 61.9, 58.9, 42.6, 18.9, 18.4, 12.3. IR (thin film NaCl) 3440, 3060, 2935, 1213, 1051 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{15}\text{H}_{32}\text{O}_2\text{SiNa}$, 295.2069; found, 295.2069.

(3-Iodoprop-1-ynyl)trimethylsilane (SI3).

A solution of propargyl alcohol (1.00 mL, 17.2 mmol) in THF (43.0 mL, 0.400 M) was cooled to $-78\text{ }^{\circ}\text{C}$, and *n*-BuLi (2.4 M in hexanes, 21.5 mL, 51.5 mmol, 300 mol %) was added dropwise. After stirring for 20 min at $-78\text{ }^{\circ}\text{C}$, TMS-Cl (7.70 mL, 60.1 mmol, 350 mol %) was added dropwise. The reaction was stirred for 5 min at $-78\text{ }^{\circ}\text{C}$, the cold bath was removed, and the reaction stirred 1 h. The reaction was quenched with 50 mL water and then 10% aq. HCl was added to the crude reaction mixture until complete consumption of the TMS-ether was observed (TLC analysis, R_f 0.90 (10% EtOAc/hex), no UV, purple (vanillin)). The reaction was poured into a separatory funnel containing 100 mL Et_2O , shaken, and the layers were separated. The aqueous layer was extracted 2 $\times$ into 100 mL Et_2O . The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and carefully concentrated *in vacuo* (10 torr, $0\text{ }^{\circ}\text{C}$). Purification by flash chromatography (silica; 5, 20% Et_2O /pent) provided the alcohol (**4**) (2.15 g, 97.6%) as a pale yellow oil. R_f 0.25 (10% EtOAc/hex), no UV, purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 4.28 (2H, s), 1.62 (1H, bs), 0.19 (9H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 104.2, 90.5, 51.5, -0.1 . IR (thin film NaCl) 3430, 2200, 1036 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_6\text{H}_{12}\text{OSiNa}$, 151.0555; found, 151.0553.

A solution of PPh_3 (313 mg, 1.20 mmol, 120 mol %) in 2.50 mL CH_2Cl_2 was cooled to $0\text{ }^{\circ}\text{C}$. To the resulting solution were added sequentially: imidazole (102 mg, 1.50 mmol, 150 mol %), iodine (342 mg, 1.35 mmol, 135 mol %), and a solution of alcohol **4** (128 mg, 1.00 mmol) in CH_2Cl_2 (2.50 mL, 0.200 M total). The reaction was stirred at $0\text{ }^{\circ}\text{C}$, protected from light, for 90 min. The reaction was quenched by addition of 35 mL 20% aq. $\text{Na}_2\text{S}_2\text{O}_3$ and 35 mL CH_2Cl_2 . The resulting layers were separated. The aqueous layer was extracted 2 $\times$ into 35 mL CH_2Cl_2 . The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and carefully concentrated *in vacuo* (10 torr, $0\text{ }^{\circ}\text{C}$). Purification by flash chromatography (silica; pent) provided the target iodide **SI3** (150 mg, 63.2%) as a pale yellow oil. R_f 0.90 (5% EtOAc/hex), UV, faint brown (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 3.71 (2H, s), 0.17 (9H, s). The propargyl iodide

proved somewhat unstable, and was utilized immediately in subsequent reactions.

(*E*)-3-(2-(Triisopropylsilyloxy)ethyl)-6-(trimethylsilyl)hex-2-en-5-yn-1-ol (2g).

Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and propargyl iodide **SI3** (357 mg, 1.50 mmol, 300 mol %), using standard CuI•P(OEt)₃ or CuCl procedure. Purification by flash chromatography (silica; 4, 6% EtOAc/hex) provided the title compound (37% and 22% yield, respectively). *R*_f 0.30 (20% EtOAc/hex), faint UV, blue (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 5.53 (1H, t, *J* = 6.8), 4.18 (2H, d, *J* = 6.8), 3.75 (2H, t, *J* = 7.2), 2.72 (2H, s), 2.32 (2H, t, *J* = 7.2), 1.41 (1H, bs), 1.03-1.05 (21H, m), 0.20 (9H, s). ¹³C NMR (125 MHz, CDCl₃) δ 138.6, 123.0, 101.2, 90.5, 63.1, 59.4, 40.3, 23.1, 18.0, 12.1, −0.1. IR (thin film NaCl) 3436, 3065, 2930, 2225, 1220, 1060 cm^{−1}. HRMS ESI (*m/z*): [M+Na]⁺ calcd for C₂₀H₄₀O₂SiNa, 391.2465; found, 391.2450.

(3-Iodoprop-1-ynyl)dimethyl(vinyl)silane (SI4).

A solution of propargyl alcohol (0.50 mL, 8.6 mmol) in THF (22 mL, 0.40 M) was cooled to −78 °C and *n*-BuLi (2.4 M in hexanes, 10.7 mL, 25.8 mmol, 300 mol %) was added dropwise. After stirring for 20 min at −78 °C, vinyltrimethylchlorosilane (4.10 mL, 30.1 mmol, 350 mol %) was added dropwise. The reaction was stirred for 30 min at −78 °C, the cold bath was removed, and the reaction stirred 5 h. The reaction was quenched with 50 mL water and then 10% aq. HCl was added to the crude reaction mixture until complete consumption of the silyl ether was observed (TLC analysis, *R*_f 0.90 (10% EtOAc/hex), faint UV, blue (vanillin)). The reaction was poured into a separatory

funnel containing 50 mL Et₂O, shaken, and the layers were separated. The aqueous layer was extracted 2× into 50 mL Et₂O. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and carefully concentrated *in vacuo* (10 torr, 0 °C). Purification by flash chromatography (silica; 5, 10, 20% Et₂O/pent) provided the alkynyl silane (763 mg, 63.3%) as a pale yellow oil. R_f 0.25 (10% EtOAc/hex), faint UV, light purple (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 6.14 (1H, dd, *J* = 14.0, 19.5), 6.03 (1H, dd, *J* = 4.0, 14.0), 5.85 (1H, dd, *J* = 4.0, 19.5), 4.30 (2H, s), 1.62 (1H, bs), 0.26 (6H, s). ¹³C NMR (125 MHz, CDCl₃) δ 136.3, 133.5, 105.2, 88.7, 51.6, −1.6.

A solution of PPh₃ (224 mg, 0.713 mmol, 120 mol %) in 1.78 mL CH₂Cl₂ was cooled to 0 °C. To the resulting solution were added sequentially: imidazole (72 mg, 1.07 mmol, 150 mol %), iodine (244 mg, 0.963 mmol, 135 mol %), and alkynyl silane (100 mg, 0.713 mmol) in CH₂Cl₂ (1.78 mL, 0.200 M total). The reaction was stirred at 0 °C, protected from light, for 45 min. The reaction was quenched by addition of 35 mL 20% aq. Na₂S₂O₃ and 35 mL CH₂Cl₂. The resulting layers were separated. The aqueous layer was extracted 2× into 35 mL CH₂Cl₂. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and carefully concentrated (10 torr, 0 °C). Purification by flash chromatography (silica; pent) provided the target iodide **SI4** (142 mg, 79.4%) as a pale yellow oil. R_f 0.90 (5% EtOAc/hex), UV, faint brown (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 6.13 (1H, dd, *J* = 14.5, 20.0), 6.04 (1H, dd, *J* = 4.0, 14.5), 5.86 (1H, dd, *J* = 4.0, 20.0), 3.73 (2H, s), 0.24 (6H, s). The propargyl iodide proved somewhat unstable, and was utilized immediately in subsequent reactions.

(*E*)-6-(Dimethyl(vinyl)silyl)-3-(2-(triisopropylsilyloxy)ethyl)hex-2-en-5-yn-1-ol (2h). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and propargyl iodide **SI4** (375 mg, 1.50 mmol, 300 mol %), using standard $\text{CuI} \cdot \text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 4, 6% EtOAc/hex) provided the title compound (62% and 47% yield, respectively). R_f 0.30 (20% EtOAc/hex), faint UV, purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 6.09 (1H, dd, $J = 13.8, 19.5$), 5.99 (1H, dd, $J = 4.0, 13.8$), 5.85 (1H, dd, $J = 4.0, 19.5$), 5.55 (1H, t, $J = 6.9$), 4.20 (2H, d, $J = 6.9$), 3.75 (2H, t, $J = 7.2$), 2.71 (2H, s), 2.33 (2H, t, $J = 7.2$), 1.04-1.06 (21H, m), 0.25 (6H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 138.5, 133.3, 123.1, 105.1, 88.5, 62.9, 59.4, 40.1, 23.3, 17.9, 12.1, -1.7, -1.7. IR (thin film NaCl) 3446, 3060, 2927, 2228, 1660, 1220, 1062 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{21}\text{H}_{40}\text{O}_2\text{Si}_2\text{Na}$, 403.2465; found, 403.2463.

(*Z*)-2-(2-(Triisopropylsilyloxy)ethyl)but-2-ene-1,4-diol (2i). Reaction was conducted with alkynol **1** (128 mg, 0.500 mmol) and *para*-formaldehyde (45.0 mg, 1.50 mmol, 300 mol %), using standard $\text{CuI} \cdot \text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 2, 5% EtOAc/hex) provided the title compound (55% and 71% yield, respectively). R_f 0.10 (30% EtOAc/hex), no UV, purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.63 (1H, t, $J = 6.9$), 4.18 (2H, d, $J = 6.9$), 4.15 (2H, s), 3.84 (2H, t, $J = 5.7$), 2.39 (2H, t, $J = 5.7$), 1.06-1.07 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 141.6, 128.8, 64.6,

60.6, 58.6, 39.7, 18.1, 12.0. IR (thin film NaCl) 3520, 3060, 1220, 1060 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{15}\text{H}_{32}\text{O}_3\text{SiNa}$, 311.2018; found, 311.2016.

(*E*)-2-Allylbut-2-ene-1,4-diol (3b). Reaction was conducted with 2-butyn-1,4-diol (86.1 mg, 1.00 mmol) and allyl bromide (0.26 mL, 3.00 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 30, 100% EtOAc/hex) provided the title compound (82% and 89% yield, respectively). R_f 0.30 (50% EtOAc/hex), no UV, purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.75-5.78 (2H, m), 5.06 (1H, dd, $J = 1.5, 18.0$), 5.03 (1H, dd, $J = 1.5, 10.5$), 4.22 (2H, d, $J = 7.0$), 4.06 (2H, s), 2.87 (2H, d, $J = 6.4$), 2.17 (2H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 140.0, 135.6, 125.4, 116.1, 66.2, 59.0, 32.7. IR (thin film NaCl) 3506, 3082, 2927, 1630 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_7\text{H}_{12}\text{O}_2\text{Na}$, 151.0735; found, 151.0739.

(*E*)-3-Allyl-2,5-dimethylhex-3-ene-2,5-diol (3c). Reaction was conducted with 2,5-dimethyl-3-hexyne-2,5-diol (39 mg, 0.273 mmol) and allyl bromide (71 μL , 0.819 mmol, 300 mol %), using standard $\text{CuI}\cdot\text{P}(\text{OEt})_3$ or CuCl procedure. Purification by flash chromatography (silica; 20, 30% EtOAc/hex) provided the title compound (82% and 91% yield, respectively). R_f 0.50 (80% EtOAc/hex), no UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 6.05 (1H, m), 5.75 (1H, m), 5.11 (1H, m), 5.08 (1H, m), 3.26 (2H, m), 1.70 (2H, m), 1.36 (3H, s), 1.35 (3H, s), 1.34 (3H, s), 1.34 (3H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 142.7, 140.6, 131.9, 115.2,

74.5, 71.9, 31.6, 31.6, 30.1.^{S7} IR (thin film NaCl) 3078, 2935, 1630, 1385, 1366, 1180 cm⁻¹. HRMS ESI (*m/z*): [M+Na]⁺ calcd for C₁₁H₂₀O₂Na, 207.1361; found, 207.1357.

(Z)-1-(2-Methylpenta-1,4-dienyl)cyclohexanol (3d). Reaction was conducted with 1-(1-propynyl)cyclohexanol (38 mg, 0.273 mmol) and allyl bromide (71 μ L, 0.819 mmol, 300 mol %), using standard CuI•P(OEt)₃ or CuCl procedure. Purification by flash chromatography (silica; 3% EtOAc/hex) provided the title compound (96% and 95% yield, respectively). *R*_f 0.40 (10% EtOAc/hex), no UV, purple (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 5.84 (1H, ddt, *J* = 6.3, 10.1, 17.1), 5.39 (1H, s), 5.07 (1H, dd, *J* = 1.8, 17.1), 5.03 (1H, dd, *J* = 1.8, 10.1), 3.14 (2H, d, *J* = 6.3), 1.72 (3H, s), 1.62 (4H, m), 1.45 (4H, m), 1.35 (2H, m). ¹³C NMR (125 MHz, CDCl₃) δ 137.2, 136.3, 133.0, 115.4, 72.1, 39.9, 37.3, 25.6, 24.8, 22.7. IR (thin film NaCl) 3040, 2920, 1625, 1170 cm⁻¹. HRMS ESI (*m/z*): [M+Na]⁺ calcd for C₁₂H₂₀ONa, 203.1412; found, 203.1414.

(E)-2-Methyl-5-phenylocta-4,7-dien-3-ol (3e). Reaction was conducted with 1-phenyl-4-methyl-1-pentyn-3-ol (48 μ L, 0.273 mmol) and allyl bromide (71 μ L, 0.819 mmol, 300 mol %), using standard CuI•P(OEt)₃ or CuCl procedure. Purification by flash chromatography (silica; 5, 7% EtOAc/hex) provided the title compound (81% and 89% yield, respectively). *R*_f 0.40 (20% EtOAc/hex), UV active, blue (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 7.40-7.42 (2H, m), 7.31-7.35 (3H, m), 5.82-5.90 (2H, m), 5.10 (1H, ddt, *J* = 1.7, 3.6, 17.1), 5.03 (1H, ddt,

(S7) In compound **3c**, two ¹³C resonances, corresponding to two sets of equivalent methyl groups, appear as a peak with a large shoulder, at 31.611 and 31.593 ppm, respectively.

$J = 1.7, 3.4, 10.2$), 4.22 (1H, dd, $J = 7.8, 7.9$), 3.35 (2H, m), 1.81 (1H, dtt, $J = 7.8, 6.7, 6.9$), 1.48 (1H, bs), 1.04 (3H, d, $J = 6.7$), 0.96 (3H, d, $J = 6.9$). ^{13}C NMR (125 MHz, CDCl_3) δ 142.4, 140.5, 136.5, 131.0, 128.5, 127.5, 126.6, 116.1, 73.9, 34.9, 34.7, 18.7, 18.5. IR (thin film NaCl) 3360, 3040, 2940, 1380, 1362, 870 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{15}\text{H}_{20}\text{ONa}$, 239.1412; found, 239.1412.

3-(Trimethylsilyl)prop-2-yn-1-ol (4).^{S8} A solution of propargyl alcohol (1.00 mL, 17.2 mmol) in THF (43.0 mL, 0.400 M) was cooled to $-78\text{ }^{\circ}\text{C}$, and $n\text{-BuLi}$ (2.4 M in hexanes, 21.5 mL, 51.5 mmol, 300 mol %) was added dropwise. After stirring for 20 min at $-78\text{ }^{\circ}\text{C}$, TMS-Cl (7.70 mL, 60.1 mmol, 350 mol %) was added dropwise. The reaction was stirred for 5 min at $-78\text{ }^{\circ}\text{C}$, the cold bath was removed, and the reaction stirred 1 h. The reaction was quenched with 50 mL water and then 10% HCl was added to the crude reaction mixture until complete consumption of the TMS-ether was observed (TLC analysis, R_f 0.90 (10% EtOAc/hex), no UV, purple (vanillin)). The reaction was poured into a separatory funnel containing 100 mL Et_2O , shaken, and the layers were separated. The aqueous layer was extracted 2 $\times$ into 100 mL Et_2O . The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and carefully concentrated *in vacuo* (10 torr, $0\text{ }^{\circ}\text{C}$). Purification by flash chromatography (silica; 5, 20% Et_2O /pent) provided the TMS-propargyl alcohol (2.15 g, 97.6%) as a pale yellow oil. R_f 0.25 (10% EtOAc/hex), no UV, purple (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 4.28 (2H, s), 1.62 (1H, bs), 0.19 (9H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 104.2, 90.5, 51.5, -0.1 . IR (thin film NaCl) 3430, 2200, 1036 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_6\text{H}_{12}\text{OSiNa}$, 151.0555; found, 151.0553.

(S8) Procedure amended from: (a) Velcicky, J.; Lanver, A.; Lex, J.; Prokop, A.; Wieder, T.; Schmalz, H.-G. *Chem. Eur. J.* **2004**, *10*, 5087-5110. (b) Jones, T. K.; Denmark, S. E. *Org. Synth.* **1986**, *64*, 182-188.

(*E*)-3-(trimethylsilyl)hexa-2,5-dien-1-ol (5). Reaction was conducted with alcohol **4** (45 mg, 0.273 mmol) and allyl bromide (71 μ L, 0.819 mmol, 300 mol %), using standard $\text{CuI} \cdot \text{P}(\text{OEt})_3$ procedure. Purification by flash chromatography (silica; 5, 10% EtOAc/hex) provided the title compound (44.3 mg, 95%). R_f 0.50 (20% EtOAc/hex), no UV, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 6.02 (1H, t, $J = 6.0$), 5.75 (1H, m), 5.00 (1H, m), 4.97 (1H, m), 4.25 (2H, d, $J = 6.0$), 2.93 (2H, d, $J = 6.1$), 1.55 (1H, bs), 0.09 (9H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 141.2, 140.0, 137.0, 115.0, 59.9, 34.0, -1.4 . IR (thin film NaCl) 3462, 3075, 2920, 1646 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_9\text{H}_{18}\text{OSiNa}$, 193.1025; found, 193.1027.

(*Z*)-3-(Trimethylsilyl)hexa-2,5-dien-1-ol (5'). Following Ziegler's procedure,^{S9} a round-bottom flask was charged with alcohol **4** (35 mg, 0.273 mmol) and Et_2O (0.55 mL, 0.50 M), and the resulting solution was cooled to -20°C . Neat $i\text{-Bu}_2\text{AlH}$ (0.14 mL, 0.955 mmol, 350 mol %) was added dropwise, and after 2 min, the cold bath was removed. The reaction was heated to gentle reflux (40°C). After 12 h, the reaction was cooled to 0°C , then MeLi (1.6 M in Et_2O , 0.64 mL, 1.02 mmol, 375 mol %) was added dropwise. The cold bath was removed and, after stirring for 90 min, the reaction was cooled to -78°C ($\text{CO}_2/\text{acetone}$). A solution of $\text{CuI} \cdot \text{P}(\text{OEt})_3$ (86 mg, 100 mol %) in THF (1.4 mL, 0.20 M) was added by cannula (exothermic!), followed by addition of neat allyl bromide (71 μ L, 0.819 mmol, 300 mol %). The reaction was stirred for 24 h, allowing the cold bath to warm slowly to rt. The crude reaction mixture was

(S9) Ziegler, F. E.; Mikami, K. *Tetrahedron Lett.* **1984**, 25, 131-134.

poured onto sat. aq. potassium sodium tartrate (30 mL), rinsed with 30 mL EtOAc, and the mixture was stirred for 30 min. The layers were separated, and the aqueous layer was extracted 2× into 30 mL EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification by flash chromatography (silica; 5, 10% EtOAc/hex) provided the title compound (20.0 mg, 43%). *R*_f 0.50 (20% EtOAc/hex), no UV, blue (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 6.15 (1H, t, *J* = 6.9), 5.81 (1H, m), 5.03 (1H, m), 5.01 (1H, m), 4.24 (2H, d, *J* = 7.1), 2.87 (2H, d, *J* = 6.4), 1.54 (1H, bs), 0.17 (9H, s). ¹³C NMR (125 MHz, CDCl₃) δ 140.8, 138.6, 137.8, 115.8, 59.2, 40.6, −1.2. IR (thin film NaCl) 3470, 3090, 2942, 1660 cm^{−1}. HRMS ESI (*m/z*): [M+Na]⁺ calcd for C₉H₁₈OSiNa, 193.1025; found, 193.1024.

Hepta-2,5-diyne-1-ol (8).^{S10} To a rt solution of EtMgCl (2.0 M in THF, 18.0 mL, 36.0 mmol, 284 mol %) was added by cannula, portionwise over 25 min, a solution of propargyl alcohol (1.01 mL, 17.1 mmol, 135 mol %) in THF (2.20 mL, 5.8 M). Once the addition was complete, the reaction was heated to reflux for 90 min. The flask was cooled to 0 °C, then solid CuCl (126 mg, 1.27 mmol, 10.0 mol %) was added. After stirring for 15 min at 0 °C, 1-bromo-2-butyne (1.14 mL, 12.7 mmol) was added dropwise over 10 min. Upon complete addition, the reaction was heated to reflux for 16 h. The reaction flask was cooled to rt, then quenched by careful addition, over 10 min, of 87 mL 10% aq. NH₄Cl (exothermic!). The mixture was added to a separatory funnel containing 80 mL Et₂O, and the layers were separated. The aqueous layer was extracted 2× into 80 mL Et₂O. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and carefully concentrated *in vacuo* (10 torr, 0 °C). Distillation

(S10) Bushby, R. J.; Whitham, G. H. *J. Chem. Soc. B* **1970**, 563-569.

(5 torr, 150 °C) provided the title compound (997 mg, 73%) as a colorless oil.^{S11} ¹H NMR (500 MHz, CDCl₃) δ 4.28 (2H, m), 3.17 (2H, q, *J* = 2.5), 1.81 (3H, t, *J* = 2.5), 1.59 (1H, bs).

(*E*)-Hept-2-en-5-yn-1-ol (SI5).^{S12} To a 0 °C solution of diynol **8** (250 mg, 2.31 mmol, 105 mol %) in Et₂O (4.62 mL, 0.50 M) was added, slowly over 10 min, solid LiAlH₄ (84.0 mg, 2.20 mmol). Upon complete addition, the reaction was heated to a gentle reflux for 7 h. To the crude mixture were carefully added in sequence: 3.0 mL Et₂O, 84 μL H₂O, 84 μL of 2 N aq. NaOH, and then 0.24 mL H₂O. A small amount of MgSO₄ was added to the flask, and the resulting mixture was stirred at rt for 5 min. After filtration through Celite, the residue was carefully concentrated *in vacuo* (10 torr, 0 °C) to afford the title compound (258 mg, 106% theoretical yield) as a colorless oil. The crude product was used without further purification. *R*_f 0.10 (5% EtOAc/hex), no UV, blue-green (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 5.92 (1H, m), 5.71 (1H, m), 4.15 (2H, dd, *J* = 1.0, 5.7), 2.93 (2H, m), 1.83 (3H, t, *J* = 2.5), 1.46 (1H, bs).

(*E*)-1-Iodohept-2-en-5-yne (9). To a 0 °C solution of PPh₃ (131 mg, 0.499 mmol, 110 mol %) in 1.30 mL CH₂Cl₂ were added sequentially: imidazole (43.0 mg, 0.636 mmol, 140 mol %), iodine (138 mg, 0.545 mmol, 120 mol %), and a solution of crude alcohol **SI5** (50.0 mg, 0.454 mmol) in CH₂Cl₂ (1.00 mL, 0.200 M total). The reaction was stirred at 0 °C, protected from light, for 30 min. The reaction was quenched by addition of 40 mL of 20% aq. Na₂S₂O₃. The aqueous

(S11) Upon storage (2 days, 4 °C), a yellow color begins to evolve. Optimum yield in subsequent steps was obtained when diynol **8** was colorless (i.e. freshly distilled, or frozen at –10 °C).

(S12) Aebi, J. D.; Deyo, D. T.; Sun, G. C.; Guillaume, D.; Dunlap, B.; Rich, D. H. *J. Med. Chem.* **1990**, *33*, 999-1009.

layer was extracted 3× into 50 mL CH₂Cl₂. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and carefully concentrated *in vacuo* (10 torr, 0 °C). Purification by flash chromatography (silica; 0, 2% Et₂O/pent) provided the title compound (93 mg, 93%) as a faint pink oil. R_f 0.60 (5% EtOAc/hex), UV active, green (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 6.00 (1H, dt, *J* = 8.0, 15.0), 5.70 (1H, dt, *J* = 5.4, 15.0), 3.90 (2H, d, *J* = 8.0), 2.91 (2H, dq, *J* = 2.5, 5.4), 1.83 (3H, t, *J* = 2.5). The iodide proved somewhat unstable, and was utilized immediately in subsequent reactions.

(2*E*,5*E*)-3-(2-(Triisopropylsilyloxy)ethyl)deca-2,5-dien-8-yn-1-ol (10) and **(*Z*)-3-(2-(Triisopropylsilyloxy)ethyl)-4-vinyloct-2-en-6-yn-1-ol (10')**.

Reaction was conducted with alcohol **1** (100 mg, 0.390 mmol) and allyl iodide **9** (257 mg, 1.17 mmol, 300 mol %), using standard CuCl procedure. Purification by flash chromatography (silica; 6% EtOAc/hex) initially provided the S_N2' product **10'** (62.9 mg, 46.0%) as a colorless oil. Further elution from the column (silica; 6, 8% EtOAc/hex) provided the S_N2 product **10** as a pale yellow oil (69.7 mg, 51.0%).

• **Compound 10.** R_f 0.30 (20% EtOAc/hex), no UV, green (vanillin). ¹H NMR (500 MHz, CDCl₃) δ 5.62 (1H, m), 5.52 (1H, t, *J* = 6.9), 5.45 (1H, m), 4.17 (2H, d, *J* = 6.9), 3.77 (2H, t, *J* = 7.1), 2.86 (2H, m), 2.84 (2H, d, *J* = 7.2), 2.29 (2H, t, *J* = 7.1), 1.81 (3H, t, *J* = 2.5). ¹³C NMR (125 MHz, CDCl₃) δ 139.2, 131.0, 129.0, 126.4, 80.3, 76.5, 62.7, 59.4, 40.5, 34.0, 22.1, 18.2, 12.2, 3.7. IR (thin film NaCl) 3465, 3040, 2926, 2270, 1240, 1060 cm⁻¹. HRMS ESI (*m/z*): [M+Na]⁺ calcd for C₂₁H₃₈O₂SiNa, 373.2539; found, 373.2537.

• **Compound 10'**. R_f 0.40 (20% EtOAc/hex), no UV, brown (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 5.83 (1H, ddd, $J = 6.3, 10.5, 17.2$), 5.66 (1H, t, $J = 7.2$), 5.09 (1H, ddd, $J = 1.3, 1.4, 10.5$), 5.07 (1H, ddd, $J = 1.4, 1.5, 17.2$), 4.27 (1H, dd, $J = 7.6, 12.3$), 4.12 (1H, dd, $J = 6.9, 12.3$), 3.78 (2H, t, $J = 7.2$), 3.45 (1H, m), 2.36 (2H, m), 2.24 (2H, t, $J = 7.6$), 1.77 (3H, t, $J = 2.5$), 1.44 (1H, bs), 1.06-1.07 (21H, m). ^{13}C NMR (125 MHz, CDCl_3) δ 140.7, 138.9, 127.3, 115.6, 77.9, 77.4, 63.5, 58.6, 44.2, 35.3, 22.5, 18.2, 12.2, 3.6. IR (thin film NaCl) 3480, 3026, 2909, 2265, 1648, 1025 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{21}\text{H}_{38}\text{O}_2\text{SiNa}$, 373.2539; found, 373.2547.

(Z)-3-(Triethylsilyl)hept-2-en-5-yn-1-ol (11). To a 0 °C solution of distilled diyne **8** (50 mg, 0.46 mmol) in CH_2Cl_2 (0.93 mL, 0.50 M) was added Et_3SiH (92 μL , 0.575 mmol, 123 mol %). Solid $[\text{Cp}^*\text{Ru}(\text{MeCN})_3]\text{PF}_6^{\text{S13,S14}}$ (3.5 mg, 6.94 μmol , 1.5 mol %) was added, and the reaction was stirred for 20 min. A second portion of Ru catalyst (3.5 mg) was added, and the reaction was stirred an additional 30 min. The reaction mixture was filtered through a $\frac{1}{2}$ in. pad of SiO_2 , rinsed with CH_2Cl_2 , and concentrated *in vacuo*. Purification by flash chromatography (silica; 5, 10% EtOAc/hex) provided the title compound (45 mg, 43%). R_f 0.30 (20% EtOAc/hex), no UV, green (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 6.57 (1H, t, $J = 7.0$), 4.23 (2H, d, $J = 7.0$), 2.94 (2H, q, $J = 2.5$), 1.84 (3H, t, $J = 2.5$), 1.37 (1H, bs), 0.95 (9H, t, $J = 8.1$), 0.70 (6H, q, $J = 8.1$). ^{13}C NMR (125 MHz, CDCl_3) δ 141.9, 136.0, 79.9, 73.1, 62.3, 27.7, 7.7, 4.2, 3.8. IR (thin film NaCl) 3465, 3091, 2946, 2960 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{13}\text{H}_{24}\text{OSiNa}$, 247.1494; found, 247.1497.

(S13) Tris(acetonitrile)pentamethylcyclopentadienylruthenium (I) hexafluorophosphate was purchased from Strem Chemicals and used without further purification.

(S14) Trost, B. M.; Ball, Z. T. *J. Am. Chem. Soc.* **2005**, *127*, 17644-17655.

(Z)-Triethyl(1-iodohept-2-en-5-yn-3-yl)silane (12). To a 0 °C solution of PPh_3 (59 mg, 0.226 mmol, 110 mol %) in 1.10 mL CH_2Cl_2 were added sequentially: imidazole (20 mg, 0.287 mmol, 140 mol %), iodine (62 mg, 0.246 mmol, 120 mol %), and a solution of alcohol **11** (46 mg, 0.205 mmol) in CH_2Cl_2 (1.00 mL, 0.200 M total). The reaction was stirred at 0 °C, protected from light, for 60 min. The reaction was quenched by addition of 40 mL of 20% aq. $\text{Na}_2\text{S}_2\text{O}_3$. The aqueous layer was extracted 3× into 50 mL CH_2Cl_2 . The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and carefully concentrated *in vacuo* (10 torr, 0 °C). Purification by flash chromatography (silica; 2, 4% Et_2O /pent) provided the title compound (91.5 mg, 93%) as a faint pink oil. R_f 0.85 (10% EtOAc /hex), faint active, blue-green (vanillin). The iodide proved somewhat unstable, and was utilized immediately in subsequent reactions.

(2E,5Z)-6-(Triethylsilyl)-3-(2-(triisopropylsilyloxy)ethyl)deca-2,5-dien-8-yn-1-ol (13). Reaction was conducted with alcohol **1** (28 mg, 0.109 mmol) and allyl iodide **12** (109 mg, 0.327 mmol, 300 mol %), using standard CuCl procedure. Purification by flash chromatography (silica; 6% EtOAc /hex) provided the title compound (36.0 mg, 70.9%) as a colorless oil. R_f 0.50 (20% EtOAc /hex), faint active, blue (vanillin). ^1H NMR (500 MHz, CDCl_3) δ 6.17 (1H, t, $J = 6.5$), 5.51 (1H, t, $J = 7.0$), 4.18 (2H, d, $J = 7.0$), 3.78 (2H, t, $J = 7.0$), 2.96 (2H, d, $J = 6.5$), 2.88 (2H, q, $J = 2.0$), 2.30 (2H, t, $J = 7.0$), 1.81 (3H, t, $J = 2.0$), 1.41 (1H, bs), 1.05-1.06 (21H, m), 0.96 (9H, t, $J = 8.0$), 0.73 (6H, q, $J = 8.0$). ^{13}C NMR (125 MHz, CDCl_3) δ 141.9, 139.9, 133.5, 126.2, 79.1, 77.7, 62.6, 59.4, 40.5, 33.5, 27.9, 18.2, 12.2, 7.7, 4.0, 3.8. IR (thin film NaCl) 3480, 3065, 2940, 2250, 1240,

1059 cm^{-1} . HRMS ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{27}\text{H}_{52}\text{O}_2\text{Si}_2\text{Na}$, 487.3404; found, 487.2404.