

Supplemental Table 1. Proportions of cells categorized as CD24-negative (Preadipocytes) or CD24-positive (Progenitors) at each level of FACS analysis.

Group	% of Live Cells		% of CD29+ Cells		% of Sca1+/CD34+ Cells	
	Preadipocytes	Progenitors	Preadipocytes	Progenitors	Preadipocytes	Progenitors
LF E2	17.5 +/- 1.3	1.0 +/- 0.2	31.8 +/- 7.5	1.7 +/- 0.3	63.2 +/- 2.7	3.7 +/- 0.7
LF E2+TAM	15.9 +/- 2.9	1.2 +/- 0.4	38.1 +/- 7.4	2.3 +/- 0.5	79.1 +/- 3.8	5.1 +/- 1.0
LF EWD	14.8 +/- 1.6	1.3 +/- 0.2	30.5 +/- 9.4	2.5 +/- 0.4	76.4 +/- 3.1	6.9 +/- 0.8
HF E2	15.3 +/- 1.2	1.3 +/- 0.2	26.2 +/- 4.2	2.2 +/- 0.4	67.0 +/- 3.0	5.9 +/- 1.1
HF E2+TAM	14.4 +/- 1.5	0.9 +/- 0.4	29.4 +/- 4.1	2.3 +/- 1.4	75.0 +/- 5.3	2.7 +/- 0.5
HF EWD	21.8 +/- 4.9	1.0 +/- 0.2	38.6 +/- 8.7	1.9 +/- 0.4	76.4 +/- 5.2	4.4 +/- 1.2