

## References to Table S2

- Amaral-Zettler, L.A., McCliment, E.A., Ducklow, H.W. & Huse, S.M. (2009): A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. - *PLoS One* 4:e6372.
- Bittner, L., Gobet, A., Audic, S., Romac, S., Egge, E.S., Santini, S., Ogata, H. et al. (2013): Diversity patterns of uncultured Haptophytes unravelled by pyrosequencing in Naples Bay. - *Mol. Ecol.* 22:87–101.
- Egge, E.S., Bittner, L., Andersen, T., Audic, S., de Vargas, C. & Edvardsen, B. (2013): 454 Pyrosequencing to describe microbial eukaryotic community composition, diversity and relative abundance: a test for marine haptophytes. - *PLoS One* 8:e74371.
- Frias-Lopez, J., Thompson, A., Waldbauer, J. & Chisholm, S.W. (2009): Use of stable isotope-labelled cells to identify active grazers of picocyanobacteria in ocean surface waters. - *Environ. Microbiol.* 11:512–25.
- Liu, H., Probert, I., Uitz, J., Claustre, H., Aris-Brosou, S., Frada, M., Not, F. et al. (2009): Extreme diversity in noncalcifying haptophytes explains a major pigment paradox in open oceans. - *Proc. Natl. Acad. Sci. U. S. A.* 106:12803–8.
- Medlin, L.K., Elwood, H.J., Stickel, S. & Sogin, M.L. (1988): The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding regions. - *Gene* 71:491–9.
- Moon-van der Staay, S.Y., van der Staay, G.W.M., Guillou, L., Vaultot, D., Claustre, H. & Medlin, L.K. (2000): Abundance and diversity of prymnesiophytes in the picoplankton community from the equatorial Pacific Ocean inferred from 18S rDNA sequences. - *Limnol. Oceanogr.* 45:98–109.
- Shalchian-Tabrizi, K., Reier-Røberg, K., Ree, D.K., Klaveness, D. & Bråte, J. (2011): Marine-freshwater colonizations of haptophytes inferred from phylogeny of environmental 18S rDNA sequences. - *J. Eukaryot. Microbiol.* 58:315–8.
- Simon, M., López-García, P., Moreira, D. & Jardillier, L. (2013): New haptophyte lineages and multiple independent colonizations of freshwater ecosystems. - *Environ. Microbiol. Rep.* 5:322–32.
- Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M.D.M., Breiner, H.W. & Richards, T.A. (2010): Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. - *Mol. Ecol.* 19:21–31.