

**S3 Table.** Prevalence of stunting of children participating in the MINIMat trial, Bangladesh, from birth to ten years.

Age	Girls		Boys		Total	
Months	n/N Stunted <sup>1</sup>	% Stunted <sup>1</sup>	n/N Stunted <sup>1</sup>	% Stunted <sup>1</sup>	n/N Stunted <sup>1</sup>	% Stunted <sup>1</sup>
0	81/513	15.8	89/541	16.5	170/1054	16.1
1	102/512	19.9	113/540	20.9	215/1052	20.4
2	94/513	18.3	133/541	24.6	227/1054	21.5
3	89/512	17.4	136/541	25.1	225/1053	21.4
4	85/513	16.6	133/541	24.6	218/1054	20.7
5	84/511	16.4	108/541	20.0	192/1052	18.3
6	96/513	18.7	152/541	28.1	248/1054	23.5
7	107/512	20.9	155/540	28.7	262/1052	24.9
8	112/512	21.9	179/541	33.1	291/1053	27.6
9	124/512	24.2	182/540	33.7	306/1052	29.1
10	135/512	26.4	191/539	35.4	326/1051	31.0
11	146/511	28.6	215/540	39.8	361/1051	34.3
12	155/509	30.5	215/540	39.8	370/1049	35.3
15	210/511	41.1	253/540	46.9	463/1051	44.1
18	236/512	46.1	284/541	52.5	520/1053	49.4
21	264/512	51.6	307/540	56.9	571/1052	54.3
24	260/513	50.7	295/541	54.5	555/1054	53.7
54	180/513	35.1	179/541	33.1	359/1054	34.1
120	166/513	32.4	142/541	26.2	308/1054	29.2

<sup>1</sup>Below minus 2 standard deviations from WHO growth reference median