READ ME - Kenya Long-term Exclosure Experiment

Principal Investigators:

Corinna Riginos, Department of Zoology and Physiology, University of Wyoming. <u>criginos@gmail.com</u>

Kari E. Veblen, Department of Wildland Resources and Ecology Center, Utah State University. kari.veblen@usu.edu

Truman P. Young, Department of Plant Sciences, University of California, Davis. tpyoung@ucdavis.edu

The KLEE experiment:

For a full site description, see Young et al. (1998). This manuscript is also included at the end of the "read me" document. KLEE was established in 1995 and consists of three replicate blocks, each containing six 200 x 200 m treatment plots. The replicate blocks are 70 - 200 m apart. The experiment uses semi-permeable barriers to allow access by different combinations of cattle ('C'), mesoherbivore wildlife 15 - 1000 kg ('W') and megaherbivores ('M'). Each of the following six treatments is replicated across the three blocks: C, W, WC, MW, MWC and O. The capital letters indicate which herbivores are allowed access (e.g., 'O' allows no herbivores >15 kg, 'W' allows mesoherbivore wildlife >15 kg, but no cattle or megaherbivores, and 'MWC' allows megaherbivores, mesoherbivore wildlife and cattle). Long-term patterns of dung deposition in the KLEE plots indicate that 1) treatments are >90% effective at excluding targeted species, and 2) experimental fences do not deter wild herbivores from using the plots intended to be accessible to them (see Young et al. 1998 for more details).

Herded groups of 100-120 head of cattle are grazed in C, WC and MWC plots for several hours on each of two to three consecutive days, typically 3-4 times per year. The precise number of grazing days and timing of grazing largely depends on forage availability, but plots rarely experience more than 16 weeks without cattle grazing. This grazing regime reflects typical cattle management strategies for both private and communal properties in the region – where livestock graze in one general area for several days at a time until forage is depleted, then move on to a different area until the forage recovers. The landscape is not fenced into paddocks, but rather herders guide livestock so that the entire range undergoes similar episodic grazing throughout the year. The stocking rate of plots is similar to the moderate overall ranch stocking rate (0.10-0.14 cattle/ha; Odadi et al. 2007). Fire has not been an active part of this ecosystem since the 1960s (R.L. Sensenig, personal communication).



Fig. 1: Layout of the Kenya Long-term Exclosure Experiment (KLEE). Each experimental plot measures 200 x 200m. Plots use semi-permeable fences to allow grazing by different combinations of large herbivores. Each capital letter represents the herbivore guild allowed in the plot: O (no large herbivores), C (cattle), W (mesoherbivore wildlife), and M (megaherbivores).

Site description:

The Kenya Long-term Exclusion Experiment (KLEE) is located at the Mpala Conservancy (36°52'E, 0°17'N) in Laikipia County, Kenya. The study area is underlain with 'black cotton' soils, poorly drained vertisols with high (>50%) clay content (Ahn and Geiger 1987). Black cotton savannas are widespread in East Africa, covering hundreds of thousands of km². Ninety-seven percent of the tree canopy cover in KLEE is *Acacia drepanolobium* Sjost. (Young et al. 1998), and total tree canopy cover averages 15-25%. Five grass species (*Pennisetum mezianum* Leeke, *P. stramineum* Peter, *Themeda triandra* Forssk., *Lintonia nutans* Stapf., and *Brachiaria lachnantha* (Hochst.) Stapf) make up 85% of herbaceous cover (Porensky et al. 2013). The site is located on virtually flat topography at an elevation of 1810m asl. The absence of distinct runoff or run-on areas, coupled with the relatively low plant diversity, makes this an ideal system to examine the effects of different herbivores on plant production independent of other factors.

Mpala Conservancy is managed for both wildlife conservation and livestock production, with mean stocking rates of 0.10-0.14 cattle/ha. Wild ungulates commonly found at the study site include the mesoherbivores plains zebra (*Equus burchelli* Gray), Grant's gazelle (*Gazella [Nanger] grantii* Brooke), hartebeest (*Alcelaphus buselaphus* Pallas), eland (*Taurotragus oryx* Pallas), and oryx (*Oryx gazella beisa* L.), as well as the megaherbivores elephant (*Loxodonta africana* Blumenbach), and reticulated giraffe (*Giraffa camelopardalis* L.). Total biomass-density (kg / km⁻²) of large wild ungulates is approximately one third of cattle biomass-density, and large wild ungulate biomass density is split almost evenly between megaherbivores and mesoherbivore wildlife (Veblen et al. 2016). In addition to these larger herbivores, one small antelope, steinbuck (*Raphicerus campestris* Thunberg), a strict browser, occurs in the area and is able to access all experimental treatment plots (Young et al. 2005). Wildlife in this region are present year-round and do not undergo large seasonal migrations.

Descriptions of datasets available for download

- 1. Rainfall
- 2. Dung
- 3. Acacia drepanolobium density and size
- 4. Plot locations
- 5. Vegetation dataset

Additional datasets available upon request

1. Rainfall

Methods and description: Rainfall measurements were recorded from manual rain gauges located along the fence lines of KLEE. There are a total of three rain gauges. Each KLEE block has one manual rain gauge. <u>Exact GPS locations are located in the "GPS locations" sheet of this document</u>. Rainfall dates are recorded and gauge measurements are taken as soon as possible after rainfall events.

This Excel worksheet has rainfall data from each of the three replicate KLEE blocks since 2003. There are separate worksheets for daily rainfall, monthly rainfall (since 1996), and yearly rainfall (since 1996). Additional monthly and yearly rainfall data since 1996 are available upon request. Measurements are taken in millimeters. Columns *North, Central*, and *South* refer to gauge readings taken in the north, central, and south blocks of KLEE.

2. Dung

Methods and description: Starting in 2006, we have conducted biennial (roughly March and October) dung surveys along six-4 m x100 m permanent transects within each of the 18 KLEE plots and three control transects outside of the KLEE plots. This Excel worksheet features dung count data for each KLEE plot. Count data is broken down by herbivore species. To avoid recounting the same dung piles during subsequent surveys, we crush all recorded dung piles during each session. For animals that defecate in middens (such as steinbuck and Grant's gazelle), we used the number of dung pellets and differences in shape and color to estimate the number of separate defecation events. Dung piles for all major herbivore species could be positively identified to species in the field, with two exceptions.

The dung of cattle and buffalo could not be distinguished, and we lumped them. The dung of plains and Grevy's zebras also could not be distinguished from each other; hence we grouped them as "zebra". However, plains zebra far outnumber Grevy's, so effectively we consider these to be plains zebra dung.

3. Acacia drepanolobium density and size survey

Methods and description: *A. drepanolobium* makes up approximately 97% of the woody cover in this system. This Excel file contains *Acacia drepanolobium* density and size distributions in each of the of the 18 KLEE plots. Surveys were conducted in 2011. Trees were surveyed along three 4 m x 100 m transects within each KLEE plot. Each row represents data from one transect. Columns indicate experimental block, plot, number of trees in each of five size classes, and total number of trees for each transect. Additional data for non-*drepanolobium* woody species, as well as a newly updated 2015 tree survey, are available upon request.

4. Plot locations

Methods and description: This zip file contains 18 .kml files. Each file corresponds with one KLEE plot and is named according to its block (first letter) and treatment (following letters). We collected GPS points from the corners of each of the 18 experimental plots using a Trimble Juno 3B GPS unit with meter-level accuracy. Points were imported into QGIS and used to created polygons of each experimental plot. Polygons were then saved as Google Earth .kml files.

5. Vegetation dataset

Methods and description: Herbaceous vegetation in all 18 KLEE plots has been sampled biannually (in February and June) or annually (in June) since 1995. Sampling periods follow rainy periods that are similar in terms of average rainfall. Due to improvements in species identification, pre-1999 surveys are not fully comparable to later data. Each of the eighteen 4 ha KLEE treatment plots contains a central hectare that is divided into a 10 x 10 m grid of 100 sampling stations. Aerial plant cover and composition are assessed at these stations by counting the number of pins hit by each species over a ten-point pin frame (with vertical pins separated by 5 cm; maximum one hit per pin per species). All 100 grid points were sampled 1999-2005, and every fifth grid point (20 per plot) was sampled from 2006 to 2013 (see Fig. 2 for current vegetation sampling layout).

This Excel datasheet contains vegetation data from 1999 - 2013. Later surveys have also been conducted and are available upon request. This Excel file is broken into the following sheets:

• "**Raw data**" sheet: Scaled sum first hits data for each year (columns) organized by species, then by treatment and block. This sheet has data for all species.

- "Lumped spp." sheet: Data are lumped together for several species that we decided needed to be treated together. (e.g. Melhania spp.)
 - Standardizing species list:
 - Aristida spp. \rightarrow Aristida kenyensis
 - Eragrostis tenuifolia, racemosus and B lumped together
 - Justicia white and Justicia dichipteroides lumped together
 - Leucas glabrata and Leucas martinensis lumped together
 - Lily A and Lily B lumped together
 - Sedges (Cyperus, Mariscus and Kyllinga) lumped together
 - After lumping, multi-species taxa include:
 - Commelina spp
 - Eragrostis spp
 - Sedges
 - Hibiscus flavifolius/aponeurus
 - Leucas spp
 - Melhania spp
 - Plectranthus spp
 - Pseudognaphalium/Helichrysum
 - Fern (<5%)
 - Justicia spp (<5%)
 - Lily spp (<5%)
- 5% of surveys sheet: Binary designations for whether a species occurred or not (in each plot in each year). Then calculation of the percentage of "surveys" (plots*time) that species occurred in.
- **Spp >5 % sheet**: first hits for only those species that occurred in at least 5% of all sampled plot*time combinations.

"Rare": First hits (rare spp) p/a	drep & Hib	Rare	drep & Hib						
drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps
Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib
drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a
Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib
drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps	drep & Hib	First & all hits, all spp, p/a; dreps
Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib
drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a	drep & Hib	Normal: First hits, all spp.; p/a
Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib
drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare	drep & Hib	Rare

Legend

Drep & Hib=*Acacia drepanolobium* and *Hibiscus* sampling: all 100 stations, **including below:**

Rare=First hits on all but five common grasses, plus presence of all spp (in 0.5 and 1.0 m plots)

Normal= First hits on all spp., plus presence of all spp. (in 0.5 and 1.0 m plots)

All hits=First and All hits on all spp, leaves and stems, & presence of all spp (in 0.5 and 1.0 m plots)

Figure 2. Vegetation sampling layout within KLEE of the 100 sampling points in each plot. Vegetation sampling is done within the central hectare (100 m x 100 m) of each KLEE plot. All KLEE plots contain a grid of rebar stakes all spaced 10 m apart. Vegetation sampling is conducted from the same sampling stations each time.

References

- Ahn, P. M., and L. C. Geiger. 1987. Soils of Laikipia District. Ministry of Agriculture, National Agricultural Laboratories, Kenya Soil Survey, Nairobi, Kenya.
- Odadi, W. O., T. P. Young, and J. B. Okeyo-Owuor. 2007. Effects of wildlife on cattle diets in Laikipia rangeland, Kenya. Rangeland Ecology & Management 60:179–185.
- Porensky, L. M., S. E. Wittman, C. Riginos, and T. P. Young. 2013. Herbivory and drought interact to enhance spatial patterning and diversity in a savanna understory. Oecologia 173:591–602.
- Veblen, K. E., L. M. Porensky, C. Riginos, and T. P. Young. 2016. Are cattle surrogate wildlife? Savanna plant community composition explained by total herbivory more than herbivore type. Ecological Applications.
- Young, T. P., B. Okello, D. Kinyua, and T. M. Palmer. 1998. KLEE: a long-term multi-species herbivore exclusion experiment in Laikipia, Kenya. African Journal of Range and Forage Science 104:92–104.
- Young, T. P., T. M. Palmer, and M. E. Gadd. 2005. Competition and compensation among cattle, zebras, and elephants in a semi-arid savanna in Laikipia, Kenya. Biological Conservation 122:351–359.

Include a list of KLEE publications?:

KLEE Publications (E numbers are from Exclosures themselves)

- 1. Young, T.P., N. Patridge & A. Macrae. 1995. Long-term glades in acacia bushland and their edge effects in Laikipia, Kenya. Ecological Applications 5: 97-108.
- 2. Young, T.P. 1995. Landscape mosaics created by canopy gaps, forest edges and bushland glades. Selbyana 16:127-134.
- 3. Young, T.P., C.H. Stubblefield & L.A. Isbell. 1997. Ants on swollen-thorn acacias: species coexistence in a simple system. Oecologia 109:98-107.
- 4. Young, T.P. and B.D. Okello. 1998. Relaxation of an induced defense after exclusion of herbivores: spines *on Acacia drepanolobium*. Oecologia 115:508-513. (E1)
- 5. Keesing, F. 1998. Ecology and behaviour of the pouched mouse, *Saccostomus mearnsi*, in central Kenya. Journal of Mammalogy 73:919-931. (E2)
- 6. Keesing, F.L. 1998. Impacts of ungulates on the demography and diversity of small mammals in central Kenya. Oecologia 116:381-389. (E3)
- 7. Young, T.P., B. Okello, D. Kinyua, & T.M. Palmer. 1998. KLEE: the Kenya long-term exclosure experiment. African Journal of Range and Forage Science 14:94-102. (E4)
- 8. Okello, D.N. & T.P. Young. 2000. Effects of fire, bruchid beetles and soil type on the germination and seedling establishment of *Acacia drepanolobium*. African Journal of Range and Forage Science 17:46-51.
- 9. Keesing, F. 2000. Cryptic consumers and the ecology of an African savanna. Bioscience 50:205-216. (E5)

- Metz M.R. & F. Keesing. 2001. Dietary choices of the pouched mouse (*Saccostomus mearnsi*) in central Kenya. Biotropica 33: 182-187. (E6)
- 11. Keesing, F., and T. Crawford. 2001. Impacts of density and large mammals on space use by the pouched mouse (*Saccostomus mearnsi*) in central Kenya Journal of Tropical Ecology 17:465-472. (E7)
- Gadd, M.E., T.P. Young & T.M. Palmer. 2001. Effects of simulated shoot and leaf herbivory in *Acacia drepanolobium*. Oikos 92:515-520. (E8)
- Okello, B.D., T.G. O'Connor, & T.P. Young. 2001 Growth, biomass estimates, and charcoal production of *Acacia drepanolobium* in Laikipia, Kenya. Forest Ecology & Management 142:143-153.
- 14. Shaw, M.T., F. Keesing, and R.S. Ostfeld. 2002. Herbivory on acacia seedlings in an East African savanna. Oikos 98:385-392. (E9)
- Ward, D. & T.P. Young. 2002. Effects of large mammalian herbivores and ant symbionts on condensed tannins of *Acacia drepanolobium* in Kenya. Journal of Chemical Ecology 28:913-529. (E10)
- 16. Young, T.P., M.L. Stanton & C. Christian. 2003. Effects of natural and simulated herbivory on spine lengths of *Acacia drepanolobium* in Kenya. Oikos 101:171-179. (E11)
- Warui, C. M., M.H. Villet, & T.P. Young. 2004. Spiders (Araneae) from black cotton soils of a highland savanna in Laikipia, central Kenya. Journal of Afrotropical Zoology 1:13-24. (E12)
- Goheen J.R., F. Keesing, B.F. Allan, D. Ogada, and R.S. Ostfeld. 2004. Net effects of large mammals on Acacia seedling survival in an African savanna. Ecology 85:1555-1561. (E13)
- 19. Huntzinger, P.M., R. Karban, T.P. Young & T.M. Palmer. 2004. Relaxation of induced indirect defenses of acacias following removal of mammalian herbivores. Ecology 85:609-614. (E14)
- 20. Warui, C.M., M.H. Villet, T.P. Young & R. Jocqué. 2005. Influence of grazing by large mammals on the spider community of a Kenyan savanna biome. Journal of Arachnology 33:269-279. (E15)
- 21. Gadd, M.E. 2005. Conservation outside of parks: attitudes of local people in Laikipia, Kenya. Environmental Conservation 32:1-14.
- 22. Young, T.P., T.M Palmer & M.E. Gadd. 2005. Competition and compensation among cattle, zebras, and elephants in a semi-arid savanna in Laikipia, Kenya. Biological Conservation 122:351-359. (E16)
- 23. Young, T.P. 2006. Declining rural economies and the future of biodiversity: missing the forest for the trees? Journal of International Wildlife Law & Policy 9:319-334.
- McCauley, D.J., F. Keesing, T.P. Young, J. Peters & B.F. Allan. 2006. Indirect effects of large herbivorous mammals on snakes in an African savanna. Ecology 87:2657-2663. (E17)
- 25. Isbell, L.A. and T. P. Young. 2007. Interspecific and temporal variation in domatia contents of the ant-plant *Acacia drepanolobium*, a staple food for patas monkeys (*Erythrocebus patas*) in Laikipia, Kenya. American Journal of Primatology 69:1387-1398.

- Goheen, J.R., T.P. Young, F. Keesing & T.M. Palmer. 2007. Consequences of herbivory by native ungulates for reproduction of a savanna tree. Journal of Ecology 95:129-138. (E18)
- Young, T.P. and D.J. Augustine. 2007. Intraspecific variation in the effects of large mammalian herbivores on reproduction in *Acacia* species in Laikipia, Kenya. Biotropica 39:559-561.
- 28. Pringle, R.M., T.P. Young and D.I. Rubenstein. 2007. Primary productivity and the strength of herbivore-initiated interaction cascades. PNAS 104:193-197. (E19)
- Odadi, W., T.P. Young, & J. Okeyo-Owuor. 2007. The effects of wild herbivores on cattle diet in Laikipia rangeland, Kenya. Rangeland Ecology and Management 60:179-185. (E20)
- 30. Riginos, C. and T.P. Young. 2007. Positive and negative effects of grasses and wild and domestic herbivores on Acacia saplings. Oecologia 153:985-995. (E21)
- 31. Riginos, C. & J.B. Grace. 2008. Savanna tree density, herbivores, and the herbaceous community: bottom-up vs. top-down effects. Ecology 89:2228-2238.
- 32. Veblen, K.E. 2008. Season- and herbivore-dependent competition and facilitation in a semiarid savanna. Ecology 89:1532-1540. (E22)
- Ogada, D., M.E. Gadd, R.S. Ostfeld, T.P. Young and F. Keesing. 2008. Impacts of large herbivores on bird diversity and abundance in an African savanna. Oecologia 156:387-397. (E23)
- Palmer, T.M., M.L. Stanton, T.P. Young, J.R. Goheen, R.M Pringle and R. Karban. 2008. Breakdown of an ant-plant mutualism following the loss of large herbivores from an African savanna. Science 319:192-195. [See also Science 319:xxx.] (E24)
- 35. McCauley, D.J., F. Keesing, T.P. Young & K. Dittmar. 2008. Effects of the removal of large herbivores on fleas of small mammals. Journal of Vector Ecology 33:263-268. (E25)
- 36. Sensenig, R.L. 2008. Fire Ecology in Laikipia, Kenya. Booklet distributed to Laikipia landowners, managers, and stakeholders. 79pp
- Okello, B.D., T.P. Young, C. Riginos, D. Kelly & T. O'Connor. 2008. Short-term survival and long-term mortality of *Acacia drepanolobium* after a controlled burn in Laikipia, Kenya. African Journal of Ecology 46:395-401.
- Riginos, C. 2009. Grass competition suppresses acacia growth across multiple demographic stages. Ecology 90:335-340.
- Odadi, W.O., T.P. Young & J.B. Okeyo-Owuor. 2009. Behavioural responses of cattle to shared foraging with wild herbivores in an East African rangeland Applied Animal Behavior Science 116:120-125. (E26)
- 40. Riginos, C., J.B. Grace, D.J. Augustine, & T.P. Young. 2009. Local versus landscape-scale effects of savanna trees on grasses. Journal of Ecology 97:1337-1345.
- 41. Goheen, J.R., T.M. Palmer, F. Keesing, C. Riginos, and T.P. Young. 2010. Large herbivores facilitate savanna tree establishment via diverse and indirect pathways. Journal of Animal Ecology 79:372-382. (E27)

- Brody, A.K. T. M. Palmer, K. Fox-Dobbs, & D.F. Doak. 2010. Termites, vertebrate herbivores, and the fruiting success of *Acacia drepanolobium*. Ecology 91:399-407. (E28)
- 43. Goheen, J.R. & T.M. Palmer. 2010. Defensive plant-ants stabilize megaherbivore-driven landscape change in an African savanna. Current Biology 20:1768-1772. (E29)
- Kinyua, D.M., L.E. McGeoch, N. Georgiadis, & T.P. Young. 2010. Short-term and longterm effects of tilling, fertilization, and seeding on the restoration of degraded rangeland in Laikipia, Kenya. Restoration Ecology 18S1:226-233.
- Veblen, K.E. & T.P. Young. 2010. Contrasting effects of cattle and wildlife on the vegetation development of a savanna landscape mosaic. Journal of Ecology 98:993-1001. (E30)
- 46. King, E.G. and K.K. Caylor. 2010. Herbivores and mutualistic ants interact to modify tree photosynthesis. New Phytologist 187:18-22. (E31)
- 47. Ogada, D.M. and F. Keesing. 2010. Decline of vultures over a three-year period in Laikipia, Central Kenya. Journal of Raptor Research 44:129-135. (E32)
- 48. Pringle, R.M., D.F. Doak, A.K. Brody, R. Jocque', and T.M. Palmer. 2010. Spatial pattern enhances ecosystem functioning in an African savanna. PLoS Biology 8(5):1-12.
- 49. Treydte, A.C., C. Riginos and F. Jeltsch. 2010. Enhanced use of beneath-canopy vegetation by grazing ungulates in African savannahs. Journal of Arid Environments 74:1597-1603.
- Gregory, N.C., R.L. Sensenig, D.S. Wilcove. 2010. Effects of controlled fire and livestock grazing on bird communities in East African savannas. Conservation Biology 24: 1606-1616.
- 51. Sensenig, R.L., M. Demment, & E.A. Laca. 2010. Allometric scaling predicts preferences for burned patches in a guild of East African grazers. Ecology 91:2898-2907.
- 52. Kuria, S.K., M.H. Villet, T.M. Palmer, & M.L. Stanton. 2010. A comparison of two sampling methods for surveying mammalian herbivore impacts on beetle communities in the canopy of *Acacia drepanolobium* in Kenya. African Entomology 18:87-98. (E33)
- Fox-Dobbs, K., D. F. Doak, A. K. Brody, and & T. M. Palmer. 2010. Termites create spatial structure and govern ecosystem function in an East African savanna. Ecology 91:1296– 1307. (E34)
- 54. Porensky, L.M. 2011. When edges meet: interacting edge effects in an African savanna. Journal of Ecology 99:923-934.
- Augustine, D.J., T.P. Young, K.E. Veblen, C. Riginos & J.R. Goheen. 2011. Pathways for positive cattle-wildlife interactions in semi-arid rangelands. Smithsonian Contributions to Zoology 632:55-71. (E35)
- Pringle, R.M., T.M. Palmer, J.R. Goheen, D.J. McCauley, & F. Keesing. 2011. Ecological importance of large herbivores in the Ewaso ecosystem. Smithsonian Contributions to Zoology 632:43-53. (E36)
- 57. Maclean, J.E., J.R. Goheen, T.M. Palmer, & T.P. Young. 2011. Small mammals limit tree population growth in an African savanna. Ecology 92:1626–1636. (E37)

- Odadi, W.O., M. Jain, S.E. Van Wieren, H.H.T. Prins and D.I. Rubenstein. 2011. Facilitation between bovids and equids in an African savanna. Evolutionary Ecology Research 13:237-252.
- 59. Odadi, W.O., A.S. Abdulrazak, M.M. Karachi & T.P. Young. 2011. African wild ungulates compete with or facilitate cattle depending on season. Science 333:1753-1755. (E38)
- 60. Riginos, R., L.M. Porensky, K.E. Veblen, W.O. Odadi, R.L. Sensenig, F. Keesing, D. Kimuyu, M.L. Wilkerson, and T.P. Young[†]. 2012. Lessons on the relationship between pastoralism and biodiversity from the Kenya Long-term Exclosure Experiment (KLEE). Pastoralism: Research, Policy and Practice 2012 3:1. (E39)
- 61. Porensky, L.M. and K.E. Veblen. 2013. Grasses and large herbivores reinforce landscape heterogeneity by excluding trees from ecosystem hotspots. Oecologia 168:749-759.
- 62. Veblen, K.E. 2012. Savanna glade hotspots: plant community development and synergy with large herbivores. Journal of Arid Environments 78:119-127.
- 63. Kuria, S.K. and M.H. Villet. 2012. The role of ants and mammalian herbivores on the structure and composition of insect communities found on canopies of *Acacia drepanolobium*. African Journal of Agricultural Research 7:5317-5331. (E40)
- 64. Odadi, W.O., S.A. Abdulrazak, M.M. Karachi, & T.P. Young. 2013. Protein supplementation-driven shifts in forage selection by cattle: implications for cattle wildlife coexistence. Ecological Applications 23:455-463. (E41)
- 65. Donihue, C.M., L.M. Porensky, J. Foufopoulus, C. Riginos and R.M. Pringle. 2013. Glade cascades: indirect legacy effects of pastoralism enhance the abundance and spatial structuring of arboreal fauna. Ecology 94:827-837.
- Porensky, L.M., S.F. Bucher, K.E. Veblen, A.C. Treydte, and T.P. Young. 2013. Megaherbivores and cattle alter edge effects around ecosystem hotspots in an African savanna. Journal of Arid Environments 96:55-63. (E42)
- 67. Porensky, L.M. & T.P. Young. 2013. Edge interactions in fragmented and patchy landscapes. Conservation Biology 27:509-519.
- Bergstrom, B.J. 2013. Would East African savanna rodents inhibit woody encroachment? Evidence from stable isotopes and microhistological analysis of feces. Journal of Mammalogy 94:436-447.
- Young, H.S., DJ. McCauley, E. Castillo, J.R. Goheen, K.M. Helgen, T.M. Palmer, R.M. Pringle, T.P. Young, R. Dirzo. 2013. Plant community response to changes in herbivore abundance and identity across abiotic gradients and experimental contexts. Journal of Ecology 101:1031-1041. (E43)
- Veblen, K.E. 2013. Impacts of traditional livestock corrals on woody plant communities in an East African savanna. The Rangeland Journal 35:349–353.
- Porensky, L.M., S.E. Wittman, C. Riginos & T.P. Young. 2013. Herbivory and drought interact to enhance diversity and spatial patterning in a savanna understory. Oecologia 173:591-602. (E44)

- 72. 2. Keesing, F., B.F. Allan, T.P. Young, and R.S. Ostfeld. 2013. Effects of wildlife and cattle on tick abundance in central Kenya. Ecological Applications 23: 1410–1418. (E45)
- 3. Wilkerson, M.L., L.M. Roche, & T.P. Young. 2013. Indirect effects of domestic and wild herbivores on butterflies in an African savanna. Ecology and Evolution 3:3672-3682. (E46)
- 4. Kirigia, A., J.T. Njoka, P.D Kinyua & T.P. Young[†]. 2013. Characterizations of livestock manure marketing chain and the annual income contribution of manure trade in Mukogodo, Laikipia, Kenya. African Journal of Agricultural Research 8:5864-5871.
- 75. a. McNally, Amy, Chris Funk, Greg Husak, Joel Michaelsen, B. Cappelaere, J. Demarty, T. Pellarin, T.P. Young, K.K. Caylor, C. Riginos & K.E. Veblen. 2013. <u>"Estimating Sahelian and East African soil moisture using the Normalized Difference Vegetation Index"</u>. Hydrology and Earth System Sciences Discussions 10, 7963-7997.
- 76. 5. Petipas, R.H. and A.K. Brody. 2014. Termites and ungulates affect arbuscular mycorrhizal richness and infectivity in a semi-arid savanna. Botany 92:233-240. (E47)
- 77. 6. Young, H. S., R. Dirzo, K. M. Helgen, D. J. McCauley, S. Billeter, M. Kosoy, L. Osikowicz, T. P. Young, and K. Dittmar. 2014. Declines in large wildlife increase landscape-level prevalence of rodent-borne disease. Proceedings of the National Academy of Sciences 111:7036-7041. (E48)
- 1. Kimuyu, D.K., R.L. Sensenig, C. Riginos, K.E. Veblen and T.P. Young[†]. 2014. Wild and domestic browsers and grazers reduce fuels, fire temperatures, and acacia ant mortality in an African savanna. Ecological Applications 24:741-749. (E49)
- 79. 2. Keesing, F. and T.P. Young. 2014. Cascading consequences of the loss of large mammals in an African savanna. Bioscience 64:487-405. (E50)
- 3. Dirzo, R., H.S. Young, M. Galetti, G. Ceballos, N.J.B. Isaac and B. Collen. 2014. Defaunation in the Anthropocene. Science 345:401-406. (NSF not acknowledged; see Figure 4[10-15% of the paper].) (E51)
- 4. Ford, A.T., J.R. Goheen, T.O. Otieno, L. Bidner, L.A. Isbell, T.M. Palmer, D. Ward, R. Woodroffe, and R.M. Pringle. 2014. Large carnivores make savanna tree communities less thorny. Science 246:346-349. (NSF not acknowledged; used our tannin data)
- S. Young, H.S., D.J. McCauley, R. Dirzo, J.R. Goheen, B. Agwanda, E.O. Castillo, A. Ferguson, S.N. Kinyua, M. McDonough, T.M. Palmer, R.M. Pringle, T.P. Young, and K.M. Helgen. 2015. Context-dependent effects of large mammals declines on small mammal communities in central Kenya. Ecological Applications 25:348-360. (E52)
- 6. Bonachela, R.M. Pringle, E. Sheffer, T.C. Coverdale, J.A. Guyton, K.K. Caylor, S.A. Levin, C.E. Tarnita. 2015. Termite mounds can increase the robustness of dryland ecosystems to climatic change. Science 347:651-655. (Used KLEE-funded satellite imagery)
- Riginos, C. 2015. Climate alters the landscape of fear in an African savanna. Journal of Animal Ecology 84:124-133.

- 8. Porensky, L.M. and K.E. Veblen. 2015. Generation of ecosystem hotspots using shortterm cattle corrals in an African savanna. Rangeland Ecology & Management 68:131-151.
- 9. Ngatia, L.W., B.L.Turner, J.T. Njoka, T.P. Young, and K.R. Reddy. 2015. The effects of herbivory and nutrients on plant biomass and carbon storage in vertisols of an East African savanna. Agriculture, Ecosystems & Environment 208:55–63. (E53)
- 10. Young, H,S., R. Dirzo, D.J. McCauley, B. Agwanda, L. Cattaneo, K. Dittmarjj, R.P. Eckerlin, R.C. Fleischer, L.E. Helgen, A. Hintz, J. Montinieri, S. Zhao, K.M. Helgen. 2015. Drivers of intensity and prevalence of flea parasitism on small mammals in East African savanna ecosystems. Journal of Parasitology 101:327-335. (E54)
- Palmer, T.M., E.G. Pringle, A. Stier, and R.D. Holt. 2015. Mutualism in a community context. Pp. 159-180 in: *Mutualism*. J.L. Bronstein (ed). Oxford University Press. (E55) (NSF not cited)
- Pringle, R.M., D. Kimuyu, R.L. Sensenig, K.E. Veblen, C. Riginos, T.M. Palmer, and T.P. Young. 2015. Synergistic indirect effects of elephants and fire in an African savanna. Journal of Animal Ecology 84:1637–1645. (E56)
- Young, H.S., R. Dirzo, K.M. Helgen, D.J. McCauley, C. Nunn, P. Snyder, K.E. Veblen, S, Zhao, and V.O. Ezenwa. 2016. Large wildlife removal drives redistribution of immune defenses in rodents. Functional Ecology, Online Early (E57)
- 90. Porensky, L.M. and T.P. Young. 2016. Development of edge effects around experimental ecosystem hotspots is affected by edge density and matrix type. Landscape Ecology, in press.
- 91. Veblen, K.E., L.M. Porensky, C. Riginos, and T. P. Young. 2016. Are cattle surrogate wildlife? Savanna plant community composition explained by total herbivory, not herbivore identity. Ecological Applications, in press. (E58)
- xx. Porensky, L.M. 201x. Biodiversity benefits of traditional livestock management are lowest at intermediate levels of landscape patchiness in an African savanna. Biological Conservation, in review.
- xx. Kimuyu, D.M., K.E. Veblen, C. Riginos, and T.P. Young. 201x. Cattle competition with grazing and browsing wildlife varies with season and presence of megaherbivores. Ecological Applications, in review. (E5x)
- xx. Charles, G., L.M. Porensky, K.E. Veblen, C. Riginos, and T.P. Young. 201x. Effects of different large herbivores on above-ground net primary productivity: grazing intensity affects the mean, but herbivore identity affects variability. Ecological Applications, in review. (E5x)

African Journal of Range and Forage Science 14:92-104 (1998)

KLEE: a long-term multi-species herbivore exclusion experiment in Laikipia, Kenya

Truman P. Young^{1,2,4}, Bell Okello^{2,3}, David Kinyua², Todd Palmer^{2,4}

Abstract. Livestock and wildlife share muches their respective ranges throughout the semi-arid ecosystems of the world. As the profitability of livestock production becomes more marginal and wildlife values increase, there is a need to understand the interactions between livestock and wild large mammalian herbivores (and other indigenous biodiversity). To address this, we have established a long-term multi-species herbivore exclusion experiment in the Laikipia ecosystem in Kenya. Using a series of 'semi-permeable' barriers, we are differentially excluding various combinations of cattle, large wild mammalian herbivores, and 'mega-herbivores' (giraffes and elephants) from a series of replicated four hectare plots. We are monitoring soil characteristics, tree and shrub population structure, herb layer vegetation structure and composition, primary productivity, range use by non-excluded large mammals, physical and chemical plant defense, populations of additional animal taxa (rodents, birds, grasshoppers), and the community of acacia ant symbionts. This multi-disciplinary project is one of the first to include controlled, replicated exclusion of combinations of multiple guilds of rangeland herbivores in the same place at the same time. Established in 1995, the exclosures offer opportunities for collaboration for a wide variety of applied and basic ecologists. We report here 1) details of the experimental design, 2) quantitative vegetation analysis of this important grazing ecosystem, 3) evidence of the effectiveness of the herbivore exclosures, and 4) a summary of some preliminary results.

Key Words: cattle, livestock, exclosure, wildlife, megaherbivore, elephant, giraffe, fencing, Africa

¹Department of Environmental Horticulture, University of California, Davis, CA USA 95616.

²Mpala Research Centre, P.O. Box 555, Nanyuki, KENYA.

³Department of Range and Forage Science, University of Natal, South Africa.

⁴Ecology Graduate Group, University of California, Davis, CA, USA 95616.

Introduction

Rangeland management is undergoing re-evaluation worldwide. There is increasing emphasis on management activities that are both sustainable and more compatible with native biodiversity. As the profitability of livestock production becomes more marginal and wildlife values increase, there is a growing need to understand the interactions between livestock and wild large mammalian herbivores and other indigenous biodiversity.

Semi-arid ecosystems cover the majority of the land area in sub-Saharan Africa, and the majority of these are acacia savannas (Cole 1986, Menault et al. 1985). These ecosystems have been used (and manipulated) by people of traditional cultures for many centuries, and are currently under rapidly increasing pressure for both intensive cattle production and arid-land farming (West 1971, Bernard et al. 1989, Gichohi et al. 1996). These pressures can result in local, regional, and global environmental problems (Gichohi et al. 1996, Herlocker 1996), including soil erosion, threats to endangered species such as elephants and rhinos, and desertification, with its potential effects on global warming. A detailed understanding of the ecology and dynamics of these ecosystems is critical to long-term development, management and conservation (Bourliere 1983, Tothill and Mott 1985, Hansen et al. 1986, Walker and Menault 1988, Christensen et al. 1996).

Virtually all indigenous large mammal species in Kenya have more individuals on multiple-use lands (mostly on semi-arid range lands) than on land inside parks and reserves (Mbugua 1986, Western 1989). The future of these populations depends on the interaction between enlightened rangeland management and wildlife needs. Interactions between livestock and wildlife in Africa are often negative (reviewed in MacMillan 1986, Prins 1992). There is growing evidence, however, that many wildlife species are not incompatible with moderate livestock production, and can even be beneficial. Because wildlife can provide much needed additional revenue in the form of tourism or game ranching, a mixed strategy of wildlife and cattle may be economically optimal and help maintain biodiversity (Hopcraft 1990). Opinions on the effects

of livestock production on biodiversity in the western United States are similarly disparate (Brown and McDonald 1995, Fleischner 1995).

African savannas are among the most intensively studied ecosystems in the world (e.g., Sinclair and Norton-Griffiths 1979, Huntley and Walker. 1982, Bourliere 1983, Tothill and Mott 1985, Cole 1986, Proctor 1989, Sinclair and Arcese 1995, Gichohi et al. 1996). Various factors determine the structure and function of savannas, including soil chemistry (Bell 1982, Chapin and McNaughton 1989, Hogberg 1989), small mammals (Belsky 1984, Happold 1983) and invertebrates (Josens 1983). However, rainfall, fire, native herbivores, and livestock are the dominant forces in short-term and long-term savanna community dynamics, and have been the subjects of numerous experimental and descriptive studies (e.g., Kelly and Walker 1976. Lock 1977. Hatton and Smart 1984. O'Connor 1985, Ernst and Tolsma 1989, Moe et al. 1990, Belsky 1992, Stuart-Hill 1992). Considerable literature exists on the effects of various cattle grazing and burning strategies on range quality in Africa (reviewed in O'Connor 1985), and on the effects of indigenous herbivores on the vegetation of protected areas (reviewed in McNaughton 1979, Gordon and Lindsay 1990, McNaughton and Banyikwa 1995, Sinclair 1995, Gichohi et al. 1996).

However, little is known about the interactions among wildlife, domestic livestock, and vegetation in Understanding these interactions requires Africa. controlled experiments that simultaneously manipulate multiple potential causative factors, and long-term monitoring of these experiments. Despite the high densities of native herbivores on many commercially managed rangelands, experimental approaches to the interactions of livestock, wildlife, and vegetation are rare (see Loft et al. 1987, Brown and Heske 1990, Hobbs et al. 1996a,b). There have been no controlled factorial experimental studies on the interactive effects of different domestic and native herbivores on rangeland vegetation in any African ecosystem (Werger 1977, Cumming 1982).

We have initiated the Kenya Long-term Exclosure Experiment (KLEE) in an attempt to address the interactions between livestock and native biodiversity on the rangeland that they share. This experiment is part of the broader Integrated Studies of Behavior and Ecology in the Laikipia Landscape (ISBELL) that includes research by a parallel team of behavioral ecologists.

Methods

Study area

The experiment is located at the Mpala Research Centre in Laikipia District, Kenya. The Centre is a collaborative trust dedicated to conservation in the Laikipia ecosystem and the welfare of its inhabitants. It is administered by a consortium consisting of George Small (the owner of Mpala Farm), the National Museums of Kenya, the Smithsonian Institution, the Kenya Wildlife Service, and Princeton University. The Centre maintains growing physical plant consisting of a laboratory, administrative offices, and full board accommodation for resident and visiting researchers. It is located on its own 1200ha parcel of land, and researchers have access to the 17,000ha Mpala Farm to the north. Several similar properties nearby also host research, most notably the adjacent Segera Ranch.

These properties are working cattle ranches that encourage native biodiversity. Their stocking rates are moderate (one livestock unit per 5-8ha), and below levels that are detrimental to the range. Year-to-year variation in rainfall is high, and averages 500mm (in the north) to 650mm (in the south), with peaks in April, July, and November. The native biodiversity on Mpala/Segera consists of an estimated 600-800 plant species (>450 collected thus far), more than 300 bird species (current list 280 spp.), and at least 70 mammal species, including 28 species of large herbivorous mammals (20 ungulates) and 19 species of large carnivorous (and insectivorous) mammals (see Appendix 1). Lists of rodents, bats, reptiles, butterflies, and grasshoppers are being compiled.

Two major soil types underlie the Laikipia On the high central plain where the ecosystem. experimental plots are located, level soils of impeded drainage predominate, especially deep clay 'black cotton' vertisols. Similar soils (with similar vegetation) occur at other sites of impeded drainage within Laikipia, and in many sites elsewhere in East Africa (including Nairobi National Park and the western extension of Serengeti National Park). These soils support some of the most productive rangelands in East Africa. Elsewhere in Laikipia, red rocky friable soils predominate on more sloping topographies. More information on the soils and vegetation of the red soil community can be found in Young et al. (1995). Approximately 10% of Mpala and virtually all of Segera are underlain by black cotton soils.

Both soil types are characterized by a landscape mosaic with numerous isolated 'glades'. These features are treeless, have high levels of mineral nutrients, and are preferentially used by wild and domestic herbivores (see Young et al. 1995; see also below). These glades are usually less than one hectare in area, but there are also some extensive areas ('plains') of glade vegetation, most notably on Segera. As in the red soils, glades on black cotton soils are higher in nutrients than are surrounding black cotton soils (Table 1), and are 'hot spots' of herbivore activity.

Compared with those on the red soils, black cotton communities are relatively depauperate in species for some but not all groups. The overstory is dominated by a single tree, Acacia drepanolobium, that accounts for >97% of the overstory cover in the study area (Young et al. 1997, see also below), and is always more than 50% of overstory cover throughout this and similar ecosystems (TPY, pers. obs. BO, unpubl. data). Five grass species and two forbs account for the vast majority of the understory (more than 90% of the relative cover). The fauna of both birds (J. Lynch, pers. comm.) and rodents (Keesing 1997, in press, in review) are also far less rich on the black cotton soils than on the adjacent red soils. However, the species richness of large ungulate herbivores (10-13 species) and carnivores (four species large enough to prey on ungulates) is equally great on both soil types (not including the specialized habitats of rivers, rocky outcrops, and escarpments).

Experimental design

Several conditions make this site ideal for our long-term large-scale experimental study: 1) the Mpala Research Centre (and its partners) are stable and supportive of this research; 2) the baseline ecology of this biome in general, and this ecosystem, are well-studied; 3) the wild and domestic mammalian herbivores are numerous and diverse; 4) the plant community on black cotton soils is both floristically simple; and 5) these widespread soils support some of the most productive East African rangelands.

Extensive preliminary surveys of the study areas within Mpala, carried out between 1992 and 1995, allowed us to identify suitable sites for the establishment of experiments, to maximize homogeneity among potential experimental plots, and to provide a baseline against which to measure long-term change. This background research consisted of the following studies:

- 1. Baseline surveys of the soils, vegetation, and herbivore use throughout the ecosystem. Some of these baseline data have already been incorporated into published papers (Young et al. 1995, 1997). We are also developing a plant checklist in collaboration with the National Herbarium and the Smithsonian Institution, a local herbarium for the Research Centre with reference collections, and a series of guides and field keys (both reproductive and vegetative) of all the plants found in the study area and in the entire Laikipia ecosystem. In addition, we are collecting daily rainfall and temperature records.
- 2. An in-depth study of the nutrient-rich glades that occur throughout the study area (Young 1995, Young et al. 1995), and that directed our decisions about the placement and size of the plots.

3. Baseline data on soils and vegetation at the specific sites of the plots before the exclosures were constructed (see below).



Herbivore barriers

We are using a series of 'semi-permeable' barriers to exclude different combinations of large mammalian herbivores. These barriers are similar to those designed by the Kenya Wildlife Services, and have a proven track record in East Africa (Hoare 1992), and in this ecosystem (Thouless and Sakwa 1995). Electric fencing is particularly effective (especially against destructive elephants) when a) the resident herbivores are already familiar with similar fences, avoiding a painful and expensive 'learning curve', and b) when the area exclosed is small, and each stretch of fence is relatively short (our longest linear section is 500m), allowing herbivores to go around rather than through. We use three types of barriers:

1. The 2.3m wildlife fence is eleven strands of bare wire, alternating live and ground. Every second, a burst of 6-7,000 volts is sent through the live wires via a battery charged by a solar panel. The lower nine wires are 16 cm apart, and the spacing for the upper two is 32 cm. The lowermost (ground) wire is at ground level. To minimize shorting of the lowest electric wire, the area along the bottom of the fence is kept free of vegetation by slashing and by regular applications of Round-up® on calm days during the growing seasons

using a backpack sprayer. In addition, any higher branches growing into the wires are cut back regularly. In three of the plots enclosed by this fence, there are 1.5m wide gates that can be opened to allow entry by cattle. These gates form a crush to aid in



pushing cattle. are hinged on the corner, so that the gate and the fence

2. The mega-herbivore fence (also called a 'dingledangle' fence) consists of a single live wire 2m off the ground, and two ground wires. One ground wire is located 32 cm above the live wire. The other runs along the ground, 60cm outside the plot (to increase the chances that an elephant or giraffe will be standing on it when contacting the live wire). Along the live wire, there are single wires ('dingle-dangles') 50 cm long hanging down every 50cm. This fence is not cleared for vegetation, except branches near the upper wires.

3. The cattle barriers are visual. Every 10m, there is a short post (60cm high) painted red on the cattle exclusion side. All cattle on the property travel in discrete herds of ~100 head and are accompanied by individual herders on foot. The cattle herders and the person documenting the cattle run use the painted post markers to ensure that cattle do not enter excluded areas. The cattle are allowed to graze on a controlled schedule comparable to that generally utilized on Mpala. The number of cattle and the duration of such grazing (in minutes) are quantified for each cattle 'run' inside the plots. Because there are no other large mammals inside the wildlife fences, we can use a physical barrier between areas where cattle are allowed or excluded. This is a single electrified wire 60cm off the ground, attached to short posts. The area below this wire is kept clear of vegetation.



In addition, nine permanent 3m x 7m rodent exclosures (and nine controls) have been placed in the total exclusion plots (Keesing 1997).

A large level tract of acacia wooded grassland of several hundred hectares in the black cotton soils was selected for the study. This site had no evidence of recent burning. Eighteen similar four-hectare plots (200m x 200m) were selected for the following treatments (see Figure 1):

1) Full fencing to exclude all large herbivores (three replicates).

2) Full fencing, but cattle allowed to graze periodically (three replicates).

3) High single-strand electric fencing to exclude elephants and giraffes only (three replicates).

4) As in #3, but cattle also excluded (three replicates).

5) Unfenced, cattle allowed to graze (three replicates)

6) Unfenced, cattle are not allowed to graze (three replicates).

We therefore have a complete $2 \ge 2$ factorial design for the effects of cattle and wildlife (including megaherbivores), and a complete $2 \ge 2$ design for the effects of cattle and non-mega-herbivore wildlife. (See 'Random stratified design' section, below.)

Our experience in this ecosystem and the experience of other ecologists indicate that three replicates per treatment will be sufficient, particularly since initial site selection was carefully carried out, and baseline data have been collected (cf. Brown and Heske 1990, Belsky 1992, Heske et al. 1994).

Because of the large size of the plots, each was demarcated into sixteen 50m x 50m subplots (see Figure 2). This allows accurate spatial information to be more easily gathered. For each plot, a variety of activities are allowed, with more manipulative activities being restricted to the edges (and even these are restricted to the minimal), and with no manipulations (only descriptive data collection) being allowed in the central hectare (see Figure 3). For any interventions allowed in the outer parts of each plot (and for the natural glades), edge effects will be quantified to protect against data biases in the innermost sampling hectare.

All eighteen plots have been bordered on their windward sides by fire breaks. Woody and herbaceous vegetation has been cleared from two parallel 5m wide strips, leaving a 2-4m buffer zone of intact vegetable along adjacent plots, and a wider buffer (20-30m) between fire breaks. These fire breaks are maintained by Mpala Farm using a road grader. The plots are large enough to include fire in sub-plot manipulations, and future decisions on experimental burning will be made in the context of ecosystem-wide patterns. Fences are also regularly patrolled to find and repair any breaks in the fence, and to test fence voltage. Any large herbivores (or their signs) seen in the plots during these patrols are recorded, and removed if inside exclosures meant to keep them out..

A stratified random design

The exclosure plots were located and oriented after we conducted extensive preliminary surveys and mapped the entire area for glades, areas of high and low Acacia drepanolobium density, and areas of differing typical tree height. These surveys, but not the baseline data (which were collected after plots locations were determined) were the basis of our decision for plot placement. We chose sites to be as homogeneous as possible, avoiding sites of low or high Acacia density. We also placed the exclosures as far as possible from graded tracks for security, and to minimize human disturbance and edge effects from the roads. We placed plot boundaries, whenever possible, across existing glades, so that different, randomly assigned treatments would occur within glades and be replicated across different glades. In addition, the six treatments sharing a type of barrier were placed so as to share a side (for cost reduction and maintenance simplicity), and similar 'access' to glades. Finally, individual plots that shared a fence were placed in differing (random) orientation to each other and to other pairs of plots (within each block).

The net result is a random stratified design that minimizes the chances that location and orientation effects (if any) will bias the results (Figure 1). An additional check against location bias is the baseline data collected before, during, and soon after the construction of the fences. These data indicate that different plots within the blocks (and to a lesser extent, the different blocks) are homogeneous for most edaphic, floristic, and physiognomic traits. Construction began in June 1995, and the exclosures have been fully operational since September 1995.

Caveats on the experimental design

We are using two cattle densities: none and moderate. Although we are considering small within-plot treatments with higher grazing intensities, we are not carrying out full treatment plots of multiple cattle densities, especially the high densities that mimic current densities in 'problem areas' and densities approaching or surpassing the carrying capacity of the land, for several reasons: 1) Many stocking density trials already have been carried out throughout Africa and in rangelands world-wide which attempt to determine which densities are associated with positive and negative environmental and economic outcomes, at least with respect to livestock production (for a review of African research, see O'Connor 1985). It is not our goal to duplicate that research. 2) We are not interested in asking how close we can get to the edge of sustainability without going over, because a) being close to this edge is by definition antithetical to the management goal of coexistence of livestock and a rich indigenous biodiversity, and b) the highly variable climates of these semi-arid and arid ecosystems make the results of such stocking experiments less than reliable for any real management use (Swift et al. 1996). 3) Instead, we are interested in the nature of the relationship between livestock and biodiversity on lands that are managed for both, which by definition are lands on which cattle densities are kept at moderate levels. 4) In addition, the current experimental design involving multiple combinations of three guilds of large herbivores is already fairly complicated (six treatment combinations), and adding even one additional level of cattle density to the two already in the design would increase the size of the project by 50%, and the managerial (and statistical!) problems would increase by even more.

Although the experiment is both large scale and long-term, it is being carried out at specific spatial (four hectare) and temporal (<20 years) scales, and this constrains the ecological issues we can and cannot address: 1) We <u>can</u> examine both functional and numerical responses of plants, invertebrates, rodents, and perhaps some birds to different herbivory

treatments. We cannot examine numerical responses of larger (>10 kg) herbivores or mammalian carnivores. 2) We can examine functional responses of birds, large herbivores, and perhaps mammalian carnivores ('Do they spend more time foraging in certain plots?'), from which we may risk models of numerical responses on a larger spatial scale. 3) We can address how year-toyear variation in climate affects the relationships among the various members of this community, and their shortterm sustainability. We probably cannot determine the sustainability of different herbivore long-term combinations, given the tremendous variation in climate on decadal scales. 4) We can ask how different herbivore combinations influence changes in vegetation structure and composition. We cannot fully describe the kinds of (endogenous or climatically driven) cyclical variation in vegetation that have been suggested to occur on the scale of several decades (Caughlev 1976, Dublin et al. 1990, Dublin 1995), although we may be able to ask which herbivores are associated with particular states or transitions in a putative ecological cycle.

Baseline data:

To maximize the power of the experimental design and opportunities for future research, we (and our collaborators) have generated several baseline data sets:

- 1. Soil structure and chemistry (pH and ten elemental nutrients) from both background vegetation and glade sites. Soils were taken from two depths (0-10 cm and 20-30 cm).
- 2. Frequency of herbaceous species from both background and glade sites, from 100 quadrats per plot.
- 3. Percent cover of herbaceous vegetation by species, from 100 pin frames (2500 pins) per plot.
- 4. Acacia drepanolobium density, size structure, and defense (thorn length, tannin content, and ant occupancy by species), from both fixed transects (n > 4000) and from stratified size sampling (n = 1800; see below).
- 5. Mapping and measurement of individuals of all other woody plant species.
- 6. Mapping of glade size, shape, and orientation.
- 7. Complete photographic coverage of the plots, from eight fixed points per plot, revisited every six months.
- (8. Rodent density by species, from 10 x 10 trapping grids- Keesing 1997).

The use of baseline data (as model covariates) and analyses of variance that explicitly incorporate block effects (to control for the remaining variation among blocks) will combine to maximize our statistical power to discern experimental effects. In addition, the replicated experimental design and diligent efforts to control for extraneous factors in the original locating of the plots should give us the power to ask about experimental effects even in the absence of baseline data (and will therefore be useful for later collaborators whose projects measure effects for which no baseline was collected).

Baseline data methods

Densities of woody species were assessed through a series of nine 5 x 50m transects in each of the 18 plots. Within each transect, each tree and shrub was identified to species, and its height and perpendicular crown diameters measured to the nearest 5cm. Herbivore utilization was assessed on 90 marked trees within each plot, stratified by tree height. On each tree, five branch tips were tagged and regularly surveyed for herbivory. Additional trees and shrubs were similarly tagged and monitored along the edges of glades.

Understory frequency was measured through three 300m line transects. Every 3m, a 0.5 x 0.5m quadrat was placed (alternating left and right), and all species recorded to species, for a total of 100 quadrats per plot. Additional (shorter) frequency transects were run within glades. Understory cover was assessed with nine 30 (or 33) m transects, surveyed every 3m with a ten-pin point frame, for a total of 100 frames (1000 pins) per plot. Plant species are being identified in collaboration with Dr. Christine Kabuye (Chief Botanist) and Mr. Joshua Mwasya of the Herbarium of the National Museums of Kenya, and Robert Faden of the Smithsonian Institution. Duplicate collection sheets are being put into an Mpala Research Centre Herbarium.

Dung densities were measured with six 50 x 2m transects in each plot. Within each transect, each dung pile was identified to species, with ambiguous cases collected and refereed to an experienced tracker. Differences between dung of cattle and cape buffalo, and between Grevy's and Burchell's zebras, could not be distinguished reliably (Stuart and Stuart 1994); therefore all dung typical of these species was classified as 'cattle' or 'zebra'. respectively.

Statistical analysis

Total aerial cover, cover by each species, and the densities of *A. drepanolobium* and other woody plants were analyzed with a Model 1 ANOVA (10 error d.f.) for the effects of blocks (2 d.f.) and treatments (5 d.f.). Cluster analysis was done using Ward's Method (JMP).

Results

Soils

The non-glade soils in the study plot are extremely heavy in clay, with values (range 52-64%) that are at the upper end of observed clay content for natural soils. As in the red soils, the glade vegetation is underlain with soils that are richer in most mineral nutrients (Table 1). Glade soils also have less clay than nonglade soils. The black cotton soils at the study site contain 60% clay on average, with lower values (mean, 45%) in the glades (F = 29.4, p < 0.001).

Table 1. Mean soil characteristics (\pm one S.E.) of glade and non-glade sites at the experimental plots, on black cotton soils. Soil was collected at 0-30cm in August 1995. Soil nitrogen (note smaller sample sizes) was analyzed at the Range Science Department of the University of Nairobi from samples collected at 10cm in September 1997.

Soil trait	Glades (N=15)	Background (N=6)
Physical	()	()
Clay (%)	45.2 <u>+</u> 1.6	59.2 <u>+</u> 0.7
Silt (%)	22.2 ± 0.9	14.8 ± 0.5
Sand (%)	32.6 <u>+</u> 0.9	26.0 <u>+</u> 0.4

Chemical (%, unless otherwise noted)

Total N	0.28 <u>+</u> 0.06 (N=2)	0.17 ± 0.02 (N=5)
С	1.95 <u>+</u> 0.11	1.43 <u>+</u> 0.13
P (ppm)	269 <u>+</u> 21	60 <u>+</u> 2
Κ	3.11 <u>+</u> 0.12	1.15 <u>+</u> 0.11
Ca	16.9 <u>+</u> 1.0	9.7 <u>+</u> 0.4
Na	1.99 <u>+</u> 0.06	1.50 ± 0.08
Mn	1.56 <u>+</u> 0.12	1.86 <u>+</u> 0.06
Mg	4.16 <u>+</u> 0.12	3.83 <u>+</u> 0.07
pН	6.72 <u>+</u> 0.10	6.28 <u>+</u> 0.07

Overstory vegetation

Mean density of *A. drepanolobium* trees is 2267/ha, with no block and no herbivore treatment differing in density more than 22% from this mean. Analysis of variance detected no significant block (p = 0.41) or treatment (p = 0.84) effects in total density. Overall, the size distribution was strongly L-shaped (Figure 4), and was similar among blocks and among treatments. The mean height of *A. drepanolobium* trees in each of the three blocks was within 7% of the grand mean (1.38m).

Several other woody species occur in the plots, but at low densities: *Cadaba farinosa, Rhus natalensis, Acacia mellifera, A. brevispica, Balanites* sp., *Boscia* sp., *and Lippia javanica.* These species accounted for 3.2% of the woody plants less than 1.5m tall, and 0.7% of the woody plants more than 1.5m tall (2.3% overall). Many of the smaller plants of these species are represented by apparently old individuals (with stems up to 1.5 cm in diameter) that have been strongly suppressed by herbivory. Treatment (herbivore) classes did not differ in the densities of these other woody species (F = 0.55, p = 0.74), but the North block had a higher density of these rarer plants than did the Central and South blocks (F = 5.48, p = 0.025).

Understory vegetation

Baseline vegetation data demonstrate considerable homogeneity among blocks and among treatment classes. Total aerial cover was within 9% of the overall mean (58.5%) in each of the 18 plots, and there were no significant Block (p = 0.44) or Treatment (p = 0.88) effects. The five most abundant species, all grasses, accounted for 88% of the relative cover (in order: Pennisetum stramineum, Lintonia nutans, Themeda triandra, P. mezianum and Brachiaria lachnantha). With the exception of T. triandra, these five species had the same rank order in each of the three blocks. The most abundant forb was the semi-woody Aerva lanata. In the South block, T. triandra and another grass, Botriochloa insculpta, were significantly more abundant (p < 0.001, p = 0.04) than in the two other blocks, at the expense of the two *Pennisetum* spp. (p = 0.025, p < 0.0250.001). The semi-woody herb Helichrysum glumaceum was also more abundant in the South block (p = 0.04). No other species differed significantly in cover among Only B. lachnantha differed significantly blocks. among treatments, being more abundant in the three control (MWC) plots (p = 0.02). In any case, these baseline data will be used as covariates in any future analyses on the effects of the herbivore treatments.



Frequency data also indicate considerable homogeneity among blocks. Frequency measures are more sensitive to smaller and less common species, and more useful for multivariate community analysis in this depauperate system. Of the 25 species found in baseline surveys (in 1800 quadrats) only one varied significantly among blocks (*Pennisetum mezianum*). The eight most frequent herbs were the same in all three blocks, with differences in rank order only among species with very similar mean frequencies (Figure 5).



Figure 5. The mean frequencies, across all 18 plots, of the 25 most common herbaceous species (not including tran-sects through glade vegetation). The line is the cumulative relative frequency accounted for. The eight most frequent species account for 93% of the total species frequencies. There was strong homogeneity among blocks. In every block, the rank abundance of the first eight species was essentially the same. Bars are standard errors, based on a sample size of three (blocks).

analysis Cluster based on frequency data demonstrates the similarity of the background vegetation of plots and blocks, compared the vegetation of the glades (Figure 6). The Euclidean distance (9.7)from the average glade to the average non-glade vegetation was more than double the cumulative distance (4.1) among all the branches of the non-glade cluster. There is no strong pattern of similarity in baseline vegetation based either on treatment or on block.

As in the red soils (Young et al. 1997), the glade vegetation of the black cotton soils differs dramatically from background vegetation. The black cotton glades share several floristic traits with the red soil glades. First, they are also treeless (somewhat trivial, since this is our definition of 'glade'). Second, several of the glades have patches of Cynodon plectostachyus, virtually absent from other black cotton sites. Third, the dominant grass inside black cotton glades is Pennisetum stramineum, the species that characteristically ringed the glades in red soils. This grass accounted for more then three times as much relative frequency as the next most frequent taxon (Ipomoea spp.), and was even more dominant in terms of biomass (TPY, pers. obs.). Also common inside the glades are the grasses Setaria verticillata and Sporobolus sp., and (during rains) the annual forb Leucas martinicensis. The shrub Lyceum

europaeum, virtually absent away from the glades, was commonly found around their peripheries.

Note: Young et al. (1995) reported that *Digitaria milanjiana* was the dominant grass in the glades on red soils, based on vouchers from one glade on deposit at the National Herbarium. We now know that many of the plants later scored in the field as *D. milanjiana* were actually *Cynodon* spp., mainly *C. plectostachyus*.



Effectiveness of the herbivore barriers

In the first two years of the experiment, there were breaches of the game fence by single zebras on three ocasions, each detected and repaired within a day or two. There have also been three breaches of the megaherbivore fence by elephants. Each of these lasted only a short time, and in each, the area visited by elephants within the plot was carefully mapped. There was also a single breach of the mega-herbivore fence by a single giraffe.

Our goal is not necessarily the absolute exclusion of target herbivores, but large and well-documented decreases in their abundance. Our acacia utilization data and dung count data indicate that the barriers are achieving this goal. In March 1996, we carried out a preliminary survey of 2300 new shoots of *Acacia drepanolobium* in the plots. This survey showed the following patterns (Figure 7): There are no differences in acacia browsing between the plots with and without

cattle, confirming the claims of herders that cattle do not eat this species. In the unfenced areas, 4-6% of the shoot tips were browsed at both 1 and 2 m above the ground. Within the mega-herbivore exclusion fence, 5% of the shoots were eaten at 1 m, but less than 1% eaten at 2 m. Within the total game exclusion fence, less than 1% of the shoots tips were eaten at either 1 or 2 m above the ground. An additional survey on shoot herbivory was undertaken in June 1996 showed similar patterns (Young and Okello, in review).



Figure 7. Herbivory of new shoots of Acacia drepanolobium in different exclosure plots in March 1996. Plots with and without cattle were lumped, because cattle do not browse this tree. N = 220 trees (~2300 shoots). Treat-ment categories as in Figure 1. Bars represent one S.E., based on three blocks.

Our dung counts are not meant to be used for comparisons across species. In particular, the cattle are put into the plots only for specific short intervals, and probably defecate less during this period than wildlife species, which have continual access. However, these counts are likely to be appropriate measures of levels of presence within herbivore classes. All three barriers appear to be effective in excluded the target herbivores (Figure 8). The visual barriers used by the herders reduced the abundance of cattle by more than 95% (5.0 + 1.9 vs. 0.2 + 0.2, n = 3 blocks). The wildlife fences reduced the abundance of wild large mammalian herbivores by more than 95% (17.9 ± 5.0 vs. 0.5 ± 0.0). The mega-herbivore fences reduced the abundance of elephants and giraffes by 75% (2.0 + 0.5 vs. 0.5 + 0.2). Note also that the mega-herbivore fences did not apparently restrict the movement of other wildlife (Figure 8). In addition, elephant tracks were common outside the fences but rare inside, and restricted to the areas in the vicinity of recent breaks (pers. obs.).

There is constancy (homogeneity) of both dung count and herbivory rates within each treatment across all blocks. This strongly suggests that there is not a tendency for the experimental design to 'funnel' animals into any particular plot, nor any strong block or orientation effects on the animals that this experiment was designed to manipulate.



As an additional measure of effectiveness, in the time since the exclosures were fully operational, there already has been tremendous of growth of several nonacacia woody species inside the game fences. These include Cadaba farinosa, Rhus natalensis, Balanites aegyptiaca, Lippia javanica, and Maerua sp. The first two species still appear to be somewhat suppressed in the plots within these exclosures accessible to cattle (for *Cadaba*). The increase in all of these shrubs is from old (thick-stemmed) but very small and suppressed individuals already in the plots. This implies that the 'bush encroachment' we are seeing comes not from an increase in the number of individual shrubs, but from the release of a large number of suppressed shrubs previously inconspicuous in the grass (Young and Okello, unpublished data). This realization could have profound effects on how we deal with the problem of bush encroachment in this ecosystem.

Initial responses to different herbivore treatments.

We have already documented significantly different responses to herbivore treatments for several dependent variables. First are the differences in herbivore presence and utilization by browsers mentioned above, which demonstrate the effectiveness of the barriers. However, we have also quantified several non-trivial responses to the experimental treatments: 1) Rodent numbers were significantly higher in treatments with fewer herbivores (Keesing 1997 and in review), suggesting competition (and compensation) between two very different guilds of herbivores. 2) Keesing (1997 and in review) has also demonstrated increased vegetation cover in rodent exclosure plots. 3) We have demonstrated a 20% decline in the length of thorns produced by branches protected from herbivory, an effect limited to branch heights accessible to the

herbivores allowed (Young and Okello, in review). 4) Preliminary surveys indicate an increase in grasshopper densities in plots from which herbivores are excluded (Palmer, unpublished data). 5) As mentioned above, there has been a release from suppression of several species of woody plants in a form of bush encroachment that does not entail increased recruitment (Young et al., unpublished data).

Current and future collaborations.

KLEE operates in the spirit of collaboration. We encourage research by scientists with a broad variety of interests. We do insist on coordination among research groups, and on the archiving of data gathered in the plots. Large amounts of baseline data are available. Areas currently under-studied include wildlife and cattle foraging behavior, the ecology of invertebrates (especially termites, ticks, Lepidoptera) and (small) carnivores, and manipulations of sub-plots with fire or intense grazing. We are open to other kinds of basic and applied research.

List of project collaborators

- Truman P. Young, University of California (Project Director)
- Bell Dedan Okello, University of Natal (biology of *Acacia drepanolobium*)
- David Kinyua, University of Nairobi (herb layer dynamics)

Maureen Stanton, University of California (acacia ants)

- Felicia Keesing, Siena College and Institute for Ecosystem Studies (rodents)
- Todd Palmer, University of California (acacias and acacia ants)
- Robert Faden, Smithsonian Institution (plants)
- David Ward, Ben Gurion University of the Negev (acacia tannins)
- James Lynch, Smithsonian Institution (birds)
- Randy Dahlgren, University of California (biogeochemistry)
- Michelle Gadd, University of California (elephants)

Acknowledgments

This research was supported by the Smithsonian Institution (James Smithson grant to the late Alan P. Smith); the National Geographic Society (grant 4691-91), the University of California at Davis, Fordham University, Mpala Farm, the Mpala Research Centre, the University of California Research Expeditions Program (UREP) to T. Young; Citibank and International Federation of Science grants to B. Okello and D. Kinyua; and NSF grant BSR 93-07477 to L. Isbell. Field assistance was provided by F. Erei and J. Lemboi. Special thanks to George Small and John Wreford-Smith. The fencing was erected by Simon Dufresne (Sanyati Ltd.), who contributed much to its design. Helpful comments on the ms were made by M. Gadd.

Literature Cited

- Bell, R.H.V. 1982. The effect of soil nutrient availability on community structure in African ecosystems. Pp. 193-216 in: *The ecology of tropical savannas*. Huntley, B.J. and B. H. Walker (eds). Springer-Verlag, Berlin.
- Belsky, A.J. 1984. Role of small browsing mammals in preventing woodland regeneration in Serengeti National Park, Tanzania. *African Journal of Ecology* 22:271-279.
- Belsky, A.J. 1992. Effects of grazing, competition, disturbance and fire on species composition in grassland communities. *Journal of Vegetation Science* 3:187-200.
- Bernard, F.E., D.J. Campbell, and D.J. Thom. 1989. Carrying capacity of the eastern ecological gradient of Kenya. *National Geographic Research* 5:399-421.
- Bourliere, F., (ed.). 1983. Ecosystems of the world 13. Tropical savannas. Elsevier, Amsterdam.
- Brown, J. H. and E.J. Heske. 1990. Control of a desertgrassland transition by a keystone rodent guild. *Science* 250:1705-1707.
- Brown, J. H. and W. McDonald. 1995. Livestock grazing and conservation on southwestern rangelands. *Conservation Biology* 9:1644-1647.
- Caughley, G. 1976. The elephant problem an alternate hypothesis. *East African Wildlife Journal* 14:265-283.
- Chapin, F. S. III and S. J. McNaughton. 1989. Lack of compensatory growth under phosphorous deficiency in grazing adapted grasses from the Serengeti Plains. *Oecologia* 79:551-557.
- Christensen, N.L. et al. 1996. The Report on the Ecological Society of America Committee on the Scientific Basis for Ecosystem Management. *Ecological Applications* 6:665-691.
- Cole, M.M. 1986. The savannas. *Biogeography and geobotany*. Academic Press, London.
- Cumming, D.H.M. 1982. The influence of large herbivores on savanna structure in Africa. Pp. 217-245 in: *The ecology of tropical savannas*. Huntley, B.J. and B.H. Walker (eds). Springer-Verlag, Berlin.
- Dorst, J. and P. Dandelot. 1972. A field guide to the larger mammals of Africa. Collins, London.
- Dublin, H.T. 1995. Vegetation dynamics in the Serengeti-Mara ecosystem: the role of elephants, fire, and other factors. Pp. 71-90 in: Serengeti II: dynamics, management, and conservation of an ecosystem. Sinclair, A.R.E. and P. Arcese (eds). University of Chicago Press.
- Dublin, H.T., A.R.E. Sinclair, and J. McGlade. 1990. Elephants and fire as causes of multiple stable states in the Serengeti-Mara woodlands. *Journal of Animal Ecology* 59:1147-1164.
- Ernst, W.H.O. and D.J. Tolsma. 1989. Mineral nutrients in some Botswana savanna types. Pp. 97-120 in: *Mineral nutrients in tropical forest and savanna ecosystems*. Proctor, J. (ed.), Blackwell Scientific Publications, Oxford.
- Estes, R.D. 1991. *The behavior guide to African mammals.* University of California Press, Berkeley.

- Fleischner, T.L. 1995. Ecological costs of livestock grazing in western North America. *Conservation Biology* 8:629-644.
- Gichohi, H., E. Mwangi, and C. Gakahu. 1996. Savanna ecosystems. Pp. 273-298 in: *East African Ecosystems and their Conservation*, T.M. McClanahan and T.P. Young (eds). Oxford University Press, NY.
- Gordon, I.J. and W.K. Lindsay. 1990. Could mammalian herbivores "manage" their resources? *Oikos* 57:1270-280.
- Hansen, R.M., B.M. Woie, and R.D. Child (eds.) 1986. Range development and research in Kenya. Winrock International Institute for Agricultural Development. Morrilton, Arizona.
- Happold, D.C.D. 1983. Rodents and lagomorphs. Pp. 363-400 in: *Ecosystems of the world 13. Tropical savannas*. Bourliere, F. (ed.). Elsevier, Amsterdam.
- Hatton, J.C. and N.O.E. Smart. 1984. The effect of long-term exclusion of large herbivores an soil nutrients in Murchison Falls National Park, Uganda. *African Journal of Ecology* 22:23-30.
- Heske, E.J., J.H. Brown, and S. Mistry. 1994. Long-term experimental study of a Chihuahuan Desert community: 13 years of competition. *Ecology* 75:438-445.
- Hoare, R.E. 1992. Present and future use of fencing in the management of larger African mammals. *Environmental Conservation* 19:160-164.
- Hobbs, N.T., D.L. Baker, G.D. Bear, and D.C. Bowden. 1996a. Ungulate grazing in sagebrush grassland: mechanisms of resource competition. *Ecological Applications* 6:200-217.
- Hobbs, N.T., D.L. Baker, G.D. Bear, and D.C. Bowden. 1996b. Ungulate grazing in sagebrush grassland: effects of resource competition on secondary production. *Ecological Applications* 6:218-227.
- Hogberg, P. 1989. Root symbioses of trees in savannas. Pp. 121-136 in: Proctor, J. (ed.), *Mineral nutrients in tropical forest and savanna ecosystems*. Blackwell Scientific Publications, Oxford.
- Hopcraft, D. 1990. Wildlife land use at the Athi River, Kenya. Pp. 332-340 in: *The Improvement of Tropical and Subtropical Rangelands*. National Academy Press, Washington, D.C.
- Huntley, B.J. and B.H. Walker. 1982. *The ecology of tropical savannas*. Springer-Verlag, Berlin.
- Herlocker, D.J. 1996. Rangeland degradation in arid and semi-arid East Africa. Pp. 266-267 in: *East African Ecosystems and their Conservation*, T.M. McClanahan and T.P. Young (eds). Oxford University Press, NY.
- Josens, G. 1983. The soil fauna of tropical savannas. III. The termites. Pp. 503-524 in: Bourliere, F. (ed). *Ecosystems of the world 13. Tropical savannas*. Elsevier, Amsterdam.
- Keesing, F.L. 1997. Ecological interactions between small mammals, large mammals, and vegetation in a tropical savanna of central Kenya. Ph.D. Diss., University of California at Berkeley.
- Keesing, F.L. Ecology of the pouched mouse in central Kenya. *Journal of Mammalogy*, in press.
- Keesing, F.L. Indirect effects of small mammals in a savanna ecosystem in central Kenya. In review.

- Kelly, R.D. and B.H. Walker. 1976. The effects of different forms of land-use on the ecology of a semi-arid region in south-eastern Rhodesia. *Journal of Ecology* 64:553-576.
- Kingdon, J. 1997. *The Kingdon field guide to African mammals*. Academic Press, New York.
- Lock, J.M. 1977. Preliminary results from fire and elephant exclusion plots in Kabalega National Park, Uganda. *East African Wildlife Journal* 15:229-232.
- Loft, E.R., J.W. Menke, J.G. Kie, and R.C. Bertram. 1987. Influence of cattle stocking rate on the structural profile of deer hiding cover. *Journal of Wildlife Management* 51:655-664.
- MacMillan, S. (ed.) 1986. Wildlife/livestock interfaces on rangelands. Inter-African Bureau for Animal Resources, Nairobi.
- Mbugua, S.W. 1986. Monitoring livestock and wildlife in Kenya pp. 413-430 in: *Range development and research in Kenya*. Hansen, R.M., B.M. Woie and R.D. Child (eds). Winrock International Institute for Agricultural Development, Morrilton, Arizona.
- McNaughton, S.J. 1979. Grassland-herbivore dynamics. Pp. 46-81 in: Serengeti: dynamics of an ecosystem. Sinclair, A.R.E. and M. Norton-Griffiths (eds). University of Chicago Press, Chicago.
- McNaughton, S.J. and F.F. Banyikwa. 1995. Plant communities and herbivory. Pp. 49-70 in: Serengeti II: dynamics, management, and conservation of an ecosystem. Sinclair, A.R.E. and P. Arcese (eds). University of Chicago Press.
- Menault, J.C., R. Barbault, P. Lavelle, and M. Lepage. 1985. African savannas: biological systems of humidification and mineralization. Pp. 14-33 in: *Ecology and management of the world's savannas*. Tothill, J.C. and J. J. Mott (eds). Australian Academy of Science, Canberra.
- Moe, S.R., P. Wegge, and E.B. Kapela. 1990. The influence of man made fires on large herbivores in Lake Burungi area in northern Tanzania. *African Journal of Ecology* 28:35-43.
- O'Connor, T. 1985. A synthesis of field experiments concerning the grass layer in the savanna regions of Southern Africa. *South African National Scientific Programme Reports* 114:1-126.
- Prins, H.H.T. 1992. The pastoral road to extinction competition between wildlife and traditional pastoralism in East Africa. *Environmental Conservation* 19:117-123.
- Proctor, J. (ed). 1989. Mineral nutrients in tropical forest and savanna ecosystems. Blackwell Scientific Publications, Oxford.
- Sinclair, A.R.E. 1995. Equilibria in plant-herbivore interactions. Pp. 91-114 in: Serengeti II: dynamics, management, and conservation of an ecosystem. Sinclair, A.R.E. and P. Arcese (eds). University of Chicago Press.
- Sinclair, A.R.E. and P. Arcese (eds). 1995. Serengeti II: dynamics, management, and conservation of an ecosystem. University of Chicago Press.
- Sinclair, A.R.E. and M. Norton-Griffiths, Eds. 1979. Serengeti: dynamics of an ecosystem. University of Chicago Press, Chicago.
- Skarpe, C. 1990. Shrub layer dynamics under different herbivore densities in an arid savanna, Botswana. *Journal* of *Applied Ecology* 27:873-885.

KLEE.-12

- Stuart, C. and T. Stuart. 1994. *A field guide to the tracks and signs of southern and East African wildlife*. Southern Book Publishers, Capetown.
- Stuart-Hill, G.C. 1992. Effects of elephants and goats on the Kaffrarian succulent thicket of the eastern Cape, South Africa. *Journal of Applied Ecology* 29:699-710.
- Swift, D.M., M.B. Cougenour and M. Atsedu. 1996. Arid and semi-arid ecosystems. Pp. 243-272 in: *East African Ecosystems and their Conservation*, T.M. McClanahan and T.P. Young (eds). Oxford University Press, NY.
- Taiti, S.W. 1992. The vegetation of Laikipia District, Kenya. Laikipia - Mt. Kenya Papers, B-2. Laikipia Research Programme, Universities of Nairobi and Bern.
- Thouless, C.R. and J. Sakwa. 1995. Shocking elephants: fences and crop raiders in Laikipia District, Kenya. *Biological Conservation* 72:99-107.
- Tothill, J.C. and J.J. Mott (eds). 1985. *Ecology and management of the world's savannas*. Australian Academy of Science, Canberra.
- Walker, B.H. and J.C. Menaut (eds). 1988. Research procedure and experimental design for savanna ecology and management. *Responses of savannas to stress and disturbance*, Publication #4. Printing Centre, Melbourne.
- Werger, M.J.A. 1977. Effects of game and domestic livestock on vegetation in East and southern Africa. Pp. 149-159 in: *Handbook of vegetation science*. Vol. 13. Krause, W. (ed). W. Junk, The Hague.
- West, O. 1971. Fire, man and vegetation as interacting factors limiting the development of climax vegetation in Rhodesia. *Proceedings of the Tall Timbers Fire Ecology Conference* 11:121-145.
- Western, D. 1989. Conservation without parks: wildlife in the rural landscape. Pp. 158-165 in: *Conservation for the Twenty-first Century*. D. Western and M.C. Pearl (eds). Oxford University Press, Oxford.
- Young, T.P. 1995. Landscape mosaics created by canopy gaps, forest edges, and bushland glades. *Selbyana* 16:127-134.
- Young, T.P. and B.D. Okello. Relaxation of an induced defense after exclusion of herbivores: spine length in *Acacia drepanolobium. Oecologia*, in press
- Young, T.P., N. Patridge, and A. Maccrae. 1995. Long-term glades in acacia bushland and their edge effects in Laikipia, Kenya. *Ecological Applications* 5:97-108.
- Young, T.P., C.H. Stubblefield, and L.A. Isbell. 1997. Coexistence among obligate acacia ants. *Oecologia* 109:98-107.

Appendix 1. The larger (>2kg) mammals of Mpala/Segera. Species marked with an asterisk have been seen at the study plots or (in similar black cotton soils within 10km). Only the largest six carnivores listed are serious predators of the ungulate herbivores. It is difficult to assess the abundance of the smaller nocturnal carnivores. Nomenclature and weights are taken from Dorst and Dandelot (1972), Estes (1991), and Kingdon (1997). Not included are several smaller 'large' mammals (Bushbaby, Black-tipped Mongoose, Dwarf Mongoose, Genet, Zorilla, Striped Ground Squirrel, Bush Squirrel, Spectacled Elephant Shrew, Hedgehog, Nutria); in addition, Gerenuk occur just across the boundary river. Black Rhinoceros have been protected or are being reintroduced on several nearby properties.

English name	Latin name	Biomass(kg)	Abundance
Herbivorous mammals			
*Elephant	Loxodonta africana Blumenbach	1500-6000	Generally low, but seasonally
abundant			
Hippopotamus	<i>Hippopotamus amphibius</i> L.	600-3200	Very few
*Giraffe	<i>Giraffa camelopardalis</i> L.	550-2000	Moderate
[Black Rhinoceros	Diceros bicornis L.	900-1400	Locally extinct]
*Cape Buffalo	Syncerus caffer Sparrman	420-870	Low
*Eland	Taurotragus oryx Pallas	340-700(-900)	Moderate
*Grevy's Zebra	<i>Equus grevyi</i> Oustalet	350-450	Low, increasing
*Burchell's Zebra	Equus burchelli Gray	175-325	Abundant
Greater Kudu	Tragelaphus strepsiceros Pallas	120-320	Low
Defassa Waterbuck	Kobus ellipsiprymnus defassa Ruppell	160-260	Moderate
*Beisa Oryx	Oryx beisa Ruppell	130-200	Low, increasing
*Jackson's Hartebeast	Alcelaphus buselaphus jacksoni Pallas	100-170	Abundant
(*)Warthog	Phacochoerus aethiopicus Pallas	45-100	Very few, increasing?
*Grant's Gazelle	Gazella granti Brooke	40-80	Abundant
Impala	Aepyceros melampus Lichtenstein	40-70	Abundant
Bushbuck	Tragelaphus scriptus Pallas	25-80	Low
Mountain? Reedbuck	Redunca sp. (fulvorufula?)	20-35	Very few
Thomson's Gazelle	Gazella (rufifrons) thomsonii Gunther	13-30	Very few (but common on Segera)
Bush Duiker	Sylvicapra grimmia L.	13-25	Low
Klipspringer	Oreotragus oreotragus Zimmerman	11-14	Low
*Steinbuck	Raphicerus campestris Thunberg	9-13	Moderate
Crested Porcupine	<i>Hystrix</i> sp.	up to 20	Low?
*Anubis Baboon	Papio cynocephalus anubis L.	10-25(-50)	Abundant
*Patas Monkey	Erythrocebus patas Schreber	(4-)7-10	Moderate on Segera
Vervet Monkey	Cercopithecus aethiops L.	(2.5-)5-9	Abundant
Kirk's dik-dik	Madoqua kirki Gunther	4-7	Abundant
Rock hyrax	Procavia capensis Pallas	1.8-5.4	Moderate
Bush Hyrax	Heterohyrax brucei Gray	1.5-2.4	Moderate
*Hare	Lepus sp.	1-4	Abundant
Carnivorous mammals (in	<i>icluding insectivores)</i>		
*Lion	<i>Panthera leo</i> L.	120-260	Low to moderate
*Spotted Hyaena	Crocuta crocuta Erxleben	50-85	Moderate
Leopard	Panthera pardus L.	30-65	Low to moderate
*Cheetah	Acinonyx jubatus Schreber	35-65	Low
(*?)Striped Hyaena	Hyaena hyaena L.	26-45	Low?
Wild Dog	Lycaon pictus Temminck	20-25	Vagrant, virtually extinct
*Aardvark	Oreycteropus afer Pallas	up to 70?	Declining?
(*)Serval	Felis serval Schreber	14-18	Low
Cape Clawless Otter	Aonyx capensis Lonnberg	10-18	Low?
Civet	<i>Civettictis civetta</i>	7-20	Low?
*Ratel	Mellivora capensis Schreber	8-15	Low
(*?)Caracal	Felis aurata Temminck	8-18	Low
Side-striped Jackal	Canis adustus Sundevall	8	Very few

*Black-backed Jackal	Canis mesomelas Schreber	7-14	Moderate, but fluctuates
Golden Jackal	<i>Canis aureus</i> L.	7-15	Very few, on Segera
(*?)Aardwolf	Proteles cristatus Sparrman	8-12	Low?
(*?)African Wild Cat	Felis (sylvestris) lybica Forster	3.2-6.5	Low to moderate?
(*?)Bat-eared Fox	Otocyan megalotis Desmarest	3.2-5.4	Moderate, but fluctuates
*White-tailed mongoose	Ichneumia albicauda G.Cuvier	3-4	Moderate
?Marsh Mongoose	Atilax paludinosus G. Cuvier	2.4-3.2	Present?