Table 1:	Effect	of	$CoCl_2$	(10	mg/kg,	i.p.,	O.D.	for	30	days)	on	body	weight	and	food
consump	tion in di	iffeı	rent gro	up at	week 5.										

Group	Body weight (gm)	Food Consumption (gm/day)
Control group	265.17 ± 3.82	29.21 ± 0.55
Uninephrectomized diabetic group	190.00 ± 3.84 ^{***}	18.26 ± 0.94**
Treatment group	243.17 ± 5.27 [#]	24.89 ± 0.58 [#]
Control group treated with CoCl ₂	264.83 ± 3.79	29.05 ± 0.57

Each value is expressed as mean \pm S.E.M. (n=06). ^{**}p<0.01 versus control, ^{***}p <0.001 versus control group, [#]p<0.05 versus uninephrectomized diabetic group

Group	pD ₂ value	E _{max} (%)
Control group	7.41 ± 0.08	100 ± 8.35
Uninephrectomized diabetic group	8.38 ± 0.08 *, fff	237.96 ± 5.78 ***
Treatment group	7.46 ± 0.17 ^{\$}	127.41 ± 13.37 ^{\$\$}
Control group treated with CoCl ₂	$7.12 \pm 0.05^*$	138.52 ± 13.21

Table 2: Effect of $CoCl_2$ (10 mg/kg, i.p., O.D. for 30 days) on pD₂ values and E_{max} in the thoracic aorta preparation in different group.

Each value is expressed as mean \pm S.E.M. (n=06). *p<0.05 versus control, ***p <0.001 versus control group, *p<0.05 versus uninephrectomized diabetic group, *p<0.01 versus uninephrectomized diabetic group, ^{fff}p< 0.001 versus control group treated with CoCl₂. ***p <0.001 versus control group. Data are expressed as pD2 values which are defined as the negative logarithm to base 10 of the EC50 values. The maximal efficacy of the drug-receptor complex to result in a graded effect is E_{max} on a graded dose-response curve.

Table 3: Effect of CoCl₂ (10 mg/kg, i.p., O.D. for 30 days) on the activities of SOD, CAT and GSH in aorta of control and experimental rats

Groups	SOD activity	Catalase activity	GSH
	(U/mg protein)	(U/mg protein)	(mM/100mg tissue)
Control group	13.72 ± 0.21	72.20 ± 2.26	41.20 ± 0.95
Uninephrectomized diabetic group	4.28 ± 0.20 ***	37.00 ± 1.74 **	21.40 ± 1.12 **
Treatment group	$10.58 \pm 0.39^{\text{fff}}$	$61.00 \pm 2.45^{\pm}$	37.40 ± 1.64 [£]
Control group treated with CoCl ₂	13.22 ± 0.29	71.00 ± 3.46	42.00 ± 1.94

Each value is expressed as mean \pm S.E.M. (n=05). ^{**}p <0.01 versus control group, ^{***}p< 0.001 versus control group, [£]p< 0.05 versus uninephrectomized diabetic group, ^{fff} p< 0.001 versus uninephrectomized diabetic group.