Online Supplemental Table

Table C
Association of Type D with biomarkers of inflammation, coagulation and myocardium

	Non Type D N=3833	Type D N=1095	OR (95% CI)*	р
Inflammation ¹⁾²⁾³⁾				
CRP	1.60 (0.50 / 3.10)	1.60 (0.50 / 3.20)	1.02 (0.94 - 1.11)**	0.64
IL-18	227.8 (181.8 / 289.9)	219.8 (174.9 / 292.7)	0.88 (0.70 - 1.11)**	0.29
IL-1-ra	316.0 (237.4 / 417.3)	319.6 (234.1 / 444.4)	1.13 (0.96 - 1.34)**	0.15
Albumin	43.2 ± 3.1	43.2 ± 3.3	0.99 (0.97 - 1.02)	0.60
White blood cell count	6.8 (5.8 / 8.2)	6.8 (5.8 / 8.1)	1.00 (0.75 - 1.32)**	0.98
Coagulation				
Fibrinogen	346.0 (304.0 / 400.0)	345.0 (301.0 / 401.0)	1.08 (0.77 - 1.50) **	0.67
Myocardium				
Troponin I	0.005 (0.003 / 0.008)	0.005 (0.003 / 0.008)	1.02 (0.97 - 1.07)**	0.37
Troponin I ≥ 0.04 ng/mI, % (N)	1.0 (39)	0.7 (7)	0.72 (0.32 - 1.61)	0.42
NT-proBNP	63.10 (29.22 / 124.90)	57.68 (24.93 / 115.80)	0.98 (0.92 - 1.04)**	0.46

^{*}Odds ratios are adjusted for age and sex

¹⁾ Data presented are mean ± standard deviation and median (25th / 75th) percentile.

²⁾ Subjects with a self reported influenza infection, common cold or other inflammatory diseases during the last week before examination are excluded

³⁾ For the logistic regression models biomarkers were log transformed