

Online Supplemental Table

Table H. Results of subgroup analyses of patients with specific preferences only

	control group	intervention group		
	n = 178	n = 190		
	M (SD)	M (SD)	p	d (95% CI)
Morbidity [#]	4.86 (9.35)	5.48 (9.21)	.514	-0.07 (-0.27-0.14)
Anxiety [§]	6.43 (4.24)	6.15 (4.16)	.515	0.07 (-0.14-0.27)
Depression [§]	5.65 (4.43)	5.64 (4.34)	.987	0.002 (-0.20-0.21)
Positive mood [§]	2.69 (0.68)	2.84 (0.66)	.037	0.22 (0.01-0.42)
Negative mood [§]	2.48 (0.79)	2.38 (0.77)	.252	0.12 (-0.09-0.32)
Satisfaction	3.60 (0.53)	3.57 (0.52)	.538	-0.06 (-0.27-0.14)
Pain	3.35 (2.28)	3.30 (2.23)	.833	0.02 (-0.18-0.23)

Patients who refused interventions or preferred no specific kind of intervention were excluded from the analysis. ANOVA intervention x propensity strata (complex propensity score; including interaction and quadratic terms); CMS computation based on N=847; [#]patients who died included: n = 182 control group, n = 196 intervention group; [§]adjusted for preoperative values