Supplemental tables

		positive SLNB	negative SLNB	
Parameter		n=86 (13.0%)	n=578 (87.0%)	p-value
Gender				
	Male	50 (58.1%)	263 (45.5%)	0.028
	Female	36 (41.9%)	315 (54.5%)	
Age				
	Mean	56.31 +/- 13.9	57.60 +/- 14.9	0.478.
Location				
	Head/Neck	6 (7.1%)	53 (9.7%)	0.541.
	Upper Extremity	14 (16.5%)	98 (17.9%)	
	Trunk	36 (42.4%)	215 (39.2%)	
	Lower Extremity	28 (32.9%)	181 (33.0%)	
	Mucous membrane	1 (1.2%)	1 (0.2%)	
Subtype		· /		
	SSM	34 (40.5%)	206 (39.2%)	0.146.
	LMM	1 (1.2%)	23 (4.4%)	
	NMM	27 (32.1%)	136 (25.9%)	
	ALM	9 (10.7%)	35 (6.7%)	
	Other	13 (15.5%)	125 (23.8%)	
Breslow thickness	outor	10 (10.070)	120 (20.070)	
	< 1mm	6 (7.2%)	213 (37.4%)	0.001
	1.01-2 mm	17 (20.5%)	191 (33.5%)	0.001
	2.01-4.0 mm	34 (41.0%)	105 (18.4%)	
	> 4.01mm	26 (31.3%)	61 (10.7%)	
	mean	3.59 ± 2.39	1.98 ± 2.12	0.001
Clark Level	mean	5.53 ± 2.53	1.90 ± 2.12	0.001
CIAIR LEVEI	I	0 (0%)	2 (0.3%)	0.001
	1	. ,	. ,	0.001
		2 (2.3%)	30 (5.2%)	
		17 (19.8%)	255 (44.4%)	
	IV	52 (60.5%)	256 (44.6%)	
	V	15 (16.3%)	30 (5.2%)	
Ulceration	N			0.004
	No	46 (53.5%)	467 (80.9%)	0.001
. .	Yes	40 (46.5%)	110 (19.1%)	
Regression				
	No	82 (95.3%)	551 (95.5%)	0.952.
	Yes	4 (4.7%)	26 (4.5%)	
_				
Progress	No	51 (60.0%)	511(88.6%)	0.001
	Yes	34 (40.0%)	66 (11.4%)	n= num

Supplemental Table 1: Characteristics of SLN positive and negative patients (n=664)

n= numbers