



ACUTE BEHAVIOURAL DISTURBANCE (ABD) AUDIT FORM

ABD Audit Form instructions:

- This form is to be completed for all patients presenting with **ABD** and **administered sedation** by QAS Paramedics in the Metro North and Metro South LASN.
- A comprehensive patient assessment, including SAT scores must be documented every 5 minutes for the first 20 minutes, then half hourly thereafter and recorded in the table below.
- All completed *ABD Audit Forms* are to be photographed (via QAS issued iPADs) and emailed to <u>ABD.audit@ambulance.qld.gov.au</u> as soon as possible following completion of case.
- The ABD Audit Form does **NOT** replace existing eARF documentation requirements.
- Questions relating to the ABD Audit are to be directed to Lachlan Parker (Lachlan.Parker@ambulance.qld.gov.au)

SEDATION ASSESSMENT TOOL ^[1]											
Score	Responsiveness	Speech									
+3	Combative, violent, out of control	Continual loud outbursts									
+2	Very anxious and agitated	Loud outbursts									
+1	Anxious/restless	Normal/talkative									
0	Awake and calm/cooperative	Speaks normally									
-1	Asleep but rouses if name is called	Slurring or prominent slowing									
-2	Responds to physical stimulation	Few recognisable words									
-3	No response to stimulation	Nil									

Case date:		/	/	Case	number:			Crew ski	II level:	□ ACP2 □	CCP
Patient's initial SAT score:				Patient age:		□ <65 yea	□ <65 years □ ≥65 years		scene:	□ Yes □ No	
Suspected ABD cause:		□ Alcoh	nol intoxication 🗆 Ampheta		etamines 🛭	mines Mental illness		☐ Self harm ☐ Other stimulants			
QAS staff or patient injured?		□ Yes □ No (if Yes, please specify)									
			1		Ī	T	1	1			T
	Time (hh:mm):		:		:	:	:	:	:	:	:
	SAT score (+3 to -3):										
Sedation	Sedative administered:		□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
	Sedative name:										
	Sedative route:		IM /I	V	IM / IV	IM / IV	IM / IV	IM / IV	IM / IV	IM / IV	IM / IV
	Sedative dose:			mg	mg	mg	mg	mg	mį	g mg	mg
	Respiration rate < 12/min:		□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
nts	SpO ₂ <90% (on room air):		□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
Adverse events	Heart rate < 60/min:		□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
	Systolic BP < 90mmHg:		□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
	Airway obstruction requiring	g any	□ Yes □ I	No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No	□ Yes □ No
Ad	airway intervention (e.g. jaw	V									
	support or airway adjunct in	sertion):									

¹ Calver, L. A., Stokes, B., & Isbister, G. K. (2011). Sedation assessment tool to score acute behavioural disturbance in the emergency department. Emerg Med Australas, 23(6), 732-740.