# **Species-Wide Genetic Atlas of Antimicrobial Resistance** in Clostridioides difficile

### **Korakrit Imwattana<sup>1</sup>, César Rodríguez<sup>2</sup>, Thomas V Riley<sup>1,3,4,5</sup> and Daniel R Knight<sup>1,3</sup>**

<sup>1</sup> School of Biomedical Sciences, The University of Western Australia, Western Australia <sup>2</sup> Facultad de Microbiología & Centro de Investigación en Enfermedades Tropicales (CIET), Universidad de Costa Rica, Costa Rica <sup>3</sup> School of Veterinary and Life Sciences, Murdoch University, Western Australia <sup>4</sup> School of Medical and Health Sciences, Edith Cowan University, Western Australia <sup>5</sup> Department of Microbiology, PathWest Laboratory Medicine, Queen Elizabeth II Medical Centre, Western Australia

## **Background and Objectives**

Antimicrobial resistance (AMR) plays an important role in the spread and pathogenesis of Clostridioides difficile infection (CDI). An association between AMR and CDI outbreaks has been identified<sup>1,2,3</sup>, however, such studies have been limited to a few strains in limited geographical regions.



This study aimed to investigate the prevalence of AMR genotypes in the global population of *C. difficile*.

## **Methods**

- 1. All publicly available read files were downloaded.
- 2. All non-redundant illumina read files were screen for quality and contamination using kraken2<sup>4</sup>.
- 3. Read files of epidemic strains (sequence types (STs) 1, 2, 11 and 37) were screened for clonality (nucleotide identity > 99.98%) using SRST2<sup>5</sup> and Sketch<sup>6</sup>, respectively.
- 4. All remaining read files were interrogated against several AMR databases.<sup>#</sup>



## **Results**

### **1. Population structure of C. difficile**









### Conclusion

There was a higher prevalence of AMR among epidemic C. difficile lineages. Despite its intrinsic resistance, some *C. difficile* strains carried aminoglycoside resistance genes, suggesting its role as a reservoir of AMR genes.



### References

Dingle KE, *et al.* (2019). **MBio**;10(2):e02790-18. He M, et al. (2013). Nat Gen; 45(1):109-13. Imwattana K, et al. (2020). Expert Rev Anti Infect Ther;1:17-25. Wood DE, et al. (2019). Genome Biol;20(1):257 Inouye M, et al. (2014). Genome Med;6(11):90. BBtools package [available at https://sourceforge.net/projects/bbmap/] PubMLST database [available at https://pubmlst.org/cdifficile/] Gubta SK, et al. (2014). Antimicrob Agents Chemother;58(1):212-20. Zankari E, et al. (2012). J Antimicrob Chemother; 67(11):2640-4.





## **Re-Defining** *Clostridioides difficile* Using Global Phylogenomic Analyses

Korakrit Imwattana<sup>1</sup>, César Rodríguez<sup>2</sup>, Thomas V Riley<sup>1,3,4,5</sup> and Daniel R Knight<sup>1,3</sup>

- <sup>1</sup> School of Biomedical Sciences, The University of Western Australia, Western Australia
- <sup>2</sup> Facultad de Microbiología & Centro de Investigación en Enfermedades Tropicales (CIET), Universidad de Costa Rica, Costa Rica
- <sup>3</sup> Medical, Molecular and Forensic Sciences, Murdoch University, Western Australia
- <sup>4</sup> School of Medical and Health Sciences, Edith Cowan University, Western Australia
- <sup>5</sup> Department of Microbiology, PathWest Laboratory Medicine, Western Australia

### **Background and Objectives**





Clostridioides difficile species problem: A diverse population classified into a single species based on 16s rRNA Study Objective: Use ANI to re-evaluate the species definition of *C. difficile* 









**1. ANI analysis reveals major** 

discontinuity in *C. difficile* taxonomy

**Total gene repertoire** 17,470 genes

**Core gene** 2,232 genes (12.8%)

#### **Potential specific phenotypes**

- C-I Fructosamine utilisation
- C-II EDTA resistance
- C-III Ethanolamine catabolism
  - Fructosamine utilisation
  - Polyamine biosynthesis

3. Pan-genome analysis identifies clade specific genes and traits





### 2. Bayesian evolutionary analysis reveals cryptic clades are ancient species

### Conclusions

There was a clear species boundary separating *C. difficile* from the 3 novel genomospecies. Several potential phenotypic differences were identified. Difference in toxin gene architecture and its divergence may complicate the diagnosis of *C. difficile* infection.





Large toxin genes
Binary toxin genes
Excisionase genes

### **<u>4. Cryptic clades harbor novel and highly</u> <u>divergent toxin gene architecture identifies</u></u>**

### References

- . Konstantinidis KT, et al. (2006) Philos Trans R Soc Lond B Biol Sci;361:1929-1940.
- Wayne LG, *et al.* (1987) Int J Syst Evol Microbiol;37:463-464.
- 3. Stackebrandt E, *et al.* (1994) **Int J Syst Evol Microbiol**;44:846-849
- 4. Jain C, *et al.* (2018) **Nat Commun**;9:5114.
- 5. Pritchard L, et al. (2016) Anal Methods;8:12-24.
- 6. Didelot X, *et al.* (2018) Nucleic Acids Res;46:e134-e134
- 7. Drummond AJ, et al. (2007) BMC Evol Biol;7:214.
- 8. Tonkin-Hill G, et al. (2020) Genome Biol;21:180.
- 9. Brynildsrud O, *et al.* (2016) Genome Biol;17:238.



