

A cryptic *vanB2* operon carried on a Tn1549-like element in *Clostridium difficile*

Grace O. Androga¹, Daniel R. Knight¹, Susan A. Ballard², Benjamin P. Howden² and Thomas V. Riley^{1,3}

Microbiology & Immunology, School of Pathology and Laboratory Medicine, The University of Western Australia, Nedlands, Western Australia¹; Microbiological Diagnostic Unit Public Health Laboratory, and Doherty Applied Microbial Genomics, Department of Microbiology and Immunology, The Peter Doherty Institute for Infection and Immunity, The University of Melbourne, Parkville, Victoria, Australia²; Department of Microbiology, PathWest Laboratory Medicine, Queen Elizabeth II Medical Centre, Nedlands, Western Australia³.



Summary: The emergence of vancomycin resistance in *Clostridium difficile* would have dramatic clinical consequences. We describe a phenotypically silent *vanB2* operon in a strain of *C. difficile* isolated from an Australian veal calf at slaughter, carried on a ~42k mobile genetic element showing significant homology and synteny to Tn1549, a conjugative transposon linked with the emergence and global dissemination of vancomycin resistant enterococci.

MATERIALS AND METHODS

C. difficile strain (AI0499, PCR RT 033) was recovered from the carcass of a neonatal calf (aged less than 7 days) at an abattoir in Victoria, Australia in 2013 [1].

The genome of strain AI0499 was sequenced in two independent runs using illumina chemistry (Miseq/ HiSeq) and investigated using short-read sequence typing (SRST2), *de novo* assembly and comparative genomic analysis [2-5]. *In vitro* susceptibility to vancomycin was determined using agar and Etest methodologies [6].

RESULTS

De novo assembly of the AI0499 genome revealed a chromosome of 4,095,918 bp and 28.75% GC with a N50 of 58,582 bp and overall coverage of ~130x.

Annotation identified 3960 unique coding sequences (CDS) and 3 predicted intact phage sequences corresponding to ΦMMP02, ΦCD119 and CDMH1.

SRST2 identified 7 vancomycin resistance genes with >99% sequence identity to *vanXB*, *vanB*, *vanHB*, *vanW*, *vanYB*, *vanSB* and *vanRB*. AI0499 was negative for *vanG_{Cd}*.

These genes comprised a *vanB2* operon and were carried on a ~42k mobile genetic element showing significant homology and synteny to Tn1549, a conjugative transposon linked with the emergence and global dissemination of vancomycin resistant enterococci (Fig. 1).

Notably, AI0499 did not show any reduced susceptibility to vancomycin (MIC 1 mg/L).

AI0499 was characterised as ST 11 within MLST clade 5, and having an S-layer cassette type 3. It contained a complete binary toxin locus (CdtLoc) and an uncommon Pathogenicity Locus (PaLoc) identified as toxinotype XI [4] (Fig. 2).

REFERENCES

1. Knight DR, Putsathit P, Elliott B *et al.* Contamination of Australian newborn calf carcasses at slaughter with *C. difficile*. *Clin. Microbiol. Infect.* [in press] doi: 10.1016/j.cmi.2015.11.017.
2. Eyre DW, Tracey L, Elliott B *et al.* Emergence and spread of predominantly community-onset *C. difficile* PCR ribotype 244 infection in Australia, 2010 to 2012. *Euro Surveill.* 2015; 20:21059.
3. Inouye M, Dashnow H, Raven L *et al.* SRST2: Rapid genomic surveillance for public health and hospital microbiology labs. *Genome Med.* 2014; 6:90
4. Rupnik M and Janezic S. An update on *C. difficile* toxinotyping. *J. Clin. Microbiol.* [in press] doi: 10.1128/jcm.02083-15.
5. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. *Bioinformatics* 2011; 27: 1009-10.
6. Knight DR, Giglio S, Huntington PG *et al.* Surveillance for antimicrobial resistance in Australian isolates of *C. difficile*, 2013-14. *J. Antimicrob. Chemother.* 2015; 70:2992-2999.
7. Ammam F, Marvaud JC, Lambert T. Distribution of the *vanG*-like gene cluster in *C. difficile* clinical isolates. *Can J. Microbiol.* 2012; 58: 547-51.
8. Ammam F, Meziane-cherif D, Mengin-Lecreux D *et al.* The functional *vanGCd* cluster of *C. difficile* does not confer vancomycin resistance. *Mol. Microbiol.* 2013; 89: 612-25.
9. Rupnik M. Heterogeneity of large clostridial toxins: importance of *C. difficile* toxinotypes. *FEMS Microbiol. Rev.* 2008; 32:541-55.




Fig. 1 Comparative genomic analysis of Tn1549-like element in *C. difficile* strain AI0499 and Tn1549 of *E. faecalis* (AF192329)
Arrows indicate open reading frames (ORFs) and direction of transcription. Excisionase (*xis*) and integrase (*int*) genes are shown in yellow, genes comprising the *vanB2* operon (*vanXB*, *vanB*, *vanHB*, *vanW*, *vanYB*, *vanSB* and *vanRB*) are shown in light green, with the remaining ORFs shown in light blue. Figure prepared using Easyfig v2.2.2 [5].

DISCUSSION AND CONCLUSION

The absence of phenotypic vancomycin resistance in strain AI0499 is not unexpected. Despite an inducible and functionally active *vanG_{Cd}* operon which is found in the genome of some strains, and significant amounts of non-native D-alanine-D-serine containing peptidoglycan precursors, *C. difficile* strains retain susceptibility to vancomycin [7,8].

It is possible that in strain AI0499 *vanB* mediated resistance is induced by vancomycin but a yet-to-be identified mechanism prevents any modifications to the peptidoglycan structure [8]. Equally, it is likely that the presence of *Bm3R1* and *orf7* within the *vanRB* gene result in a loss of function and subsequent lack of transcription of vancomycin resistance genes. Further studies are underway to investigate these hypotheses.

In conclusion, the isolation of a Tn1549-like element from *C. difficile* is alarming and vancomycin resistance expressed *in vivo* could have dramatic consequences. These data further confirm that anaerobes, particularly those of the animal gut microbiota, represent a reservoir of clinically important *vanB*-like resistance operons.




Fig. 2 Uncommon PaLoc of strain AI0499 compared to toxinotype 0
Strain AI0499 contains a deletion within the PaLoc as shown above; a trait exhibited by toxinotype XI strains [9]. The remaining fragments of the *tcdA* gene are 3231bp and 915bp in size and correspond to the A2 and A3 fragments of the *C. difficile* 630 *tcdA* gene, respectively.