

Emergence of *Clostridium difficile* ribotype 106 in Western Australia

Korakrit Imwattana¹ , Daniel R Knight² , Larry K Kociolek^{3,4} , Egon A Ozer⁵ , David W Eyre^{6,7} , Thomas V Riley^{1,2,8,9}

1. School of Biomedical Sciences, The University of Western Australia, WA, Australia

2. School of Veterinary and Life Sciences, Murdoch University, WA, Australia

3. Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, USA

4. Division of Infectious Diseases, Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, USA

5. Department of Medicine, Division of Infectious Diseases, Northwestern University Feinberg School of Medicine, Chicago, USA

6. Nuffield Department of Clinical Medicine, John Radcliffe Hospital, University of Oxford, Oxford, UK

7. National Institute for Health Research (NIHR) Oxford Biomedical Research Centre, John Radcliffe Hospital, Oxford, UK

8. School of Medical and Health Sciences, Edith Cowan University, WA, Australia

9. Department of Microbiology, PathWest Laboratory Medicine, WA, Australia

Background & Objectives

Clostridium difficile is a common cause of antibiotic-associated diarrhoea, which is mediated by one or more of three toxins: toxin A (TcdA), toxin B (TcdB) and binary toxin (CDT).^{1,2} *C. difficile* can be classified into different ribotypes (RTs). Recently, there has been a change in the epidemiology of *C. difficile* infection (CDI) in the USA, as *C. difficile* RT 106 has become the most common cause of both community-acquired and hospital-associated CDI supplanting RT 027.³ In WA, there were no reports of RT 106 before September 2015, when the first strain was isolated. Since then, there has been a steady increase in the prevalence of RT 106 in WA (Figure 1).

The objectives of this study were to investigate the increased prevalence of *C. difficile* RT 106 in WA and to evaluate the screening tool used in the outbreak investigation of CDI.

Methods

- All *C. difficile* strains that had been reported as RT106 were sent to the reference laboratory where PCR ribotyping and toxin gene detection were performed as previously described.⁴⁻⁷
- Eleven *C. difficile* strains from 5 different hospitals in WA (Figure 2) were selected for further whole genome sequence analysis.
- In silico* multi-locus sequence typing (MLST) was performed on 11 *C. difficile* strains and core genome single nucleotide polymorphisms (CGSNPs) analysis was performed on 11 *C. difficile* strains. For comparative analysis, we included genomes of RT 106 circulating in the United Kingdom (UK)⁸ and the USA.⁹





Fig 2 – Allele differences between STs 28 and 42. ST 28 differs from ST 42 by two different SNPs in a single allele.

Fig 3 – Phylogenetic tree of *C. difficile* ST 28 (A) and ST 42 (B) from CGSNPs analysis. Eleven WA isolates are displayed in blue lettering (PWs). The coloured squares correspond to individual wards in different hospitals. SRR and LK strains were from USA and ERR strains were from UK. Asterisks (* and **) in the red boxes indicate two clonal groups of *C. difficile*.



Symbol	Name	Details
*	Clonal group I	A group of 2 <i>C. difficile</i> isolates from 2 patients from a single ward in a hospital. The 2 strains were separated by 55 days.
**	Clonal group II	A group of 3 <i>C. difficile</i> isolates from 3 patients from 3 different hospitals. PW029 and PW033 were separated by 65 days. PW033 and PW043 were separated by 98 days.




Fig 1 – Number of *C. difficile* RT 106 isolated in WA. The first and only strain in 2015 was isolated in September. Nine, twelve and sixteen strains were isolated in 2016, 2017 and 2018, respectively.

Results

- Six RT 106 strains belonged to ST 28 and five belonged to ST 42. These two STs are single-locus variants of one another (Figure 2).
- The phylogenetic tree of ST 28 (Figure 3A) shows that ST 28 in WA formed a single cluster which was distinct from strains from both the UK and USA.
- Two ST 28 strains from two patients in the same ward (clonal group I in Figure 3A) were identical (CGSNP = 0).
- The phylogenetic tree for ST 42 (Figure 3B) showed that ST 42 in WA did not form a single cluster.
- Three ST 42 strains from three patients in different hospitals (clonal group II in Figure 3B) were identical (CGSNP = 0).

Discussion and Conclusions

- This study demonstrates two closely related lineages of *C. difficile* RT 106 (STs 28 and 42).
- Two clonal groups of *C. difficile* RT 106 were identified in this study. Clonal group I demonstrates a possible source of transmission within a hospital. Clonal group II demonstrates a possible source of transmission in the community.
- C. difficile* RT 106 strains in WA were distinct from RT 106 strains found in other countries.
- PCR ribotyping can be used as a screening test for outbreak investigations, but confirmation testing (e.g. MLST or core genome typing) is also needed.

References

- Chumbler NM, Rutherford SA, Zhang Z, Farrow MA, Lisher JP, Farquhar E, Giedroc DP, Spiller BW, Melnyk RA, Lacy DB. 2016. Crystal structure of *Clostridium difficile* toxin A. Nat Microbiol 1:15002.
- Bauer MP, Notermans DW, van Benthem BH, Brazier JS, Wilcox MH, Rupnik M, Monnet DL, van Dissel JT, Kuijper EJ, ECDIS Study Group. 2011. *Clostridium difficile* infection in Europe: a hospital-based survey. Lancet 377:63-73.
- Kociolek LK, Gerding DN, Hecht DW, Ozer EA. 2018. Comparative genomics analysis of *Clostridium difficile* epidemic strain DH/NAP11/106. Microbes Infect 20:245-253.
- Kato H, Kato N, Watanabe K, Iwai N, Nakamura H, Yamamoto T, Suzuki K, Kim SM, Chong Y, Wasito EB. 1998. Identification of toxin A-negative, toxin B-positive *Clostridium difficile* by PCR. J Clin Microbiol 36:2178-82.
- Stubbs SL, Brazier JS, O'Neill GL, Duerden BI. 1999. PCR targeted to the 16S-23S rRNA gene intergenic spacer region of *Clostridium difficile* and construction of a library consisting of 116 different PCR ribotypes. J Clin Microbiol 37:461-3.
- Kato N, Ou CY, Kato H, Bartley SL, Brown VK, Dowell VR, Jr., Ueno K. 1991. Identification of toxigenic *Clostridium difficile* by the polymerase chain reaction. J Clin Microbiol 29:33-7.
- Stubbs S, Rupnik M, Gibert M, Brazier J, Duerden B, Popoff M. 2000. Production of actin-specific ADP-ribosyltransferase (binary toxin) by strains of *Clostridium difficile*. FEMS Microbiol Lett 186:307-12.
- Eyre DW, Cule ML, Wilson DJ, Griffiths D, Vaughan A, O'Connor L, Ip CLC, Golubchik T, Batty EM, Finney JM, Wyllie DH, Didelot X, Piazza P, Bowden R, Dingle KE, Harding RM, Crook DW, Wilcox MH, Peto TEA, Walker AS. 2013. Diverse sources of *C. difficile* infection identified on whole-genome sequencing. N Engl J Med 369:1195-205.
- Stone NE, Sidak-Loftis LC, Sahl JW, Vazquez AJ, Wiggins KB, Gillette JD, Hicks ND, Schupp JM, Busch JD, Keim P, Wagner DM. 2016. More than 50% of *Clostridium difficile* Isolates from Pet Dogs in Flagstaff, USA, Carry Toxigenic Genotypes. PLoS One 11:e0164504.

