# Higuera, P.E. and P.V. Dunnette. 2014. Data, code, and figures from Dunnette et al. 2014, Biogeochemical impacts of wildfires over four millennia in a Rocky Mountain subalpine watershed. *New Phytologist* In Press (doi: 10.1111/nph.12828). figShare:

http://dx.doi.org/10.6084/m9.figshare.988687

## Overview

This archive includes data and scripts needed to reproduce the analyses in Dunnette et al. 2014; it includes 49 files and is 23 MB when downloaded as a .zip archive. After extracting the .zip archive, sort by file type. Most data files are in .csv format, while some are in .xls format. Scripts and functions are written for MATLAB software (\*.m file type; www.mathworks.com), and each is file fully commented, including dependencies. For some scripts or functions, the MATLAB statistics or curve fitting toolbox is required; all other functions required are provided in this archive.

**Contact:** Philip Higuera, University of Idaho, PaleoEcology and Fire Ecology Lab, phiguera@uidaho.edu

#### **Citation Information**

Original Reference - please cite if you use data, code, or figures in your own work

Dunnette, P.V., P.E. Higuera, K.K. McLauchlan, K.M. Derr, C.E. Briles, and M.H. Keefe. 2014. Biogeochemical impacts of wildfires over four millennia in a Rocky Mountain subalpine watershed. *New Phytologist* In Press doi: 10.1111/nph.12828

\*The definitive version of this paper is available at http://www.newphytologist.com

**Online Resources** 

Higuera, P.E. and P.V. Dunnette. 2014. Data, code, and figures from Dunnette et al. 2014, Biogeochemical impacts of wildfires over four millennia in subalpine watershed. *New Phytologist* In Press (doi: 10.1111/n] http://dx.doi.org/10.6084/m9.figshare.988687

Also available from:

ŧ

#### 4/21/2014

#### README.md

- github: https://github.com/HigueraLab/Dunnette\_et\_al\_2014\_NewPhytologist/
- University of Idaho PaleoEcology and Fire Ecology Lab: http://www.uidaho.edu/cnr/paleoecologylab

#### Contents

- 1. Chickaree Lake Biogeochemical Data
- 2. Chickaree Lake Biogenic Silica Data
- 3. Chicakree Lake Loss-on-ignition Data
- 4. Chickaree Lake Charcoal Data
- 5. Chickaree Lake Magnetic Susceptibility Data
- 6. Chickaree Lake Chronology Data
- 7. Chickaree Lake Pollen Data
- 8. MATLAB Code to generate figures
- 9. Figures

# 1. Chickaree Lake Biogeochemical Data

#### File: CH10\_biogeochemData.csv

#### Includes the following raw data (by column):

- 1. core\_ID: Sediment core identifier
- 2. drive\_ID: Drive identifier
- 3. top\_sam #: Top sample number
- 4. bot\_sam #: Bottom sample number
- 5. top\_cm: Top depth of sample (cm)
- 6. bot\_cm: Bottom depth of sample (cm)
- 7. top\_age: Top age of sample (cal. yr before CE 1950)
- 8. bot\_age: Bottom age of sample (cal. yr before CE 1950)
- 9. d15N: Nitrogen isotopic composition (delta 15N; per mil)
- 10. %N: Percent Nitrogen (by weight)
- 11. d13C: Carbon isotopic composition (delta 13C; per mil)
- 12. %C: Percent Carbon (by weight)
- 13. C:N\_atomic\_ratio: Ratio of %C to %N (atomic)
- 14. bulk\_density: Bulk density (dry g wet cm-3)
- 15. C\_acc: Carbon accumulation rate (g cm-2 yr-1)

16. MS\_SI : Magnetic susceptibility (SI units)

#### **Missing Values: NaN**

#### **Checksum values:**

- 635 rows (with headers), 16 columns
- Column 3 (top\_sam #): 58551
- Column 5 (top\_cm): 201836
- Column 10 (%N): 626.83
- Column 15 (C\_acc): 341.58

# 2. Chickaree Lake Biogenic Silica Data

#### File: CH10\_BSiData.csv

#### Includes the following raw data (by column):

- 1. top\_cm: top depth of sample (cm)
- 2. bot\_cm: bottom depth of sample (cm)
- 3. top\_age: top age of sample (cal. yr before CE 1950)
- 4. bot\_age: bottom age of sample (cal. yr before CE 1950)
- 5. %BSi: Percent biogenic silica (by weight)
- 6. d15NAIR: Nitrogen isotopic composition (delta 15N; per mil)
- 7. %N: Percent Nitrogen (by weight)
- 8. d13CVPDB\_17O\_corrected: Carbon isotopic composition delta 13C; per mil)
- 9. %C: Percent Carbon (by weight)
- 10. C:N\_atomic\_ratio: Ratio of %C to %N (atomic)
- 11. bulk\_density: Bulk density (dry g / wet cm3)

#### **Missing Values: None**

#### **Checksum values:**

- 41 rows (with headers), 11 columns
- Column 3 (top\_cm): 69935
- Column 5 (%BSi): 1042.45
- Column 10 (C:N): 591.52

# 3. Chickaree Lake Loss-on-ignition Data

#### File: CH10\_LOI\_Data.csv

#### Includes the following raw data (by column):

- 1. core\_ID: Sediment core identification number
- 2. drive\_ID: Drive identifier
- 3. top\_sam #: Top sample number
- 4. bot\_sam #: Bottom sample number
- 5. top\_cm: top depth of sample (cm)
- 6. bot\_cm: bottom depth of sample (cm)
- 7. top\_age: top age of sample (cal. yr before CE 1950)
- 8. bot\_age: bottom age of sample (cal. yr before CE 1950)
- 9. d15N: Nitrogen isotopic composition (delta 15N; per mil)
- 10. %N: Percent nitrogen (by weight)
- 11. d13C: Carbon isotopic composition (delta 13C; per mil)
- 12. %C: Percent carbon (by weight)
- 13. C:N\_atomic\_ratio: Ratio of %C to %N (atomic)
- 14. bulk\_density: Bulk density (dry g wet cm-3)
- 15. LOI\_550: Loss on ignition at 550 C (% organic matter; multiply by 100)
- 16. LOI\_1000: Loss on ignition at 1000 C (% organic matter; multiply by 100)

#### **Missing Values: None**

#### **Checksum values:**

- 124 rows (with headers), 16 columns
- Column 3 (top\_samp #): 12010
- Column 5 (top\_cm): 31833.7
- Column 10 (%N): 161.05
- Column 15 (LOI\_550): 38.63

### 4. Chickaree Lake Charcoal Data

# Files: CH10\_charData.csv, CH10\_charParams.csv, CH10\_charResults.csv

#### README.md

Three files provide the raw input data, the parameters used, and the output data for charcoal analysis via the program *CharAnalysis* (see *Materials and Methods* in main text, and web site https://sites.google.com/site/charanalysis/).

#### 4.1 CH10\_charData.csv

#### Includes the following raw data (by column):

- 1. cmTop: top depth (cm) of the sample
- 2. cmBot: bottom depth (cm) of the sample
- 3. ageTop: estimated age at top of sample (cal. yr before CE 1950)
- 4. ageBot: estimated age at bottom of sample (cal. yr before CE 1950)
- 5. charVol: volume of sediment subsample from which charcoal was prepared (cm3)
- 6. charCount: pieces of charcoal counted in the sample (#)

#### **Missing values: None**

#### **Checksum values:**

- 1202 rows (with headers), 6 columns
- Column 3 (ageTop): 2446519
- Column 6 (charCount): 38218

#### 4.2 CH10\_charParams.csv

See CharAnalysis User's Guide for description of parameters file, available at the web site linked to above.

#### Missing values: -9999 (column 3) or blank cell (all others)

#### **Checksum values:**

- 26 rows (with headers), 5 columns
- Column 3 (Parameters): -39779

#### 4.3 CH10\_charResults.csv

#### Includes the following derived data (reflecting interpolation):

- 1. cmTop\_i: top depth (cm) of interpolated sample
- 2. ageTop\_i: bottom depth (cm) of interpolated sample
- 3. charCount\_i: pieces of charcoal in interpolated sample
- 4. charVol\_i: volume of interpolated sample

- 5. charCon\_i: charcoal concentration in interpolated sample (pieces/cm3)
- 6. charAcc\_i: charcoal accumulation rate, based in interpolated concentration and age (pieces/cm2 yr)
- 7. charBkg: background charcoal, Cback, smoothed based on methods selected in \_\_charParams.csv file, (pieces/cm2 yr)
- 8. charPeak: peak charcoal, Cpeak, based on methods selected in \_charParams.csv file, (pieces/cm2\*yr)
- 9. thresh1: threshold value (pieces/cm2 yr) based on first threshold entered in \_\_charParams.csv file
- 10. thresh2: same as thresh1, but for second threshold entered
- 11. thresh3: same as thresh1, but for third threshold entered
- 12. threshFinalPos: positive threshold value (pieces/cm2 yr), based on fourth threshold value entered in \*charParams.csv file
- 13. threshFinalNeg: negative threshold value (pieces/cm2 yr), based on fourth threshold value entered in \*charParams.csv file
- 14. SNI: signal-to-noise index values, based on threshdFinalPos values
- 15. threshGOF: P value from KS goodness-of-fit test between fitted noise distribution and empirical data below the sample-specific threshold
- 16. peaks1: samples that exceed thresh1 values are identified by "1"; only the first sample is identified
- 17. peaks2: same as peaks1, but for the second threshold value
- 18. peaks3: same as peaks1, but for the third threshold value
- 19. peaksFinal: same as peaks1, but for the final threshold value
- 20. peaksInsig.: peaks that exceeded threshFinalPos (the final threshold), but did not pass the minimum-count test are identified with a "1"
- 21. peakMag: peak magnitude is the total pieces of charcoal accumulated in a given peak (pieces/cm2\*peak); if a peak is only one sample long, then peak magnitude is simply CHAR minus the positive threshold. If a peak is more than one sample long, then each sample exceeding threshFinal is summed
- 22. smPeak Frequ: frequency of peaks (from peaksFinal) smoothed over time, as set in \*\_charParams.csv file
- 23. smFRIs: fire return intervals (from peaksFinal) smoothed over time, as set in \*\_charParams.csv file

#### Missing values: NaN

#### **Checksum values:**

• 458 rows (with headers), 23 columns

- Column 2 (age Top\_i): 1014540
- Column 6 (char Acc\_i): 753.8938
- Column 19 (peaks Final): 36

# 5. Chickaree Lake Magnetic Susceptibility Data

# Files:CH10\_MS\_charParams.csv,CH10\_MS\_charData.csv,CH10\_MS\_charResults.csv

Three files provide the raw input data, the parameters used, and the output data for peak analysis of magnetic susceptibility (MS) data, using the program *CharAnalysis* (see *Materials and Methods* in main text, and web site

https://sites.google.com/site/charanalysis/). The column headers for all files are the same as for charcoal analysis (CH10\_char\* files), but NOTE that the units for MS have been manipulated to be able to be used in CharAnalysis (see 7-9 below).

#### 5.1 CH10\_MC\_charData.csv

#### Includes the following raw data (by column):

- 1. cmTop: top depth (cm) of the sample
- 2. cmBot: bottom depth (cm) of the sample
- 3. ageTop: estimated age at top of sample (cal. yr before CE 1950)
- 4. ageBot: estimated age at bottom of sample (cal. yr before CE 1950)
- 5. dummyVar: dummy variable set to 1, to facilitate use in CharAnalysis program
- 6. MS\_shifted\_trans: Shifted MS data (from col. 8), transformed by dividing by the sediment accumulation rate (cm/yr). Thus, when these transformed values are multiplied by the sediment accumulation rate (cm/yr), as done in CharAnalysis, the result is the transformed MS values (column 8).
- 7. MS\_raw\_SI: Raw magnetic susceptibility measurements (SI units)
- 8. MS\_shifted: Raw MS values, shifted by adding the minimum values in MS\_raw\_SI to each value, such that the minimum value becomes 0. CharAnalysis cannot work with negative data.

#### **Missing values: None**

#### **Checksum values:**

- 1477 rows (with headers), 8 columns
- Column 3 (ageTop): 3970165
- Column 6 (setRate/MS): 0.3626761

#### README.md

# 6. Chickaree Lake Chronology Data

#### Files: CH10\_210Pb\_data.xls, CH10\_ageDepthData.xls, CH10\_ageDepthData.csv, CH10\_radiometricSamples.csv, CH10\_14Cdates.zip

These files provide the chronology data presented in Table S1 (CH10*radiometricSamples.csv), the age-depth data presented in Figure 2* (*CH10\_ageDepthData.csv), and the input files used in the MATLAB functions \_CRSModel* and *MCAgeDepth* (\*.xls), publically available at http://code.google.com/p/crsmodel/ and http://code.google.com/p/mcagedepth/, as used in Higuera et al. (2009, Ecological Monographs, 79: 201-219). See the user manuals for each program at these web sites for details on the two .xls files. The .zip archive includes the .Boo` files output from CALIB (see *Materials and Methods* in main text) and needed to run *MCAgeDepth*. The age-depth data output from *MCAgeDepth* is described below.

#### 6.1 CH10\_ageDepthData.csv

Includes the following raw data (by column):

- 1. Top depth of sample (cm)
- 2. Calibrated age at sample top (cal yr BP)
- 3. Upper 95% confidence intervals for data in column two
- 4. Lower 95% confidence intervals for data in column two
- 5. Sedimentation rate (cm/yr)
- 6. Upper 95% confidence intervals for data in column five
- 7. Lower 95% confidence intervals for data in column five
- 8. Sample resolution (yr/sample)
- 9. Upper 95% confidence intervals for data in column eight
- 10. Lower 95% confidence intervals for data in column eight

#### **Missing values: None**

#### **Checksum values:**

- 1596 rows (with headers), 10 columns
- Column 1 (sampleCm): 642150.9
- Column 2 (calAge): 4679522
- Column 5 (sedAcc): 211.236
- Column 8 (sampleRes): 6860.85

# 7. Chickaree Lake Pollen Data

#### Files: CH10\_PollenCounts.xls, CH10\_pollenPercentages.csv

Raw pollen data needed to use in the function pollenDiagram\_ROMO.m, to create Fig. S2 (*.xls*), *and pollen percentages* (.csv).

#### 7.1 CH10\_PollenCounts.xls

Includes the following raw data (by column):

- 1. Sample\_ID : Identification of sample, by lake (CH), year (07 or 10), core (1 or 2), and drive (A, B,C,...).
- 2. Sample\_number: Sample number for each drive
- 3. Top\_cm: Top depth of each sample (cm)
- 4. age\_yrBP: Age of each sample, in calibrated years before present (CE 1950)
- 5. Pinus hap: Haploxilon Pinus pollen grains counted
- 6. Pinud dip: Diploxilon Pinus pollen grains counted
- 7. Pinus undiff: Undifferentiated Pinus pollen grains counted
- 8. Picea: Picea pollen grains counted 9...112: Column heads are the taxonomic identification for each pollen grain counted
- 9. Charcoal: NOT COUNTED
- 10. EU: Exotic pollen grains, added as spike.

#### Missing values: NaN.

#### **Checksum values:**

- 46 rows (with headers), 113 columns
- Column 2 (sample\_number): 6143
- Column 4 (age\_yrBP): 86683
- Column 10 (Pseudotsuga/Larix): 42
- Column 113 (EU): 8793

#### 7.2 CH10\_pollenpercentages.csv

Includes the following raw data (by column):

- 1. Sample\_ID : Identification of sample, by lake (CH), year (07 or 10), core (1 or 2), and drive (A, B, C,...).
- 2. Sample\_number: Sample number for each drive

- 3. Top\_cm: Top depth of each sample (cm)
- 4. age\_yrBP: Age of each sample, in calibrated years before present (CE 1950)
- 5. Pinus hap: Haploxilon Pinus pollen grains counted
- 6. Pinud dip: Diploxilon Pinus pollen grains counted
- 7. Pinus undiff: Undifferentiated Pinus pollen grains counted
- 8. Picea: Picea pollen grains counted 9...107: Columns heads are the taxonomic identification for each pollen grain counted

#### Missing values: None.

#### **Checksum values:**

- 46 rows (with headers), 106 columns
- Column 2 (sample\_number): 6143
- Column 4 (age\_yrBP): 86683
- Column 10 (Pseudotsuga/Larix): 12
- Column 102 (Nuphar): 13.19

# 8. MATLAB code to generate figures in Dunnette et al. 2014

Files: All are MATLAB files; figure scripts inclues "\_scripts" while all other files are functions called upon by the scripts. To run the script files, the function files must be in the working directory. See individual files for readme information, including dependencies.

- auto.m
- corr\_p\_adj\_N.m
- Fig\_3\_script.m
- Fig\_4\_5\_script.m
- Fig\_6\_S4\_script.m
- Fig\_S1\_script.m
- Fig\_S2\_script.m
- pollenDiagram\_ROMO.m
- SEA.m
- SEA\_CI.m

# 9. Figure files from Dunnette et al. 2014

Figures come in two formats .tif and .fig. The latter is the MATLAB figure file format. *NOTE:* There is no .fig file for figure 1.

#### Files:

- Fig\_1.jpg
- Fig\_2.jpg
- Fig\_3.jpg
- Fig\_4.jpg
- Fig\_5.jpg
- Fig\_6.jpg
- Fig\_S1.jpg
- Fig\_S2.jpg
- Fig\_S3.jpg
- Fig\_S4.jpg
- Fig\_2\_800\_dpi.fig
- Fig\_3\_800\_dpi.fig
- Fig\_4\_800\_dpi.fig
- Fig\_5\_800\_dpi.fig
- Fig\_6\_800\_dpi.fig
- Fig\_S1.fig
- Fig\_S2.fig
- Fig\_S3.fig
- Fig\_S4.fig