


## Supporting Information

# **pH/Reduction dual-responsive oxidized alginate-doxorubicin (mPEG-OAL-DOX/Cys) prodrug nanohydrogels: Effect of complexation with cyclodextrins**


*Tingting Zhou, Jiagen Li, Xu Jia, Xubo Zhao, and Peng Liu\**

State Key Laboratory of Applied Organic Chemistry and Key Laboratory of  
Nonferrous Metal Chemistry and Resources Utilization of Gansu Province, College of  
Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000, China.







\* Corresponding Author. Tel./Fax: 86 0931 8912582. Email: [pliu@lzu.edu.cn](mailto:pliu@lzu.edu.cn).




**Figure S1.** FTIR spectrum of  $\alpha$ -CD obtained with the KBr pellet technique.










**Figure S1.** The plotted curves: zero-order model mechanism of mPEG-OAL-DOX/Cys(ND) (a); first-order model mechanism of mPEG-OAL-DOX/Cys(ND) (b); zero-order model mechanism of mPEG(ND)-OAL-DOX/Cys (c); and first-order model mechanism of mPEG(ND)-OAL-DOX/Cys (d).





**Figure S3.** The plotted curves of the Higuchi and Korsmeyer-Peppas drug release models of the mPEG-OAL-DOX/Cys(CD) prodrug nanohydrogels.





**Figure S4.** The plotted curves of the Higuchi and Korsmeyer-Peppas drug release models of the mPEG(CD)-OAL-DOX/Cys prodrug nanohydrogels.