




**United States Department of Interior
Bureau of Ocean Energy Management**




Biostratigraphic Chart of the Gulf of Mexico Offshore Region, Jurassic to Quaternary



BOEM Gulf of Mexico Resource Evaluation
Robert D. Witrock (504) 736-2718 or 1-800-200-GULF

For public release May 1, 2003



LEGEND	
All foraminiferal markers are extinction points (LADs) unless otherwise indicated.	
(O) indicates an ostracode species.	
The calcareous nanoplanktonic markers are mainly coccolith species LADs. Dinoflagellates and palynomorphs (pollen) are occasionally used.	
LAD = Last Appearance Datum	
FAD = First Appearance Datum	
Species LADs separated by a virgule (the mark "/") indicates either synonymous species or synchronous species.	
Species LADs separated by "=" indicates species that are not synonymous.	
Two or more species LADs separated by commas indicate synchronous species LADs.	
(for example, Discaster bellii, Catmister calyculus are two separate but time-equivalent coccolith extinction points).	
BOEM Chronzone: A body of rock formed during the same time span, bounded by biostratigraphic or correlative seismic markers.	

- References used for Mesozoic and Cenozoic age determination and stratigraphic units**
- Ammerman, J.M., 1989, High-resolution sequence biostratigraphy: examples from the Gulf of Mexico Plio-Pleistocene, in J.A. Howell and F.J. Aitken, eds., High-resolution sequence stratigraphy: innovations and applications: Geological Society Special Publication 104, p. 65-86.
- Ascoli, P., 1984, Episubtidial biostratigraphy across the Jurassic-Cretaceous boundary on the northwestern Atlantic shelf: Benito 83, International Symposium on Benthic Foraminifera, Pau, v. 23-34.
- Ascoli, P., C.W. Poag, and J. Bernier, 1984, Microfacial zonation across the Jurassic-Cretaceous boundary on the Atlantic margin of North America, in G.E. Weissenberg, ed., Jurassic-Cretaceous biostratigraphy and paleogeography of North America: Geological Association of Canada Special Paper 27, p. 31-111.
- Bergen, J.A., and R. Skora, 1984, Molluscan distribution in southern Norwegian Sea chalk: Valhall and Snorre fields, in R. Skora and D. Simone, eds., Biostratigraphy in production and development geology: Geological Society Special Publication 152, p. 85-111.
- Bergen, W.A., D. V. Kent, C. C. Swisher III, and M.-P. Aubry, 1986, Plankton stratigraphy, volume 1: planktic foraminiferal calcareous nanofossils and calcipeloids: Cambridge University Press, 509 p.
- Botti, H.M., J. B. Saunders, and J. Perch-Nielsen, 1984, Benthic foraminiferal biostratigraphy of the south Caribbean region: Cambridge University Press, 408 p.
- Braunholtz, T.J., R.M. Lester, V. Star, and H.R. Thiet, 1995, An integrated Cretaceous microfacies biostratigraphy: a comparison of planktonic foraminifera and nanofossils, parts I and II: Oligocene through Pleistocene foraminifera: Gulf Coast Society of Economic Paleontologists and Mineralogists Special Publication 54, p. 65-79.
- Bread, S.O., A.D. CalleNDER, and M. Naul, 1997, Application of bathyal foraminiferal events to the Neogene through Plio-Quaternary: biostratigraphic framework of the Gulf of Mexico basin: Gulf Coast Association of Geological Societies Transactions, v. 47, p. 111-116.
- Bread, S.O., A.D. CalleNDER, and M. Naul, 2000, Gulf of Mexico basin Late Tertiary deep-water biostratigraphic zonation: relationship to standard shelf foraminiferal and calcareous nanofossil marker terminology: Gulf Coast Section Society of Economic Paleontologists and Mineralogists Foundation 17th Annual Research Conference Deep-water Reservoirs of the World, December 3-6, 2000, Houston.
- Brenchley, P.J., E. Di Stefano, A. Sulli, P.F. Vitale, S. Infuso, and P.R. Vail, 1998, Sequences and system tracts calibrated by high-resolution biostratigraphy of the central Mediterranean Plio-Pleistocene system, in P.-C. de Graciansky, J. Hardenbol, T. Jacquin, and P.R. Vail, eds., Mesozoic and Cenozoic sequence stratigraphy of Europe's basins: Society of Economic Paleontologists and Mineralogists Special Publication 61, p. 155-177.
- Fritzell, D.L., 1954, Handbook of Cretaceous foraminifera of Texas: Bureau of Economic Geology Report of Investigations 22, 23 p.
- Gradstein, F.M., 1976, Biostratigraphy and biogeography of Jurassic Grand Banks foraminifera: Maritime Sediments Special Publication 18, p. 557-583.
- Gradstein, F.M., F.P. Agterberg, and M.A. Torino, 1990, Time in quantitative stratigraphy, in T.A. Cross, ed., Quantitative dynamic stratigraphy, p. 519-542.
- Gradstein, F.M., F.P. Agterberg, J.G. Ogg, J. Hardenbol, P. Veen, J. Thierry, and Z. Huang, 1995, A Triassic, Jurassic and Cretaceous time scale, in W.A. Berggren, D.V. Kent, M.-P. Aubry, and J. Hardenbol, eds., Geochronology, time scales and global stratigraphic correlation: Society of Economic Paleontologists and Mineralogists Special Publication 54, p. 95-126.
- Jones, R.W., 1998, Microfacial biostratigraphy in petroleum exploration: Oxford, Oxford University Press, 432 p.
- Jones, R.W., and M.D. Simmons, eds., 1999, Biostratigraphy in production and development geology: Geological Society Special Publication 152, p. 318-319.
- Lord, A.R., ed., 1982, A stratigraphic index of calcareous nanofossils: Chichester, British Micropalaeontological Society, 192 p.
- Poag, C.W., Jr., B.F. Perkins, N.C. Boers, and J.J. Naar, eds., 1999, Gulf of Mexico basin biostratigraphic index: intercalibration of planktonic foraminifera and nanofossils, parts I and II: Oligocene through Pleistocene foraminifera: Gulf Coast Society of Economic Paleontologists and Mineralogists Foundation, 215 p., 3 charts, 2 tables of synforms (Text and CD-ROM).
- Poehl, D., 2001, Magnetostratigraphic tests of sequence stratigraphic correlations from the southern California Paleogene, Journal of Sedimentary Research, v. 71, p. 528-536.
- Simmons, M.D., W.H. Bergeron, R.O. Keay, R.W. Scott, and W. Ziegler, 1997, Biostratigraphy and geochemistry in the 21st century: Paleontology in the 21st century, an International Senckenberg Conference and Workshop, reports and recommendations, Senckenberg Conference, 215 p.
- Stainforth, R.M., J.L. Lamb, H. Lutzebacher, J.H. Beard, and R.M. Jeffords, 1975, Cenozoic planktonic foraminiferal zonation and characteristic index forms: University of Kansas Paleontological Institute, Article 62, 162 p.
- Styan, M.J., 1996, Late Cenozoic biostratigraphy of the Gulf of Mexico: Gulf Coast Society of Economic Paleontologists and Mineralogists Foundation, chart, 2 sheets.
- Total Biostratigraphic Services, 1994, Miocene/Oligocene biostratigraphic chart (Gulf of Mexico): Total Biostratigraphic Services, Inc., New Orleans.
- Wateman, A.S., 1999, Biostratigraphic chart, Gulf of Mexico Neogene (version 9909): Paleo Data, Inc., New Orleans.
- Wood, L.J., 2000, Chronostratigraphy and lithostratigraphy of the Columbus basin, eastern offshore Trinidad: American Association of Petroleum Geologists Bulletin, v. 84, p. 1905-1928.