

**Supplemental Table 1: Unadjusted and adjusted risk for readmission for all patients with preeclampsia restricted to hospitalizations with deliveries occurring within first two days of hospitalization**

		Unadjusted RR, 95% CI	Adjusted RR, 95% CI	
<b>Postoperative length of stay</b>				
Length of stay <3 days		Reference	Reference	
Length of stay 3 days	1.02	0.78, 1.35	0.92	0.76, 1.11
Length of stay 4 days	0.92	0.68, 1.25	0.75	0.61, 0.94
Length of stay 5 to 7 days	0.62	0.41, 0.95	0.46	0.34, 0.62
Length of stay > days	0.60	0.24, 1.48	0.39	0.19, 0.79
<b>Preeclampsia diagnosis</b>				
Mild		Reference	Reference	
Severe	1.28	1.03, 1.60	1.37	1.17, 1.60
Superimposed	1.44	1.10, 1.90	1.37	1.12, 1.66
<b>Age</b>				
15-19 years old	0.76	0.45, 1.29	0.91	0.66, 1.25
20-24 years old	0.79	0.57, 1.1	0.81	0.65, 1.01
25-29 years old		Reference	Reference	
30-34 years old	0.99	0.75, 1.31	1.06	0.88, 1.29
35-39 years old	1.42	1.07, 1.89	1.56	1.27, 1.90
40-54 years old	1.35	0.91, 2.01	1.30	0.96, 1.76
<b>Payer information</b>				
Medicare	1.86	0.88, 3.95	1.77	1.02, 3.05
Medicaid	1.08	0.88, 1.32	1.22	1.05, 1.42
Private insurance		Reference	Reference	
Self-pay	0.62	0.20, 1.93	0.46	0.18, 1.17
No Charge	n/a	n/a	n/a	n/a
Other	0.69	0.32, 1.47	1.09	0.71, 1.68
<b>Hospital location</b>				
Central counties		Reference	Reference	
Fringe counties	0.78	0.60, 1.00	0.83	0.70, 1.00
Counties in metro areas >250k	0.69	0.52, 0.91	0.66	0.54, 0.80
Counties in metro areas 50-250k	0.74	0.49, 1.11	0.71	0.53, 0.95
Micropolitan	0.54	0.33, 0.90	0.69	0.48, 0.99
Not metro/micropolitan	0.88	0.56, 1.38	0.90	0.64, 1.25
<b>Median Income by ZIP Code</b>				
Income Quartile 1 <sup>st</sup>	0.97	0.73, 1.27	1.03	0.83, 1.28
Income Quartile 2 <sup>nd</sup>	0.88	0.66, 1.18	1.00	0.8, 1.24
Income Quartile 3 <sup>rd</sup>	0.96	0.72, 1.29	1.02	0.82, 1.27
Income Quartile 4 <sup>th</sup>		Reference	Reference	
<b>Diabetes</b>	0.92	0.59, 1.45	0.87	0.64, 1.19
<b>Maternal cardiac condition</b>	1.29	0.41, 4.02	1.03	0.42, 2.55
<b>Endometritis</b>	0.57	0.18, 1.78	0.78	0.36, 1.67
<b>Postpartum hemorrhage</b>	0.83	0.50, 1.39	0.99	0.70, 1.39
<b>Hospital Bed Size</b>				
Small		Reference	Reference	
Medium	0.86	0.62, 1.20	0.88	0.70, 1.10
Large	0.91	0.67, 1.23	0.92	0.75, 1.12
<b>Hospital Teaching Status</b>				
Metro non-teaching		Reference	Reference	
Metro teaching	1.27	0.99, 1.63	1.11	0.93, 1.33
Non-metro	1.07	0.67, 1.72	1.05	0.72, 1.54

The adjusted model included all of the variables listed in this table. *RR*, risk ratio. *CI*, confidence interval.

**Supplemental File 2: Unadjusted and adjusted risk for readmission for patients with severe preeclampsia restricted to hospitalizations with deliveries occurring within first two days of hospitalization**

	Unadjusted		Adjusted	
	RR	95% CI	RR	95% CI
<b>Postoperative length of stay</b>				
Length of stay <3 days		Reference		Reference
Length of stay 3 days	0.72	0.47, 1.12	0.65	0.48, 0.88
Length of stay 4 days	0.71	0.45, 1.11	0.67	0.49, 0.92
Length of stay 5 to 7 days	0.21	0.10, 0.45	0.16	0.09, 0.28
Length of stay > 7 days	0.40	0.12, 1.31	0.30	0.12, 0.78
<b>Age</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
15-19 years old	0.93	0.48, 1.79	1.32	0.88, 1.99
20-24 years old	0.88	0.56, 1.39	0.94	0.68, 1.29
25-29 years old		Reference		Reference
30-34 years old	0.96	0.63, 1.46	1.14	0.85, 1.53
35-39 years old	1.40	0.91, 2.16	1.64	1.20, 2.22
40-54 years old	1.05	0.53, 2.07	1.17	0.70, 1.95
<b>Payer information</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
Medicare	1.89	0.60, 5.99	2.28	1.04, 5.01
Medicaid	1.01	0.74, 1.37	1.09	0.86, 1.37
Private insurance		Reference		Reference
Self-pay	1.23	0.39, 3.88	0.86	0.34, 2.19
No Charge	n/a	n/a	n/a	n/a
Other	1.09	0.44, 2.68	1.73	1.05, 2.83
<b>Patient location</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
Central counties		Reference		Reference
Fringe counties	1.00	0.70, 1.42	1.02	0.79, 1.31
Counties in metro areas >250k	0.52	0.32, 0.83	0.43	0.31, 0.61
Counties in metro areas 50-250k	0.71	0.38, 1.33	0.57	0.36, 0.89
Micropolitan	0.74	0.37, 1.48	0.82	0.50, 1.35
Not metro/micropolitans	0.77	0.35, 1.66	0.58	0.33, 1.01
<b>Median Income by ZIP Code</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
Income Quartile 1 <sup>st</sup>	1.05	0.69, 1.61	1.20	0.86, 1.66
Income Quartile 2 <sup>nd</sup>	0.94	0.60, 1.48	1.02	0.73, 1.43
Income Quartile 3 <sup>rd</sup>	1.25	0.81, 1.92	1.33	0.98, 1.81
Income Quartile 4 <sup>th</sup>		Reference		Reference
<b>Diabetes</b>	0.76	0.34, 1.72		0.34, 1.18
<b>Endometritis</b>	0.37	0.05, 2.66	0.48	0.12, 1.91
<b>Postpartum hemorrhage</b>	0.68	0.32, 1.45	0.99	0.61, 1.61
<b>Hospital Bed Size</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
Small		Reference		Reference
Medium	0.54	0.34, 0.86	0.54	0.39, 0.73
Large	0.61	0.41, 0.92	0.63	0.48, 0.83
<b>Hospital Teaching Status</b>	<b>RR</b>	<b>95% CI</b>	<b>RR</b>	<b>95% CI</b>
Metro non-teaching		Reference		Reference
Metro teaching	1.27	0.86, 1.86	1.18	0.90, 1.56
Non-metro	1.40	0.67, 2.94	1.22	0.67, 2.22

The adjusted model included all of the variables listed in this table. RR, risk ratio. CI, confidence interval.