

A cosmic background image featuring a large, vibrant nebula. The nebula has a central blue region surrounded by orange and red filaments, with numerous small white stars scattered throughout the dark space.
$$1 + 2 + 3 + 4 + 5 + 6 + 7 + \dots + \infty = -\frac{1}{12}$$

Blank page



Universidad
Católica del Norte



lawphysics
Latin American Webinars on Physics

Neutrino and Dark Matter connection from spontaneous lepton number violation

Roberto A. Lineros

Departamento de Física, Universidad Católica del Norte

Seminar IFT – 21 February 2019


Science around Antofagasta




ALMA



LLAMAS




VLT






CTA

Outline



Dark Matter




Observations support Dark Matter


- Dynamics of clusters and galaxies
- Structure formation
- CMB anisotropies
- Baryon Acoustic Oscillation

$$\Omega_{\text{DM}} h^2 = 0.1196 \pm 0.0031$$



Dark Matter Searches




Neutrinos



AGM2015: Antineutrino Global Map 2015



Forero, Tortola and Valle PRD 90 (2014) 093006




The **SM** predicts zero neutrino mass

Beyond SM physics is required to explain
mass spectrum and mixing angles


The Standard Model (so far)

SM matter families



- Massless neutrinos
- Lepton number conserved
- Baryon number conserved

The Standard Model (so far)



Beyond SM



Case 1

(light) Dark Matter candidate
and neutrino masses

Majoron dark matter from a spontaneous inverse seesaw model.
N. Rojas, R. A. Lineros, F. Gonzalez-Canales. [[arxiv:1703.03416](#)]

Neutrino mass mechanisms

A large fraction of the models uses the 5-dim **Weinberg operator** to generate **majorana** neutrino masses


$$\mathcal{O}_{5ij} = \frac{1}{\Lambda} (L_i H)^T (L_j H)$$

This operator preserves SM symmetries but it breaks lepton number in **2 units**

$$\mathcal{O}_{5ij} = \frac{v^2}{\Lambda} \nu_i \nu_j = M_{ij} \nu_i \nu_j$$

Neutrino mass mechanisms

The commonly known schemes are **see-saw mechanisms**




Enters the Majoron

The Type-I seesaw can be generated via the **spontaneous** breaking of the **global U(1) lepton** symmetry

$$\mathcal{L} \supset -y_L \bar{L} H N^c - \frac{y_S}{2} S \bar{N}^c N + h.c.$$

$\begin{matrix} -1 & 0 & 1 \\ 2 & -1 & -1 \end{matrix}$

$$S = \frac{v_S + \sigma + iJ}{\sqrt{2}}$$



Enters the Majoron

$$m_D = \frac{y_L v_H}{\sqrt{2}}$$

$$M_N = \frac{y_S v_S}{\sqrt{2}}$$

After the SSB, we get the Type-I seesaw

$$\mathcal{L} \supset -m_D \bar{\nu}_L N^c - \frac{M_N}{2} \overline{N^c} N + h.c.$$

and 2 scalars: σ and J

$$m_\sigma \simeq v_S \quad m_J = 0$$

Enters the Majoron

$$m_D = \frac{y_L v_H}{\sqrt{2}}$$

$$M_N = \frac{y_S v_S}{\sqrt{2}}$$

After the SSB, we get the Type-I seesaw

$$\mathcal{L} \supset -m_D \bar{\nu}_L N^c - \frac{M_N}{2} \overline{N^c} N + h.c.$$

and 2 scalars: σ and J  DM candidate


$$m_\sigma \simeq v_S \quad m_J = 0$$

Majoron as DM _(pros)

- Neutral
- Weakly coupled to the SM
- Long lived

$$\Gamma_{J \rightarrow \nu\nu} = \frac{m_J}{32\pi} \frac{\sum_i (m_i^\nu)^2}{2v_1^2}$$

$$\Gamma_{J \rightarrow \gamma\gamma} = \frac{\alpha^2 m_J^3}{64\pi^3} \left| \sum_f N_f Q_f^2 \frac{2v_3^2}{v_2^2 v_1} (-2T_3^f) \frac{m_J^2}{12m_f^2} \right|^2$$



Details in arxiv: 1406.0004

Majoron as DM _(cons)

$$m_J = 0 \quad ! ! ! !$$

... but global symmetries are not protected under gravity effects

Therefore

$$m_J \neq 0$$

... and the majoron DM is just a **pseudo Nambu-Goldstone boson**

Majoron as DM (our proposal)

ArXiv:1703.03416

What defines a **majoron DM**?

- It is a (pseudo)scalar
- It is part of the neutrino mass mechanism
- Its signature is the decay into neutrinos
- It is massive

Inverse seesaw

The **standard** inverse seesaw

$$\mu \ll m_D \ll M$$

$$\mathcal{L} = -\frac{1}{2} n_L^T C \mathcal{M} n_L + h.c.$$

$$n_L^T = (\nu_L, N_1^c, N_2)$$

$$\mathcal{M} = \begin{pmatrix} 0 & m_D & 0 \\ m_D^T & 0 & M \\ 0 & M^T & \mu \end{pmatrix}$$

Inverse seesaw

The **standard** inverse seesaw


$$\mathcal{L} = -\frac{1}{2}n_L^T C \mathcal{M} n_L + h.c.$$

$$n_L^T = (\nu_L, N_1^c, N_2)$$

$$\mu \ll m_D \ll M$$

$$\mathcal{M} = \begin{pmatrix} 0 & m_D & 0 \\ m_D^T & 0 & M \\ 0 & M^T & \mu \end{pmatrix}$$

Lepton number
violating term



Inverse seesaw

The **standard** inverse seesaw

$$\mu \ll m_D \ll M$$

Active neutrinos

$$m_\nu = \left(\frac{m_D}{M} \right)^2 \mu$$

Heavy neutrinos

$$m_{\mathcal{N}'} = M - \frac{m_D^2}{M} + \frac{\mu}{2}$$
$$m_{\mathcal{N}} = M - \frac{m_D^2}{M} - \frac{\mu}{2}$$

Inverse seesaw

The usual inverse seesaw hierarchy:

$$\mu \ll m_D \ll M$$

Some numerology:

$$M \sim 100 \text{ TeV} \quad m_D \sim 10 \text{ GeV} \quad \mu \sim 10 \text{ MeV}$$

$$m_\nu \sim 0.1 \text{ eV}$$

$$\alpha = \frac{\mu}{M} \sim 10^{-7}$$

Spontaneous Inverse seesaw

To generate the **inverse seesaw** scheme we need **2 complex scalars**

$$\mathcal{L} = -y_L \bar{L} H N_1^c - y_S S^\dagger \bar{N}_2 N_1^c - \frac{y_X}{2} X^\dagger \bar{N}_2^c N_2 + h.c.$$

$$m_D = \frac{y_L v_h}{\sqrt{2}}, \quad M = \frac{y_S v_S}{\sqrt{2}}, \quad \text{and} \quad \mu = \frac{y_X v_X}{\sqrt{2}}$$

Spontaneous Inverse seesaw

To generate the **inverse seesaw** scheme we need **2 complex scalars**

$$\mathcal{L} = -y_L \bar{L} H N_1^c - y_S S^\dagger \bar{N}_2 N_1^c - \frac{y_X}{2} X^\dagger \bar{N}_2^c N_2 + h.c.$$

$$v_S > 50 \text{ TeV} \quad v_X > 5 \text{ MeV}$$

Spontaneous Inverse seesaw

The **charge assignments** do not follow the standard one of an Inverse See Saw

	L	N_1	N_2	S	X
$SU(2)_L$	2	1	1	1	1
$U(1)_Y$	1/2	0	0	0	0
$U(1)_I$	1	-1	x	$1 - x$	$2x$

Other assignments
are possible

$$x = 3/5$$

$$\mathcal{L} = -y_L \bar{L} H N_1^c - y_S S^\dagger \bar{N}_2 N_1^c - \frac{y_X}{2} X^\dagger \bar{N}_2^c N_2 + h.c.$$

Scalar potential

The **assignment** fixes the potential

$$\omega = \frac{v_X}{v_S}$$

$$V_{\text{scalar}} = V_{XS} + V_{HXS} + V_I$$

$$V_I = \lambda_{\text{cp}} e^{i\delta} X S^{\dagger 3} + h.c.$$

$$S = \frac{v_S e^{i\theta} + \sigma_S + i\chi_S}{\sqrt{2}}$$

$$X = \frac{v_X e^{i\tau} + \sigma_X + i\chi_X}{\sqrt{2}}$$


The tadpole equations relate the CP phases:

$$\tau = 3\theta - \delta - \pi$$

Mass spectrum

$$\omega = \frac{v_X}{v_S}$$

Now we have 5 spin-0 fields: 4 related to L breaking
1 related to EW breaking



Majoron DM stability

The only candidate is the **lightest massive scalar** i.e.



$$\zeta_2 = J_{\text{DM}}$$

We still has to satisfy the stability condition keV decaying DM:

$$\Gamma_{\text{DM}} < 10^{-43} \text{GeV}$$

Decay modes

There are potentially dangerous decay modes:



Decay into neutrinos

$$\alpha = \frac{\mu}{M} \sim 10^{-7}$$

The decay rate vanishes for:


$$\omega_0 = \sqrt{2/3}$$

$$\Gamma_\nu = \Gamma_{0\nu}(\omega_0) 4\alpha^2$$

$$\Gamma_{0\nu}(\omega_0) \simeq 10^{-40} \text{ GeV} \left(\frac{m_\nu}{0.1 \text{ eV}} \right)^2 \left(\frac{M_J}{1 \text{ keV}} \right) \left(\frac{v_S}{100 \text{ TeV}} \right)^{-2}$$

Decay into neutrinos

$$J_{\text{DM}} \rightarrow \nu\nu$$



Stability thanks to
vev alignment

Decay into scalars



$$J_{\text{DM}} \rightarrow \zeta'_{\text{S}}$$

Without a protective symmetry, this channel is not suppressed

However we can find the parameter space where the **mode vanishes**

Decay into scalars


$$J_{\text{DM}} \rightarrow \zeta' \text{'s}$$



The interplay of different diagrams allows to vanish the decay mode

Decay into scalars

$$J_{\text{DM}} \rightarrow \zeta' \text{'s}$$



There is a whole volume that satisfy this condition

Conclusions (of this part)


- The **spontaneous inverse seesaw** provides a well suited majoron DM candidate
- Our **majoron DM** is phenomenologically equivalent to the PNGB
- The **vev alignment** has a relevant role in the DM stability

Case 2

Spontaneously generated Scotogenic model

Fermion Dark Matter from Spontaneous Breaking of Lepton Number in the Scotogenic Model
C. Bonilla, L. dl Vega, J. M. Lamprea, R.L, E. Peinado [appearing soon]

Scotogenic model



Neutrino masses are generated at one loop

An extra symmetry is required to protect the loop

Dark Matter can be part of the loop


$$\mathcal{O}_{5ij} = \frac{1}{\Lambda} (L_i H)^T (L_j H)$$

Lepton number is explicitly broken

See a models zoology in Restrepo et al. arxiv:1308.3655

Scotogenic model


The simplest **scotogenic** models



E. Ma, Phys.Rev.D73:077301,2006

"Type-I"

SU(2) doublet




E. Ma, D. Suematsu Mod.Phys.Lett.A24:583-589,2009

"Type-III"

Scotogenic model


The simplest **scotogenic** models



E. Ma, Phys.Rev.D73:077301,2006

"Type-I"


$$(\mathcal{M}_\nu)_{ij} = \sum_{k=1}^3 \frac{Y_{ik}^\nu Y_{kj}^\nu m_{N_k}}{16\pi^2} \left[\frac{m_{\eta_R}^2}{m_{\eta_R}^2 - m_{N_k}^2} \log \frac{m_{\eta_R}^2}{m_{N_k}^2} - \frac{m_{\eta_I}^2}{m_{\eta_I}^2 - m_{N_k}^2} \log \frac{m_{\eta_I}^2}{m_{N_k}^2} \right]$$



Arxiv:1804.04117

$$m_N > 100 \text{ GeV}$$

Spontaneous Scotogenic




$$(\mathcal{M}_\nu)_{ij} \simeq \frac{\lambda_5 v_h^2}{16\pi^2} \sum_k \frac{Y_{ik}^\nu Y_{jk}^\nu}{m_{N_k}}$$

The scotogenic model emerge when lepton symmetry is spontaneously broken

The new scalar opens new annihilation channels


	$\bar{L}_i$	ℓ_i	H	η	N_i	ϕ
SU(2)	2	1	2	2	1	1
$U(1)_L$	1	-1	0	0	-1	2
Z_2	+	+	+	-	-	+

Channels




Leptophilic annihilation

Common to all models




majoron-higgs portal




Annihilation into majorons
< 10%

New!

Annihilation cross section



Direct detection



Conclusions (of this part)

- Scotogenic mechanism for neutrino masses give an interplay with Dark Matter
- The spontaneous version opens DM phenomenology thanks the new channels

Final words

- Neutrinos observables and DM are keys to unveil New Physics
- Spontaneously broken lepton symmetry produces an appealing DM candidate
- Scotogenic mechanism connects DM stability and neutrino masses




lawphysics

Latin American Webinars on Physics

Recent detections of gravitational waves

Isabel Cordero-Carrión
University of Valencia, Spain

Host: Joel Jones-Perez
Wednesday 13 December 2017 15:00 GMT



 /lawphysicsw

 @lawphysics

 /lawphysics

 lawphysics.wordpress.com





Thanks

The background of the slide is a deep space image featuring a large, vibrant nebula. The nebula has a central region of bright blue and white light, surrounded by swirling clouds of orange, red, and yellow gas. Numerous small, distant stars are visible as white points of light against the dark cosmic background.
$$1 + 2 + 3 + 4 + 5 + 6 + 7 + \dots + \infty = -\frac{1}{12}$$

Backup slides

Charge assignments

5 possible models

	L	N_1	N_2	S	X
$n = 1$	1	-1	$1/7$	$6/7$	$2/7$
$n = 2$	1	-1	$1/3$	$2/3$	$2/3$
$n = 3$	1	-1	$3/5$	$2/5$	$6/5$

$$m+n = 4$$

$$V_I = \lambda_{\text{cp}} e^{i\delta} X^m S^{\dagger n}$$

$$m+n = 3$$

	L	N_1	N_2	S	X
$n = 1$	1	-1	$1/5$	$4/5$	$2/5$
$n = 2$	1	-1	$1/2$	$1/2$	1

The rest of the scalar potential

$$V_{SX} = -\mu_S^2 |S|^2 + \frac{\lambda_S}{4} |S|^4 - \mu_X^2 |X|^2 + \frac{\lambda_X}{4} |X|^4 + \lambda_5 |S|^2 |X|^2 + V_I$$

$$V_{HSX} = -\mu_H^2 H^\dagger H + \frac{\lambda_H}{4} (H^\dagger H)^2 + \lambda_{HS} |S|^2 H^\dagger H + \lambda_{HX} |X|^2 H^\dagger H$$

Mass spectrum

$$m_h^2 \simeq \frac{v_h^2}{2} \left\{ \frac{\lambda_H}{2} + 2 \left(\frac{\lambda_{HX}^2 \lambda_S + \lambda_{HS}^2 \lambda_X - 4\lambda_5 \lambda_{HS} \lambda_{HX}}{4\lambda_5^2 - \lambda_S \lambda_X} \right) \right\}$$

$$M_{\zeta_3}^2 \simeq \frac{v_S^2}{2} \left(\frac{-A + A\psi + 2\lambda_X \omega \psi}{2\psi} \right)$$

$$M_{\zeta_4}^2 \simeq \frac{v_S^2}{2} \left(\frac{A + A\psi + 2\lambda_X \omega \psi}{2\psi} \right)$$

$$\lambda_S = A + \lambda_X \omega^2$$

$$\lambda_5 = -A \left(\frac{\sqrt{1 - \psi^2}}{4\omega\psi} \right)$$

Numerology

Parameter	Value
M	100 TeV
μ	10 MeV
m_D	10 GeV
v_S	$10^8 - 10^{12}$ GeV
ω	0.4 – 1.6

$$\lambda_{\text{cp}} \simeq \frac{M_J^2}{v_S^2} < 10^{-22}$$