Identification of the Minimum Effective Dose for Normally Distributed Endpoints Using a Model Selection Approach: Supplementary Material

Rebecca M. Kuiper¹, Daniel Gerhard², Ludwig A. Hothorn²

¹ Utrecht University, Department of Methodology and Statistics, PO box 80140,
NL-3508TC Utrecht, The Netherlands (E-mail: *R.M.Kuiper@uu.nl*).

² Leibniz University Hannover, Institute for Biostatistics, Herrenhäuser Straße
2, D-30419 Hannover, Germany.

Figure 1: Simulated MED selection rates for the GORIC and the Dunnett- and Williams-type multiple contrast test ($\alpha = .05$) given a non-centrality parameter based on a **linear** dose-response profile. Several relevance margins ($\delta = 0, 1, 2, 3$) are presented in the columns.

Figure 2: Simulated MED selection rates for the GORIC and the Dunnett- and Williams-type multiple contrast test ($\alpha = .05$) given a non-centrality parameter based on a **convex** dose-response profile. Several relevance margins ($\delta = 0, 1, 2, 3$) are presented in the columns.

Figure 3: Simulated MED selection rates for the GORIC and the Dunnett- and Williams-type multiple contrast test ($\alpha = .05$) given a non-centrality parameter based on a **semi-concave** dose-response profile. Several relevance margins ($\delta = 0, 1, 2, 3$) are presented in the columns.

Figure 4: Simulated MED selection rates for the GORIC and the Dunnett- and Williams-type multiple contrast test ($\alpha = .05$) given a non-centrality parameter based on a **concave** dose-response profile. Several relevance margins ($\delta = 0, 1, 2, 3$) are presented in the columns.