

Gravitational fields of tesserooids with variable density

Santiago R. Soler^{1,2}, Agustina Pesce^{1,2}, Mario E. Gimenez^{1,2} and Leonardo Uieda³

¹ CONICET ² IGSV, UNSJ ³ Department of Earth Sciences, SOEST, University of Hawai'i at Mānoa, USA.


Introduction

When computing gravitational fields on regional or global scales it's necessary to take into account the curvature of the Earth. One way to do this is using spherical prisms, also known as **tesserooids**. Most methodologies consider only Tesserooids with homogeneous, linear or polynomial densities in depth. We present a new methodology to compute the **gravitational fields** generated by **tesserooids** with any **continuous density** variation through a **Gauss-Legendre Quadrature** and two **discretization algorithms**.

Gauss-Legendre Quadrature


The gravitational potential generated by a tesserooid with variable density in depth can be numerically approximated through a **Gauss-Legendre Quadrature (GLQ)**.

The GLQ is equivalent to approximating the **tesserooid** with **point masses** located on the **nodes** of the Quadrature.




2D Adaptive Discretization

The **approximation** becomes **less accurate** when the **computation point** gets **closer** to the **tesserooid**. To reduce this error we use a **two dimensional adaptive discretization** algorithm.





Density-based Discretization

We have developed a complementary discretization algorithm that reduces the errors due to the variations of the density function. In short, it **divides the tesserooid** along the **radial dimension** at the depths at which the **maximum density variation** takes place.




Density-based discretization examples



The **density-based** discretization **automatically** adjust the **number of splits** depending on the **variability of the function**.


Determination of D and delta

We want to find the **optimal values** of **D** and **delta** that guarantee **accuracy** while minimizing the **computation time**. In order to do that we **compared the numerical results** with the **analytical solutions** of a **spherical shell**.



Forward Model: Neuquén Basin

We modelled the **Neuquén basin** using an **exponential density profile** in order to account for the **compaction of the sediments**.




Conclusions

- New **fully automated** method to compute **gravitational fields** of **continuous density tesserooids**.
- The **density-based discretization** algorithm **automatically** adjusts number of **radial splits**.
- **Linear density** tesserooids **don't need** density-based **discretizations**.
- To achieve a **0.1% accuracy**:
 - ♦ **D = 1** for potential,
 - ♦ **D = 2.5** for accelerations,
 - ♦ **delta = 0.1** sufficient for most geophysical applications.
- Starting point for **non-linear gravity inversion** of large scale **sedimentary basins**.

References

- Fukushima, T. (2018) doi: 10.1007/s00190-018-1126-2
 Li, Z., Hao, T., Xu, Y. and Xu, Y. (2011) doi: 10.1016/j.jappgeo.2011.01.004
 Grombein, T., Seitz, K. and Heck, B. (2013) doi: 10.1007/s00190-013-0636-1
 Lin, M. & Denker, H. (2018) doi: 10.1007/s00190-018-1193-4
 Uieda, L., Barbosa, V. C. and Braitenberg, C. (2016) doi: 10.1190/geo2015-0204.1


Feel free to photograph and share this poster



Content licensed under Creative Commons Attribution 4.0 International License.



Download the source code
github.com/pinga-lab/tesserooid-variable-density



Download the poster
github.com/santisoler/lapis2019



santisoler



santiago.r.soler@gmail.com

This research was possible thanks to a PhD Scholarship granted by CONICET.



INSTITUTO
GEOFÍSICO-
SISMOLÓGICO
ING. VOLPONI

