

Supporting information for
A comparison of different oxidative pretreatments on polysaccharide
hydrolyzability and cell wall structure for interpreting the greatly
improved enzymatic digestibility of sugarcane bagasse by
delignification

Yazhu Han ^{1,2}, Yuchen Bai ^{1,2}, Jingzhi Zhang ^{1,2}, Dehua Liu ^{1,2}, Xuebing Zhao ^{1,2,*}

¹ Key Laboratory of Industrial Biocatalysis, Ministry of Education, Tsinghua University, Beijing 100084, China;

² Institute of Applied Chemistry, Department of Chemical Engineering, Tsinghua University, Beijing 100084, China

** Corresponding authors: Email: zhaoxb@mail.tsinghua.edu.cn; Tel: +86-10-62772130; Fax: +86-10-62772130*

Figure S1 Effects of 5g/L Tween on enzymatic hydrolysis of different oxidative pretreatment of sugarcane bagasse. Cellulase loading was 15 FPU/g solid




Figure S2 XRD diagrams (A) and enzymatic hydrolysis (B) of filter paper treated by sodium chlorite (SC) for different time




Figure S3 Effects of addition of wheat straw klason lignin on enzymatic hydrolysis of filter paper. The cellulase cocktail use was Novozyme CTec 2.

