

SUPPLEMENTARY DATA

Anti-Allergic Effects of Phenolic Compounds Isolated from *Stellera chamaejasme* on RBL-2H3 Cells

Beom-Geun Jo^{1,#}, Sim-Kyu Bong^{2,#}, Jonghwan Jegal¹, Su-Nam Kim², and Min Hye Yang¹


¹College of Pharmacy, Pusan National University; Busan 46241, South Korea

²Natural Products Research Institute, Korea Institute of Science and Technology; Gangneung 25451, South Korea


Supporting Information Contents:

- S1.** The ¹H NMR spectrum of compound **9**
- S2.** The ¹³C NMR spectrum of compound **9**
- S3.** The ¹H-¹H COSY spectrum of compound **9**
- S4.** The ¹H-¹H NOESY spectrum of compound **9**
- S5.** The HMQC spectrum of compound **9**
- S6.** The HMBC spectrum of compound **9**


S1. The ^1H NMR spectrum of compound **9**




S2. The ^{13}C NMR spectrum of compound **9**




S3. The ^1H - ^1H COSY spectrum of compound **9**



S4. The ^1H - ^1H NOESY spectrum of compound **9**



S5. The HMQC spectrum of compound **9**



S6. The HMBC spectrum of compound **9**

