Hepatalin

The missing link in prediabetes, obesity, and type 2 diabetes

W. Wayne Lautt

Professor Emeritus

Max Rady Faculty of Health Sciences

University of Manitoba

260 Brodie Center

727 McDermot Avenue

Winnipeg, Manitoba R3E 3P5

Canada

and

Cofounder and Chief Scientific Officer

SciMar Ltd.

119 Main Street South

Dauphin, Manitoba R7N 1K4

Canada

Wayne.Lautt@scimar.ca

Abstract

Hepatalin is a hormone secreted by the liver in response to pulses of insulin after a mixed nutrient meal, but only if the liver receives two permissive synergistic feeding signals from the stomach. Hepatalin stimulates glucose uptake and storage as glycogen in skeletal muscle, heart, and kidney but not liver, guts, or adipocytes. Insulin acts primarily on liver and fat. Reduced hepatalin action results in postprandial hyperglycemia, compensatory elevation of insulin secretion, and a resultant shift in partitioning of nutrient energy storage from glycogen in muscle, to fat. Chronic hepatalin suppression leads to a predictable chronology of dysfunctions, first diagnosable as Absence of Meal-induced Insulin Sensitization (AMIS) which progresses to prediabetes, adiposity, and type 2 diabetes (see graphic abstract). The focus on nutrient partitioning and the role of hepatalin allows AMIS to be diagnosed, prevented, and treated, including through the use of lifestyle interventions.

Keywords: AMIS, AMISS, RIST, hepatalin, HISS, prediabetes, obesity, nutrient partitioning, type 2 diabetes, insulin

Graphic Abstract



Contents	
1. IS THERE A MISSING LINK?	5
2. SCREENING:	6
2.1 POSTPRANDIAL HYPERGLYCEMIA	6
2.2 POSTPRANDIAL HYPERLIPIDEMIA	6
2.3 POSTPRANDIAL HYPERINSULINEMIA	7
2.4 POSTPRANDIAL REACTIVE OXIDATIVE STRESS	7
3. HEPATALIN AND THE AMISS PARADIGM	9
3.1 THE EARLY SCIENCE OF HEPATALIN	9
3.2 TERMINOLOGY – HISS, MIS, AMIS, AMISS	. 10
4. QUANTIFYING DYNAMIC INSULIN AND HEPATALIN ACTION, THE RIST	. 12
5. EVIDENCE OF MEAL-INDUCED INSULIN SENSITIZATION (MIS)	. 14
5.1 NUTRIENT PARTITIONING	. 15
6. REGULATION OF HEPATALIN PRODUCTION	. 17
6.1 THE PARASYMPATHETIC SIGNAL	. 18
6.1.1 ATROPINE	. 21
6.2 THE GLUTATHIONE (GSH) SIGNAL	. 23
6.3 THE INSULIN SIGNAL	24
6.4 THERAPEUTIC MIMICKING OF THE NERVE AND GSH FEEDING SIGNALS	. 25
6.5 REDUNDANT CONTROL SYSTEMS	. 26
7. ABSENCE OF MEAL-INDUCED INSULIN SENSITIZATION (AMIS)	. 27
8. HEPATALIN AND THE AMIS SYNDROME (AMISS)	. 28
8.1 AGE AND AMISS	. 30
8.2 ANTIOXIDANTS AND AMISS	.31
8.3 EXERCISE AND AMISS	34
8.4 DIET AND AMISS	. 34
8.5 STRESS AND AMISS	. 35
8.5.1 ACUTE STRESS AND AMISS	. 35
8.5.2 PSYCHOLOGICAL STRESS AND AMISS	. 35
8.6 GLUCAGON AND AMISS	.36
8.7 HEART FUNCTION AND AMISS	36
8.8 VASCULATURE AND AMISS	37
8.9 OBESITY AND AMISS	. 38
8.10 SEX AND AMISS	41
8.10.1 FETAL ALCOHOL EFFECT AND AMISS	. 41
8.10.2 GESTATION AND AMISS	. 41
8.10.3 HUMAN FEMALE DATA	. 42
9.IMPACTS OF THE NEW PARADIGM	. 44
9.1 ACCELERATED KNOWLEDGE	.44
9.2 NEW PARADIGM-BASED PRODUCTS	44
a) SciMar NuPa Daily®,	.44
b) SciMar NuPa Test [®]	.44
c) SciMar NuPa Renew [®]	.44
9.3 FOCUS ON LIFESTYLE	.45
REFERENCES	48

1. IS THERE A MISSING LINK?

Approximately 450 million people around the world have currently been diagnosed with type 2 diabetes (International Diabetes Federation, 2019), and those are only the ones with access to conventional diagnostics. Since the discovery of insulin 100 years ago, the paradigm for dealing with diabetes has focused on insulin, with blood glucose levels as the primary diagnostic and the index of therapeutic control. A multitude of pharmaceutical interventions have come and gone, but the 'pandemic' of obesity and type 2 diabetes continues to accelerate, threatening to overwhelm health care capacity in even the wealthiest countries. Insulin analogs and secretagogues have been shown to lower blood glucose levels, but progression to the pathologies associated with diabetes is only delayed, not prevented. The problem is that the focus of research and intervention has not been on the initiating event. While standard screening has been based on data from the stable fasted state, the actual dysfunction is in the dynamic postprandial processing of the nutrient energy from a meal, i.e., in the fed state. By the time the diagnosis of type 2 diabetes is made based on fasting glucose levels or HbA1c, the process of what we have termed the AMIS syndrome is well underway (Section 8, Figure 5). There is clearly a missing link in the understanding of this disease.

2. SCREENING:

The current approach of screening for type 2 diabetes using the fasting metabolic status is not effective. Ceriello (2000) concluded that "... paradoxically, the vast majority of the studies on cardiovascular disease risk factors have been conducted by measuring them in strictly fasting conditions. This simply means that most of the data available to date may not reflect the real situation". Postprandial hyperglycemia, hyperinsulinemia, hypertriglyceridemia, and oxidative stress have all been reported to be independent risk factors for cardiovascular disease (reviewed in Lautt, 2007). Similarly, these postprandial, rather than fasted metabolic markers, are also better indicators of prediabetic and diabetic dysmetabolism.

2.1 POSTPRANDIAL HYPERGLYCEMIA: The strongest age- and sex-adjusted risk for allcause and cardiovascular mortality has been associated with two-hour post-load plasma glucose levels (de Vegt et al., 1999). Increased mortality risk has been associated with two-hour post-load plasma glucose levels to a much greater extent than with fasting plasma glucose (Simon et al., 1987; Hanefield et al., 1996; DECODE Study Group, 1999; Shaw et al., 1999; Vaccaro et al., 1999; Engelgau et al., 2000; Simon and Brandenberger, 2002). Postprandial plasma glucose has been shown to be a determinant of both onset and development of nephropathy in type 2 diabetic patients (Shichiri et al., 2000). Acute hyperglycemia was associated with an acute increase in clotting factor VII (Ceriello et al., 1988) and enhanced thrombin activity that was proportional to the level of hyperglycemia (Ceriello et al., 1996).

2.2 POSTPRANDIAL HYPERLIPIDEMIA: Postprandial elevation of blood lipids is prevalent in diabetic patients even with normal fasting triglyceride concentrations (Mero et al., 1998). The postprandial triglyceride increment after a fat meal was paralleled by an inflammatory

leukocyte increment due to an increase in neutrophils in the first two postprandial hours (Van Oostrom et al., 2003). Low-density lipoprotein oxidation increased after meals and was directly related to the degree of hyperglycemia (Ceriello et al., 1999). Postprandial hypertriglyceridemia has been recognized as an independent predictor of cardiovascular pathology (Anderson et al., 2001).

2.3 POSTPRANDIAL HYPERINSULINEMIA: Ingelson et al. (2005) discussed postprandial hyperinsulinemia as a risk factor for cardiovascular disease and proposed several possible mechanisms. Typically, as anomalies of an existing paradigm accumulate, the defense of the existing paradigm becomes convoluted. The discovery of a previously unknown hormone, hepatalin, allows re-interpretation of much of the literature. Absence of postprandial hepatalin secretion accounts for compensatory postprandial hyperinsulinemia, which then has consequences. Insulin has an impact on many systems, some acutely controlling glucose metabolism, and some affecting protein and lipid metabolism. The inability to secrete hepatalin after a meal is proposed to explain the hyperinsulinemia as a compensatory reaction to the absence of hepatalin action. The assumption is that restoration of postprandial hepatalin secretion would be insulin-sparing and avoid hyperinsulinemia.

2.4 POSTPRANDIAL REACTIVE OXIDATIVE STRESS: From his review in 2001, Brownlee concluded that hyperglycemia-induced overproduction of superoxide by the mitochondrial electron transport chain likely accounts for the main molecular mechanisms implicated in glucose-mediated vascular damage associated with blindness, renal failure, nerve damage, atherosclerosis, stroke, and limb amputation. Ceriello (2000) suggested that increased

postprandial oxidative stress may be the common pathway through which the majority of diabetic complications occur.

Although postprandial hyperglycemia, hyperinsulinemia, hyperlipidemia, and oxidative stress are well accepted cardiovascular risk factors associated with prediabetes and type 2 diabetes, the mechanism connecting them has been unknown. There is a missing link.

3. HEPATALIN AND THE AMISS PARADIGM:

The bold claim made in this review is that the discovery of a hepatic hormone that accounts for the majority of glucose uptake after feeding, represents the missing link in understanding the aetiology and mechanism of metabolic dysfunction in prediabetes, which in turn leads to obesity and type 2 diabetes, followed by the full array of associated pathologies. Until this review, the tentatively named putative hormone has been referred to in the scientific literature as Hepatic Insulin Sensitizing Substance (HISS). Over time, the biological activity of this hormone has been quantified, and descriptions of its regulatory systems, its actions, and its role in pathology, have been published. As research continues, we introduce the new name "hepatalin" in this review.

The discovery of hepatalin and ongoing research have presented opportunities for novel early diagnosis, prevention, and treatment of prediabetes and type 2 diabetes.

3.1 THE EARLY SCIENCE OF HEPATALIN: Electrical stimulation of hepatic parasympathetic nerves resulted in the net fasting efflux of glucose from the liver to be completely blocked within minutes (Lautt and Wong, 1978). Hepatic parasympathetic neuropathy was suggested to initiate type 2 diabetes via an ineffective parasympathetic signal and a potential therapeutic direction was proposed (Lautt, 1980). In the absence of knowledge of the existence of hepatalin, this suggestion was not pursued by us or others.

In 1993 (Xie et al., 1993) we reported that denervation of the anterior hepatic nerve plexus resulted in a decreased whole body hypoglycemic response to insulin. The initial hypothesis was that the impaired response to insulin was a consequence of impaired hepatic glucose uptake. However, in 1996 (Xie and Lautt, 1996a) we reported that surgical denervation

of the liver or intraportal atropine administration resulted in a reduced response to insulin in skeletal muscle but not liver or guts, and that the effect on the hindlimbs could be restored by intraportal infusion of acetylcholine at doses too low to recirculate to the systemic circulation (Xie and Lautt, 1996b). The signal from the liver to the periphery was via the blood. The action of a previously unknown hormone had been revealed.

The research that followed for the next 30 years was pure basic research focused on understanding the role and control of this newly discovered hormone. The first demonstration of hepatalin action in humans was carried out by the Macedo team in Lisbon. The findings were strikingly similar to the data from several other species (Patarrao et al., 2008) (Figure 1 in humans, Figure 3 in rats).

3.2 TERMINOLOGY – HISS, MIS, AMIS, AMISS: As studies of this new paradigm progressed, new methodology and terminology were required. Hepatic Insulin Sensitizing Substance (HISS) was the tentative name we gave to the substance that was secreted from the liver and acted on skeletal muscle to potentiate the response to insulin after feeding (Lautt, 1988). The first review of the HISS story in 1999 (Lautt, 1999) left the question open as to whether HISS sensitized muscle to insulin or had direct glucose uptake action, thus the term 'sensitizing substance' was tentatively retained for convenient reference. It was later shown that the hepatic hormone does not interact with the insulin signal, but has independent and selective action on glucose uptake, and is tightly regulated by different neuroendocrine signals. Accordingly, to recognize the relationship between the liver and insulin, we more appropriately named this hormone hepatalin.

As a result of hepatalin action, the glucose uptake response to insulin is at least doubled after a meal. We named this response Meal-induced Insulin Sensitization (MIS). If hepatalin secretion is absent during a meal, we referred to this condition as Absence of Meal-induced Insulin Sensitization (AMIS). Finally, the predictable progression of related pathologies seen with chronic AMIS has been referred to as the AMIS Syndrome (AMISS) (Ming et al., 2009; Lautt et al., 2010).

4. QUANTIFYING DYNAMIC INSULIN AND HEPATALIN ACTION, THE RIST:

Endogenous insulin is secreted in pulses of oscillations of about 13 minutes in both humans and rats (Lang et al., 1979; Chou et al., 1991) and 9 minutes in monkeys (Goodner et al., 1977). Increased insulin secretion is associated with an increase in magnitude, not period. Non-pulsatile administration is not physiological and results in a linear duration-dependent inhibition of hepatalin action beginning at infusion duration of 10 minutes, with insignificant hepatalin action remaining after one hour of constant infusion (Reid and Lautt, 2004). Hepatalin action can therefore not be assessed by the classical two-to-three-hour hyperinsulinemic euglycemic clamp.

The dynamic action of insulin and hepatalin can be determined using the insulin tolerance test based on the degree of induced hypoglycemia in the first 30 minutes after injection and before counter-regulatory mechanisms act to restore glucose to basal levels (Reid et al., 2002; Afonso et al., 2016, Martins et al., 2016). However, the ensuing hypoglycemia limits the use of this test. To avoid the hypoglycemia, and to detect the dynamic response to pulses of insulin, the Rapid Insulin Sensitivity Test (RIST) was developed and has been used in cats (Xie et al., 1996), anesthetized (Lautt et al., 1998) or conscious (Latour and Lautt, 2002) rats, mice (Latour and Chan, 2002), rabbits (Peitl et al., 2006), and humans (Patarrao et al., 2007, 2008, 2012).

The RIST is a rapidly sampled and adjusted euglycemic clamp to quantify the glucose uptake response to a pulse of insulin. The RIST provides reproducible test scores up to four consecutive times in anesthetized rats (Lautt et al., 1998). An operating procedure has been described for the RIST in animals (Lautt et al., 1998) and humans (Patarrao et al., 2008). The

RIST index is the amount of glucose that is required to be infused to compensate for glucose uptake while maintaining a constant arterial glucose level after administration of a pulse of insulin. An advantage of using the RIST index is that protocols can be designed to exploit the multiple and reproducible tests in anesthetized and conscious preparations, providing both greater statistical power and economy of animal use.

5. EVIDENCE OF MEAL-INDUCED INSULIN SENSITIZATION (MIS):

The concept of MIS was derived from the observation that the dynamic response to insulin, determined after a 24-hour fast, is at least doubled when tested 100 minutes after administration of a mixed meal in conscious rats (Latour and Lautt, 2002b; Sadri et al., 2006) and humans (Patarrao et al., 2007, 2008, 2012). This greatly amplified response to insulin is through the action of the hormone hepatalin and is regulated by parasympathetic nerves.

The effect of feeding on insulin action (MIS) was reported in 2001 (Lautt et al., 2001). This increased postprandial response to insulin was attributed to a previously unknown hormone which we tentatively named Hepatic Insulin Sensitizing Substance (HISS). An overview of its hypothetical regulation and role was described by Lautt in 1999. Several reviews followed the progress in concepts, including: the effects of adult and fetal alcohol exposure on MIS (Ting and Lautt, 2006); the effect and interactions of diet, exercise, and antioxidant supplementation on the AMIS syndrome and hepatalin action (Chowdhury et al., 2013b); adiposity as an early component of the AMIS syndrome (Lautt and Wang, 2014a); AMISS and fatty heart and liver (Lautt et al., 2015); and a review of the role of the parasympathetic nerves in metabolic regulation in health and disease (Macedo et al., 2013).





Figure 1: Rapid Insulin Sensitivity Test (RIST) in fasted and fed humans showing Meal-induced Insulin Sensitization (MIS). The rate of glucose infusion required to maintain baseline euglycemia is shown for healthy male volunteers after administration of a bolus of 50 mU/kg insulin in the 24-hour fasted state, and 100 minutes after consumption of a mixed test meal. The mean RIST curves were obtained by averaging glucose infusion rates at 0.1-minute intervals. The RIST index increased from a fasted level of 215 to 681 mg glucose/kg after feeding due to the action of hepatalin. Hepatalin action shown in the second panel is calculated from the difference in the curves in the first panel. (Figure modified from Patarrao et al., 2008, Can. J. Physiol. Pharmacol. 86: 880–888.)

5.1 NUTRIENT PARTITIONING: The nutrient energy contained in a meal must be rapidly processed and stored for use between meals. The primary storage sites are as glycogen and lipids. Insulin action is primarily on liver and adipose tissue. Hepatalin action is primarily on muscle, heart, and kidney (Section 8). A healthy balance between the actions of hepatalin and insulin results in the majority of nutrient partitioning going to glycogen in muscle. As hepatalin action decreases, the partitioning shifts from glycogen to lipids, with accumulated adiposity and development of an early prediabetic state.

Regulation of nutrient partitioning is dynamic and affected by many factors (as shown in Figure 5). For example, gestation requires a dramatic shift in nutrient energy use and storage for the changing needs of the mother and fetus. In the equivalent of the first trimester, while tissues and organs are being formed in the rat fetus, hepatalin action predominates. By the third trimester, to the benefit of the future nursing mother and newborn, lipid accumulation in the fetus and mother is facilitated by reduced hepatalin action and elevated insulin levels (Lovat et al., 2020). A sucrose-supplemented diet shifts the balance away from hepatalin and results in an earlier hyperinsulinemia with resultant increased adiposity (Lovat et al., 2021).

Nutrient partitioning is also affected by physical activity (reviewed by Chowdhury et al., 2013b). Hepatalin and insulin action in rats were determined at 9, 14, and 21 weeks of age with and without access to a running wheel for 1 week. At all ages, exercise increased the postprandial response to insulin, because of increased hepatalin secretion. The hepatalin-independent, direct action of insulin was not significantly improved. Decreased hepatalin action correlated positively with adiposity.

6. REGULATION OF HEPATALIN PRODUCTION:

Hepatalin secretion from the liver is regulated by three signals mediated by parasympathetic nerves, elevated hepatic glutathione (GSH), and pulses of insulin.



Hepatalin Production

Figure 2: Hepatalin synthesis hypothesis. Hepatalin production is regulated by the synergistic permissive action of feeding signals from the stomach activating hepatic cholinergic muscarinic stimulation of nitric oxide synthase to produce NO and activate generation of cGMP. Hepatic glutathione levels increase by about 50%. In the presence of both synergistic, permissive feeding signals, pulses of insulin stimulate secretion of pulses of hepatalin. The degree of production of hepatalin determines the ratio of partitioning of nutrient energy storage as glycogen or lipids. At this time, we speculate that hepatalin is a modified fragment of the insulin molecule, probably nitrosylated (Lautt and Wang, 2014b).

6.1 THE PARASYMPATHETIC SIGNAL: The nerve signal is activated by the presence of a mixed-nutrient meal in the stomach or upper GI tract (Afonso et al., 2016) and travels via the vagus nerve to the hepatic branch of the parasympathetic nerves (Latour and Lautt, 2002a). Neither pure glucose nor sucrose activate the nerve signal (Sadri et al., 2006; Afonso et al., 2016). Also, given separately, glucose, amino acids, and lipids do not activate MIS (Afonso et al., 2016).

The hepatic parasympathetic nerves release acetylcholine (ACh), acting on muscarinic receptors (Xie and Lautt, 1995b) to activate nitric oxide synthase, leading to hepatic nitric oxide (NO) synthesis, which activates hepatic guanylate cyclase (Guarino et al., 2004). Atropine or surgical denervation of the hepatic parasympathetic nerves blocks hepatalin secretion which can be reversed by either a cholinergic agonist (Xie and Lautt, 1996a, b) or an NO donor (Sadri and Lautt, 1999); the effect of blocking NO synthase production of NO can be reversed by an NO donor but not ACh (Guarino et al., 2001). Thus, NO production is dependent on a cholinergic signal.

The parasympathetic signal is absent in obstructive hepatitis. Chronic bile duct ligation resulted in AMIS that was reversed by intraportal infusion of ACh (Lautt and Xie, 1998), suggesting that the GSH signal had not been blocked. In contrast, diet-induced AMIS results in absence of both the nerve and GSH signal (Lautt et al., 2011) (Section 6.4, Figure 5). The earliest manifestation of diabetic autonomic neuropathy is associated with parasympathetic abnormality (Hosking et al., 1978) (Section 8.1).

For at least six hours after a meal, the response to insulin in the anesthetized state does not decrease. However, fasting in the conscious state results in a decrease of hepatalin

secretion in response to insulin by about 10% per hour until becoming insignificant after a 24hour fast (Lautt et al., 2001; Schafer et al., 2010). The reduced AMIS seen after eight hours of fasting can be restored to fed levels by potentiating the parasympathetic signal using an acetylcholinesterase antagonist (Schafer et al., 2010), suggesting that at this time point GSH levels have not been decreased, and that the early reduction in hepatalin secretion after fasting is regulated by the nerve signal.



Parasympathetic Control of MIS

Figure 3: Blocking hepatalin secretion via the nerve signal by three methods. Elimination of hepatalin action in fed rats was done by blocking the parasympathetic feeding signal by surgical denervation, blocking cholinergic muscarinic receptors (atropine 3 mg/kg), and prevention of production of nitric oxide by blocking nitric oxide synthase (L-NMMA 0.73 mg/kg). The mean RIST curves were obtained by averaging glucose infusion rates at 0.1-minute intervals throughout the test response to 50 mU/kg insulin. The control RIST was done in the stable fed state and included the combined action of hepatalin and insulin. Blocking hepatalin action. The dynamic hepatalin-dependent component of insulin action is calculated from the difference between the curves on the left. Hepatalin action commenced at 3.1 minutes after the onset of insulin administration and continued for 9 to 10 minutes after completion of hepatalin-independent insulin action. (Figure modified from Lautt et al., 2001, Am. J. Physiol. Gastrointest. Liver Physiol. 281: G29–G36).

6.1.1 ATROPINE: Atropine is a useful tool to block hepatalin secretion in a postprandial steady state (which is required before a valid RIST can be carried out (Lautt, 2003)). Atropine did not, however, produce a further reduction in response to insulin after surgical denervation had already blocked hepatalin secretion (Lautt et al., 2001). Atropine produces a dose-dependent degree of AMIS by blocking hepatalin secretion but has no effect on the response to insulin in the fasted state when hepatalin secretion has already been physiologically suppressed by absence of the nerve signal and reduced GSH level (Lautt et al., 2011).

The use of atropine as a tool to differentiate hepatalin and insulin action has been controversial because of the suggestion that atropine at an appropriate dose and at an appropriate time, blocks the ability of insulin to stimulate hepatalin secretion without having other major metabolic consequences relevant to the acute dynamic glucose uptake response to a pulse of insulin.



Two Methods to Quantify MIS Using the RIST

Figure 4: The hepatalin component of the response to insulin can be quantified in two ways using the RIST. This example is from 24-hour fasted, conscious, unrestrained rats showing both approaches. To show the development of MIS, the first RIST is done before a meal (left panel) and the second after a stable glycemic baseline has been determined after the meal (center panel). The RIST index was increased by 90%, representing combined insulin and hepatalin action (this procedure was used in the human RISTs (Figure 1)). The second method is to do the first test in the fed stable state (center panel), and then block hepatalin secretion using atropine (right panel) or blocking nitric oxide production or surgically denervating the liver (Figure 3). Atropine completely reversed MIS with no effect on hepatalin-independent insulin action. (Figure modified from Sadri et al., 2006, Br. J. Nutr. 95: 288–295).

The glucose-uptake response to insulin in a 24-hour fasted conscious unrestrained rat doubled when measured 90 minutes after gastric injection of a test meal. Atropine administration eliminated this potentiated response resulting in a RIST index identical to that seen in the fasted state despite other maintained postprandial conditions (Sadri et al., 2006). Atropine has no effect on insulin action in fasted rats, where physiological suppression of hepatalin release already exists (Lautt, 2003). Similarly, if hepatic nerves are cut to eliminate the permissive neural feeding signal, hepatalin release is blocked in the fed state, and the remaining direct response to insulin is not altered by atropine and is not different from the fasting response (Lautt et al., 2001). Atropine has no effect on the RIST index if hepatalin release has already been blocked by chronic dietary sucrose (Ribeiro et al., 2005; Ming et al., 2009) or fat supplementation (Afonso et al., 2007, 2010) or stress (Seredycz et al., 2006).

A precautionary note is that any drug used to block the hepatalin pathway must be administered at a stable postprandial glycemic state to avoid neuroendocrine involvement with breakdown and absorption of nutrients from the guts.

6.2 THE GLUTATHIONE (GSH) SIGNAL: Hepatic GSH levels increase by ~40–50% after a meal (Lautt et al., 2011; Guarino et al., 2004). Cholinergic mimicking of the parasympathetic signal does not reverse diet-induced AMIS unless GSH levels are also raised to normal fed levels (Section 6.4). The source of hepatic GSH is unclear but blocking production blocks hepatalin release (Guarino et al., 2004).

To reverse chronic sucrose-induced AMIS, GSH levels must be elevated to postprandial levels using, for example N-acetyl cysteine (Lautt et al., 2011). The role of GSH in this signaling pathway is unclear. However, nitrosylated GSH can trigger guanylate cyclase to stimulate cyclic GMP synthesis (Hogg, 2002) which in turn suggests that nitrosylated GSH may be an intermediate between NO synthesis and guanylate cyclase in the hepatalin synthesis pathway. Nitrosylated GSH can also trigger cyclic GMP independent S-nitrosylation of proteins (Guarino et al., 2004; Hogg, 2002).

The GSH signal is subject to many insults and regulations. Glucagon levels are increased during fasting when hepatic GSH levels are low. Feeding results in reduced glucagon and elevated GSH. Glucagon suppression of hepatalin secretion is mediated by cAMP (Patarrao et

al., 2015). A cAMP analog (DBcAMP) and glucagon produced a decrease of insulin sensitivity in a dose-dependent manner correlated with a reduction of hepatic GSH content. The response to insulin was restored by administration of the GSH analogue (Patarrao et al., 2015).

Acute alcohol also results in AMIS associated with reduced hepatic GSH (Lautt et al., 2005; Ting and Lautt, 2006). Fasting and diets high in sugars reduce the GSH feeding signal (Lautt et al., 2011). The GSH signal is absent after a 24-hour fast but appears intact after an 8-hour fast, as potentiating the nerve signal by blocking acetylcholine catabolism with an acetylcholinesterase antagonist (neostigmine) fully restores MIS to fed levels (Schafer et al., 2010).

6.3 THE INSULIN SIGNAL: To elicit maximal hepatalin release in response to insulin, the dose must be administered as a pulse over a duration of less than 10 minutes. Administration of 50 mU/kg insulin over 30 seconds or five minutes resulted in similar dynamic responses, but infusions over a longer period resulted in a linear duration-dependent suppression of hepatalin release that was complete by one hour of constant infusion (Reid and Lautt, 2004; Reid et al., 2002). The prolonged constant administration of insulin used in the 'gold standard' hyperinsulinemic euglycemic clamp is not capable of detecting hepatalin action, but actually blocks MIS.

Hepatalin action is linearly related to the dose of insulin with approximately 56% of the postprandial response to insulin being hepatalin-dependent at doses from 10 to 100 mU/kg in rats (Lautt et al., 2001). The background steady state insulin concentration does not affect the response to pulses, as shown by similar RIST indexes in fasted rats during intravenous or intraportal infusions that raised background insulin levels eightfold (Peitl et al., 2009). The RIST

index is the same in the fasted state and in the fed state after atropine although insulin levels are elevated in the fed state (Figure 4).

Insulin release is pulsatile and must be duplicated as such in relevant research.

6.4 THERAPEUTIC MIMICKING OF THE NERVE AND GSH FEEDING SIGNALS: Both the nerve and GSH feeding signals are absent in the diet-induced model of diabetes and can be mimicked by drugs. After 9 weeks of access to a 35% sucrose supplemented drink, rats showed complete absence of hepatalin action. A single pre-meal combination of bethanechol and Nacetylcysteine (BENAC) fully restored hepatic GSH levels and the hepatic cholinergic signal and the ability of a pulse of insulin to stimulate secretion of a pulse of hepatalin. The signals are synergistic and permissive in that they do not result in direct action on glucose uptake in the absence of a pulse of insulin but restore the response to insulin when mimicked by drugs. Doseresponse curves indicate a maximal response for both signals that are not potentiated by higher doses (Lautt et al., 2011).



One Pre-meal Dose of Drugs to Mimic the Nerve (BE) and the Glutathione (NAC) Signals Restores MIS

Figure 5: BENAC synergistic treatment of AMIS. The % MIS (hepatalin/insulin action x 100%) calculated from the RIST index in 24-hour fasted anesthetized rats determined before and 90 minutes after intragastric injection of a mixed liquid meal in a normal control group and in four groups that had been treated with nine weeks of sucrose supplementation. The sucrose diet completely blocked MIS. Bethanechol (BE) produced a modest restoration of MIS in the diabetic model. N-acetyl cysteine (NAC) elevated GSH levels to that seen after feeding but had no effect on the RIST index. The combination of the two feeding signals (BENAC) resulted in a full restoration of MIS. Increasing the doses of either NAC or BE restored MIS to levels seen after feeding but not above. This model demonstrates that in sucrose-induced AMIS, both the nerve and GSH signals were absent but were able to be restored by mimicking these signals with pharmaceuticals. The synergistic permissive nature of the feeding signals is demonstrated. (Figure modified from Lautt et al., 2011, Can. J. Physiol. Pharmacol. 89: 135-142).

6.5 REDUNDANT CONTROL SYSTEMS: Positive redundant control systems in the liver are known. The hyperglycemic response to hemorrhage is maintained in the absence of hepatic sympathetic nerves or adrenal catecholamine secretions. However, absence of both signals prevents the hyperglycemic response (Lautt et al., 1982; Lautt and Cote, 1977). Similarly, the important hepatic vascular compensation for blood loss that occurs by activation of the hepatic capacitance vessels can be regulated in the absence of hepatic sympathetic nerves, adrenaline, noradrenaline, and angiotensin, all of which independently can contract the capacitance vessels. The capacitance response is particularly redundantly controlled in that elimination of the active regulators also results in a greater degree of pressure drop for the same amount of blood loss and therefore the passive contraction of the liver is also increased (Lautt et al., 1980). These are examples of positive hepatic redundant control systems whereby absence of one signal is compensated by another signal to ensure activation of the response.

The regulatory systems controlling hepatalin secretion represent a negative redundant control system whereby absence of either signal blocks the response. The nerve and GSH signals are both required for pulses of insulin to stimulate the secretion of pulses of hepatalin from the liver. These signals are also unique in that they are permissive only. Activation of the feeding signals in the absence of insulin pulses does not result in the secretion of hepatalin (Lautt et al., 2011).

7. ABSENCE OF MEAL-INDUCED INSULIN SENSITIZATION (AMIS):

AMIS results from diminished hepatalin action following a meal. A healthy MIS involves the actions of insulin and hepatalin, with the result being tight glycemic control and storage of nutrient energy primarily as glycogen in muscle. The partitioning and storage of nutrient energy via glucose is essentially complete by 2–3 hours in the healthy state. Acute AMIS results in increased and prolonged postprandial hyperglycemia (Lautt et al., 2011) with compensatory hyperinsulinemia. The elevated insulin acts primarily on the liver and adipose tissue and results

in hyperlipidemia, elevated free radical production, and a shift in nutrient storage from glycogen to fat.

The presence of hepatalin also accounts for the vasodilation in skeletal muscle traditionally associated with insulin (Figure 8). The absence of hepatalin is associated with lack of an increased blood flow response to insulin administration (Ming and Lautt, 2011). Chronic AMIS results in a progressive predictable series of metabolic and cardiovascular dysfunctions that can be increased or decreased by manipulating hepatalin release (Ming et al., 2009; Ming et al., 2011).

A sucrose-supplemented diet for nine weeks resulted in complete elimination of MIS. Intraportal bethanechol (nerve signal) produced a modest restoration of MIS whereas Nacetylcysteine alone (GSH signal) conferred no improvement. The synergistic effect of the two feeding signals is shown by the full restoration of MIS in response to mimicking both feeding signals with one dose prior to the meal (Figure 6). AMIS that occurred with age and was potentiated by a sucrose supplement was attenuated by an antioxidant cocktail (Ming et al., 2009).

8. HEPATALIN AND THE AMIS SYNDROME (AMISS):

The AMISS paradigm proposes that the initiating metabolic defect that progressively leads to type 2 diabetes is an absence of postprandial hepatalin action. Hepatalin is released from the liver in the immediate postprandial state until nutrient energy has been processed, at which time insulin no longer stimulates hepatalin secretion. Postprandially, hepatalin acts in concert with insulin and accounts for the majority of glucose uptake following a meal: 55% in rats (Lautt et al., 2001), 45% in mice (Latour and Chan, 2002), and 67% in humans (Patarrao et al., 2008).

Hepatalin acts selectively on skeletal muscle, kidney, and heart to form glycogen (Fernandes et al., 2011), but not liver, guts, or fat (Xie and Lautt, 1996a).

Absence of hepatalin secretion results in Absence of Meal-induced Insulin Sensitization (AMIS). AMIS for one meal results in elevated postprandial hyperglycemia with compensatory hyperinsulinemia and resultant hyperlipidemia and elevated free radical generation. The vasodilator action of hepatalin is also absent. If AMIS becomes chronic, a predictable sequence of pathologies comprising the AMIS syndrome progresses from prediabetes to obesity and type 2 diabetes. Based on this paradigm of postprandial nutrient energy processing, diagnosis of prediabetes is possible at a much earlier stage than previously available.

The clustering of pathologies and dysfunctions associated with AMIS has previously been given multiple names which simply reflect a clustering of symptoms. These names have included syndrome X, metabolic syndrome, insulin resistance syndrome, dysmetabolic syndrome, plurimetabolic syndrome, cardiometabolic syndrome, dyslipidemic hypertension, hypertriglyceridemic waist, and deadly quartet (Grundy et al., 2005), with the most recent recommendation by the American Diabetes Association to use the term, cardiometabolic risk (Stern and Izkhakov, 2006). However, none of the past names reflect knowledge of the mechanistic basis for the syndrome. Equally importantly, these previous descriptive categorizations do not include the concept of chronological progression. We have proposed the term AMIS Syndrome (AMISS) to define the predictable chronology of pathologies that is initiated by the acute absence of hepatalin action. Chronic AMISS progresses from early-stage prediabetes associated generally with cardiovascular alterations and body fat accumulation, to

dysfunctions in every organ system including heart, blood vessels, kidneys, nerves, eyes,

endocrine organs, and the central nervous system.



Absence of Meal-Induced Insulin Sensitization Syndrome (AMISS)

TIME

Figure 6: The AMIS syndrome is the progressive, predictable, chronological accumulation of signs and symptoms of homeostatic disturbances caused by absence of hepatalin action after each meal. Absence of hepatalin action in the fasted state is appropriate, but if it is not rapidly reversed after a meal, leads to postprandial increases in glucose, insulin, lipids, and reactive oxygen species (ROS). AMISS progresses to prediabetes, obesity, and type 2 diabetes, and associated dysfunctions in major organ systems. By the time fasting glucose is elevated the AMISS is well underway. The specific order of appearance of dysfunctions is an estimate that requires verification. A variety of contributing factors lead to suppression of hepatalin secretion. Treatment options vary with the duration and extent of AMISS. ROS = reactive oxidative species, PP = postprandial. (Figure modified from Lautt et al., 2010, Can. J. Physiol. Pharmacol. 88: 313–323).

8.1 AGE AND AMISS: Ageing is associated with a generalized reduction in

parasympathetic nerve function demonstrated for the cardiovascular system (O'Brien et al.,

1986; Ingall et al., 1990), gastrointestinal tract (Phillips and Powley, 2007), urinary bladder

(Schneider et al., 2005) and eyes (Fitzgerald et al., 2005). Parasympathetic dysfunction can be identified before the development of diabetes (reviewed Hosking et al., 1978) and is later associated with the metabolic syndrome (Britton et al., 2007) and obesity (Skrapari et al., 2007; Von Kanel et al., 2007). Parasympathetic dysfunction in the liver results in the blocking of hepatalin release and resultant AMIS as the earliest stage of AMISS.

The free radical (oxidative stress) theory of ageing was first proposed by Harman in 1956. According to this theory, endogenously produced free radicals continuously and progressively cause permanent DNA and tissue damage, finally leading to ageing. Since many free radicals are produced from the mitochondria, the current version of this theory is the mitochondrial free radical theory of ageing. Oxidative damage to proteins has been found in association with age in a variety of tissues and cells including fibroblasts, brain, liver, heart, and skeletal muscle (reviewed Merker et al., 2001). Free radicals are involved with, or possibly even trigger, the process of ageing (Biesalski, 2002).

Insulin action in rats decreased until nine weeks of age and remained unchanged thereafter, whereas the hepatalin-dependent component decreased from nine weeks of age throughout ageing (Ribeiro et al., 2008). Figure 7 shows the percent of glucose uptake response to insulin that is attributable to HISS action, decreasing dramatically with age. Sucrose supplementation increased AMIS and accelerated AMISS whereas SAMEC (a combination of antioxidants as described in Section 8.2) protected against both age- and diet-induced AMIS (Ming et al., 2009).

8.2 ANTIOXIDANTS AND AMISS: The effects of antioxidants or free radical scavengers have been widely tested for the prevention and treatment of acute and chronic diseases.

However, the efficacy of pharmaceutical antioxidant treatment in clinical trials has been generally unimpressive. Although the production of free radicals with various chemical properties is widely spread out throughout the different tissues and cellular components, the chemical property of an individual antioxidant can only allow it to scavenge the free radicals located in a limited cellular component, for example, the lipid or aqueous phase.



Age, Sugar, and SAMEC Ageing results in AMIS, made worse by sugar supplement;

Figure 7: The RIST index (amount of glucose required to maintain a glycemic baseline in response to a 5 min pulse of 50 mU/kg insulin) was determined in the fed state before (insulin plus hepatalin action) and after atropine (insulin action only), to determine the relative glucose uptake response to insulin that is attributable to insulin or hepatalin. The ratio of insulin and hepatalin in partitioning of storage of nutrient energy was manipulated by interacting normal ageing (9, 26, 52 weeks) with sucrose-induced AMIS, and the effect of a synergistic antioxidant combination (SAMEC = S-adenosyl-L-methionine, vitamin E, and vitamin C) on both conditions. Normal ageing resulted in a severe decline in hepatalin action that was strongly protected by SAMEC. A 5% sucrose water supplement potentiated the age-related decline, which was also protected by SAMEC. The direct effect of insulin in the absence of hepatalin action remained unchanged in these conditions. The variation in response to insulin can be attributed to hepatalin action. This model allowed monitoring the development of components of the AMIS syndrome. (Figure modified from Ming et al., 2009 Can. J. Physiol. Pharmacol. 87: 873–882).

We have reported a unique synergistic interaction among three antioxidants selected to specifically target different cellular components: the combination of S-adenosyl-L-methionine and vitamins E and C (SAMEC). All three components play an important and different but interacting role in scavenging free radicals and in cell health. The water-soluble property of vitamin C makes it the first order antioxidant to protect cell components from free radicalinduced damage by quenching various water-soluble radicals, for example, superoxide anion. Vitamin E is a lipid soluble molecule and can transfer its phenolic hydrogen to a peroxyl free radical of a peroxidized polyunsaturated fatty acid, thereby breaking the radical chain reaction, and thus preventing the lipid peroxidation in cellular and subcellular membrane phospholipids especially those of mitochondria and microsomes. S-adenosyl-L-methionine is a natural regulator of glutathione (GSH) and is involved with transmethylation and sulfation reactions. GSH is the main intracellular defence against free radicals. GSH levels are significantly depleted in liver injury induced by oxidative stress. Administration of S-adenosyl-L-methionine is an effective means of restoring intracellular GSH stores in mitochondria, thus improving the cellular ability to scavenge free radicals (Lieber, 1999).

The unique synergistic activity of SAMEC was demonstrated by the absence of any beneficial effect of vitamins E and C taken together or by S-adenosyl-L-methionine administered separately. In contrast, when administered together, the three conferred very significant synergistic protection against liver damage and the development of hepatalindependent insulin resistance that results from acute exposure to the free radical generating hepatotoxin, thioacetamide (Ming at al., 2006).

8.3 EXERCISE AND AMISS: Voluntary exercise using a running wheel for seven days led to an improvement in the response to insulin in both healthy, sucrose-induced, and high fat dietinduced prediabetic rats (Chowdhury et al., 2013c). The improvement was attributable to an increase in hepatalin action with no significant improvement in direct insulin action. The AMIS that was diet-induced was significantly reversed by seven days of access to a running wheel (Chowdhury et al., 2013a). Hepatalin action correlated strongly with the distance run. The animals that benefited the most per distance run had the lowest pre-training hepatalin action (Chowdhury et al., 2011). Antioxidant supplementation with SAMEC did not impair or improve the beneficial effect of exercise on AMIS. Clinical testing of the effect of exercise on AMIS has not been done.

8.4 DIET AND AMISS: Rat chow supplemented with sucrose (Ribeiro et al., 2005; Lautt et al., 2011) or lipids (Afonso et al., 2007, 2010) resulted in a reduction of MIS that is attributable to absence of the feeding signals. A limited low dose of sucrose supplementation (50 ml of 5% solution shared with two cage mates) resulted in a progression of the AMIS syndrome with hepatalin action being essentially absent after 52 weeks (Ming et al., 2009). A 35% sucrose solution, in contrast, results in complete AMIS after only two weeks with accumulation in body weight being seen by six weeks (Ribeiro et al., 2005). The sucrose-induced prediabetic state was prevented by daily supplementation with SAMEC (Ming et al., 2009).

Diet also affected gestational nutrient partitioning in rats by affecting hepatalin action. A sucrose supplement shifted the balance of hepatalin and insulin so that fat accumulation occurred early during gestation (Lovat et al., 2021). Antioxidant supplementation (SAMEC) prevented the impact of the sugar diet on hepatalin and obesity (Lovat and Lautt, unpublished).

8.5 STRESS AND AMISS: Acutely stressful physical or psychological events are often associated with development of diabetes. The link between stress and AMIS is unclear and possibly mediated by several independent mechanisms.

8.5.1 ACUTE STRESS AND AMISS: As with all factors that can trigger AMIS, the acute response to stress, such as blood loss, is overlaid by other homeostatic adjustments. In response to blood loss, the hepatic sympathetic nerves and adrenal catecholamines activate glycogen breakdown in the liver that can double glucose levels within minutes (Lautt et al., 1982). The secretion of insulin in response to the hyperglycemia is blocked by somatostatin secreted by the pancreas (Lautt et al., 1983).

Acute blood loss leads to complete AMIS that was not prevented by blocking either the alpha- or beta-adrenergic receptors. However, the somatostatin receptor antagonist, cyclosomatostatin, completely blocked the development of hemorrhage-induced AMIS. Exogeneous somatostatin produced AMIS that was prevented or reversed by blocking somatostatin receptors (Seredycz and Lautt, 2006). The hemorrhage experiments were done under full anesthesia, yet the stress of extensive surgical intervention does not activate these responses as shown by the RIST index in response to intragastric injection of a liquid test meal being similar in conscious unrestrained or anesthetized rats (Sadri et al., 2006)

8.5.2 PSYCHOLOGICAL STRESS AND AMISS: Mental stress suppresses MIS, which is an appropriate response preserving blood glucose levels for emergency use in a fight or flight situation. The relation between stress and metabolic dysfunction was defined as a component of the General Adaptation Syndrome by the Canadian physiologist, Hans Selye (1950), primarily focusing on the pituitary and adrenal hormonal roles. Diabetes and change in body composition

were components. Similar to the general adaptation syndrome, the acute induction of AMIS is useful in the immediate fight or flight situation. However, chronically maintained, both responses will lead to pathologies. Personal observation revealed that rats experiencing loud noises showed behavioral evidence of anxiety, and consistently resulted in the rats having AMIS. Chronic inflammation was also associated with AMIS.

8.6 GLUCAGON AND AMISS: A cAMP (cyclic adenosine monophosphate) analog or glucagon produced a dose-dependent AMIS that correlated with decreased hepatic GSH content and was reversed by administration of a GSH donor. Neither hepatic nor plasma nitric oxide levels were affected, suggesting that only the GSH feeding signal had been impacted (Patarrao et al., 2015). Glucagon levels are increased during normal fasting and diabetes; and may account for the reduced GSH levels seen in the fasted and diabetic state.

8.7 HEART FUNCTION AND AMISS: Parasympathetic dysfunction leads to AMIS. Cardiac parasympathetic dysfunction is seen in early-stage prediabetes as a reduced heart rate variability coupled with the respiratory cycle. In health, inhalation and expiration result in increases and decreases in heart rate that is mediated by parasympathetic nerves. This variability is seen in young and healthy subjects and is absent in the elderly and diabetics (Svensson et al., 2016; Benichou et al., 2018; Hoshi et al., 2019).

Cardiac performance in rats was evaluated using a Millar pressure volume conductance catheter system. Cardiac dysfunctions and AMIS developed with age and were increased by sucrose supplementation and inhibited by the antioxidant (SAMEC) treatment. Cardiac dysfunction correlated with AMIS and showed as reduction in cardiac index, heart rate, maximal rate of contraction and relaxation, reduced ejection fraction, and increased left

ventricular end diastolic pressure. Total peripheral vascular resistance also increased. These dysfunctions occurred prior to accumulation of cardiac lipids (Ming et al., 2011).

8.8 VASCULATURE AND AMISS: Hindlimb blood flow and the metabolic response to insulin were tightly coupled and attributable to hepatalin action. Severing the hepatic nerves resulted in a loss of hepatalin action as demonstrated by reduced hindlimb glucose uptake and blood flow in response to a pulse of insulin. Intraportal infusion of acetylcholine to mimic the permissive nerve signal, fully restored the metabolic and vascular response to a pulse of insulin (Ming and Lautt, 2011). Hepatalin action is seen only in the fed state, accounting for much of the debate as to the existence of a vascular response to insulin.



Hepatalin Action on Blood Flow

Figure 8: The effect of a pulse of insulin quantified by the RIST showing the combined action of insulin (black) and hepatalin (orange) in the fed state on hindlimb blood flow and glucose uptake, and inhibition of both effects by atropine. In contrast, in the 24-hour fasted rats,

hepatalin secretion was already physiologically suppressed and therefore atropine had no additional effect on insulin action. In the same study, not shown in the figure, denervation of the liver eliminated the vascular and metabolic action of hepatalin, which was restored by intraportal infusion of acetylcholine. (Data from Ming and Lautt, 2011, J. Appl. Physiol. 110: 60–68).

Myocardial blood flow was similar in fasted diabetic and healthy subjects but increased significantly in response to a mixed meal in the control subjects but did not in the diabetic patients (Scognamiglio et al., 2005). The superiority of a mixed meal test verses a pure sugar test was shown by the inability of the oral glucose tolerance test to elicit increased cardiac blood flow that was seen in response to a mixed meal. Oral glucose does not activate the feeding signals and does not result in hepatalin secretion (Sadri et al., 2006; Afonso et al., 2016).

8.9 OBESITY AND AMISS: It is widely suggested that adiposity causes insulin resistance and type 2 diabetes. But guidance from this new paradigm suggests instead that reduced or absent hepatalin action leads to adiposity.

The glucose uptake response to a bolus of insulin in the fasted state is potentiated after feeding by 232% in humans (Patarrao et al., 2008). Meal-induced Insulin Sensitization was significantly less in subjects with a mild degree of adiposity (466 mg/kg glucose uptake, BMI 22.7 verses 211 mg/kg glucose in the overweight group, BMI 27.7). The glucose uptake in response to insulin was less in the overweight group although the insulin concentration profiles were not different (Patarrao et al., 2012). The difference was attributable to reduced hepatalin action, equivalent to very mild AMIS.

Adiposity does not precede insulin resistance. Rather, absence of hepatalin action leads to adiposity. Several studies show a correlation of reduced hepatalin action and generalized obesity (reviewed in Lautt and Wang, 2014a).

Postprandial elevations in glucose, insulin and triglycerides are a first sign of AMIS. Blood glucose levels may not change significantly in early AMISS due to compensatory insulinemia. Complete AMIS results after two weeks of a 35% sucrose supplemented diet, whereas adiposity was detected only after 6 weeks (Ribeiro et al., 2005), suggesting that adiposity follows, not precedes AMIS. Liver and plasma levels of lipids were elevated in the 2week sucrose model of diabetes (Martins et al., 2016).

To test the hypothesis that AMIS precedes obesity, several chronic protocols known to have a predictable effect on hepatalin action were used to produce chronic graded slow onset of degrees of impairment of MIS related to age; a chronic sucrose supplement to potentiate AMIS (Ribeiro et al., 2005, 2008; Lautt et al., 2008); and an antioxidant cocktail to attenuate AMIS (Lautt et al., 2008; Ming et al., 2006). By both hastening and attenuating the rate of decline in MIS, the intention was to produce a sufficient range of dysfunctions to permit interpretable correlations. Adiposity increased as hepatalin action decreased. The strong correlation is similar if adiposity is assessed by individual fat pads, or whole-body adiposity determined by bioelectric impedance (Ming et al., 2009)).

Hepatalin and Adiposity

Normal Diet



Figure 9: The relation between hepatalin-dependent insulin action on percent whole body fat mass calculated from bio electrical impedance, and from total weighed fat pad masses as percent of body weight in rats at 9, 26, and 52 weeks of age. Hepatalin action is strongly and negatively correlated with both indices of adiposity. The correlation remains similar in the control groups on normal diet, the 5% sucrose supplemented groups, and the groups on normal and sucrose diets that had been chronically treated with SAMEC (S-adenosyl-L-methionine, vitamin C, and vitamin E). The data points move along the regression line with lower hepatalin action and increased fat moving to the left AMIS) and the data from the treated group with improved hepatalin action moving to the right. (Figure modified from Ming et al., 2009, Can. J. Physiol. Pharmacol. 87: 873–882).

Petersen et al. (2007) showed that insulin resistance in lean subjects was associated with a 60% increase in plasma triglyceride and reduction of muscle glycogen synthesis by 60% that occurred prior to the development of abdominal obesity

The data show that AMIS leads to adiposity. However, the effect of adiposity resulting from overeating on AMIS is yet unknown. Chronic accumulated fat stores and increased lipid metabolism is inflammatory (Sun et al., 2014) which may suppress hepatalin secretion.

8.10 SEX AND AMISS: To say that the physiology of the human male has more similarity to that of lab rats than it does to that of the human female is perhaps overstated, but the lifelong shifting milieu of female hormones must be assumed to impact nutrient partitioning.

8.10.1 FETAL ALCOHOL EFFECT AND AMISS: The metabolism of offspring is affected by maternal alcohol exposure (Taylor et al., 1982; Kelly et al., 1991; Weinberg, 1992). The RIST index and % hepatalin contribution are similar in male and virgin female rats (Ribeiro et al., 2002). Pregnant rats fed alcohol develop AMIS with normal insulin action but impaired hepatalin action, and the offspring are pre-diabetic in young adulthood depending on the dose of maternal alcohol (Minuk et al., 1998; Sadri et al., 2003). The female offspring are more severely impacted by the fetal alcohol exposure (Sadri et al., 2005).

8.10.2 GESTATION AND AMISS: Maternal obesity is among the most prevalent risk factors in pregnancy (Ladyman et al., 2010). It is associated with macrosomia, delivery complications, neural tube defects, as well as increased perinatal mortality (Begum et al., 2011; Ladyman et al., 2010; Nelson et al., 2010; Olson and Blackwell, 2011). Macrosomic infants have significantly more adipose tissue, with maternal hyperinsulinemia fueling a large portion of this excess fetal growth (Ferraro et al., 2012). Cord blood triglycerides are elevated, reflecting

enhanced hepatic and adipose tissue fat synthesis in fetal tissues (Zhu et al., 2010). In the mother, obesity and insulin resistance result in increased risks for preeclampsia, hypertension, and diabetes (Begum et al., 2011; Levin, 2006; Olson et al., 2003).

The relative predominance of hepatalin and insulin shift during gestation with hepatalin action being more critical in early gestation when muscle and tissue growth are maximal. By the latter part of gestation, the nutrient partitioning shifts away from glycogen formation to lipid metabolism and fat accumulation in both the fetus and mother, a condition similar to prediabetes but of value in storing lipid reserves for both mother and the newborn during the postpartum period (Lovat et al., 2020). A sucrose-supplemented diet shifted the timing of the appearance of AMIS during gestation and resulted in increased and earlier postprandial elevations in insulin and lipids and accumulated adiposity (Lovat et al., 2021).

The timing of the shift from hepatalin to insulin dominance determines the degree of accumulated adiposity and components of AMISS over the gestation term.

8.10.3 HUMAN FEMALE DATA: At this point there are no clinical studies directly related to MIS and hepatalin action that would allow assessment of the changes in nutrient partitioning or hepatalin action that occur throughout the complex hormonal life of the human female. Females experience puberty very differently from males. The monthly hormonal changes that occur thereafter in females involve a flux of hormones that individually have metabolic impact. The role of hepatalin versus insulin throughout this cycle is unknown.

Although the obvious change in body composition in elderly humans from muscle to fat is consistent with the animal data, the effect of age on AMIS has not been specifically verified in humans, and female ageing is also made more complex by the appearance of menopause. To

complicate matters further, the effect of hormonal contraceptives and hormone replacement therapy on hepatalin action and nutrient partitioning has not been examined. 9. IMPACTS OF THE NEW PARADIGM:

9.1 ACCELERATED KNOWLEDGE: Before settling on the name 'insulin'—followed by its formal introduction in May of 1922 by Banting, Best, Collip and McLeod—the hormone had been previously named insuline and iletin, (insuline p126, iletin p147 in Bliss, 1984). But the structure of insulin was not described until a half century later (Sanger, 1959). If hepatalin is the missing link in our understanding of prediabetes, obesity, and type 2 diabetes, the discovery of new tools through focusing on the new paradigm should accelerate the growth of knowledge in this area at a much faster rate than it has developed over the past 100 years.

9.2 NEW PARADIGM-BASED PRODUCTS: The focus on nutrient partitioning and the role of hepatalin has resulted in basic science tools being developed into health-promoting products:

a) SciMar NuPa Daily[®], a version of SAMEC (Section 8.2., Figure 7) has been approved as a nutraceutical preventative and is available online.
b) SciMar NuPa Test[®], a mixed reconstituted powdered test meal calibrated to body weight, has completed clinical evaluation. Full postprandial profiles of glucose, insulin, triglycerides, and a calculated scoring index based on meal-induced insulinemia and glycemia (MIG score) will soon be published.
c) SciMar NuPa Renew[®] is a diagnostic with timing and dosing clinical trials soon to be underway. The diagnostic can be validated clinically on an individual basis with two meal tests, one with placebo and one with SciMar NuPa Renew[®] given prior to consumption of the test meal. Only those showing a reduced MIG score

(a score combining glucose and insulin levels after a meal) after the SciMar NuPa Renew would be appropriate for chronic pre-meal therapeutics (a theranostic).

9.3 FOCUS ON LIFESTYLE: Over the last few decades, the diabetes literature has begun to focus on the impact of lifestyle changes such as weight loss, exercise, and healthy diet. But reliance on the old metabolic syndrome paradigm has prevented a clear understanding of exactly how lifestyle impacts diabetic pathologies. The focus of the new paradigm on AMIS, AMISS, and the role of hepatalin in nutrient partitioning has allowed the development of tools to quantitatively evaluate the impact of changing one's lifestyle and other therapeutic interventions.

As one example, The Wellness Transformation Network (WTN)

(wellnesstransformation.org) has been developed by SciMar (see Disclosure) as a not-for-profit community-based program supported by and involving industry, government, wellness groups, First Nations, and academic scientists. The objective is to encourage but also evaluate lifestyle wellness approaches related to prediabetes, obesity, and type 2 diabetes. As part of the process of evaluation, a full clinical screening procedure is being developed to examine a wide range of components of the AMIS syndrome including: the effect of the test meal assessed from the MIG score (combined insulin and glucose induced by the meal); total body regional fat, muscle, and water content; parasympathetic health of eyes (iris response to light) and heart (heart rate variability with the respiratory cycle); airway resistance (preclinical data indicate increased airway resistance associated with AMIS); retinal vascular health, muscle strength, and standard blood assays including HbA1c, and a lifestyle survey. The impact of various lifestyle and other

interventions on ameliorating the initiating dysfunction (AMIS) and progressive pathologies (AMISS) will be able to be tested.

This new paradigm provides the scientific basis for individuals and groups to make quantitatively evaluated wellness choices, and to take control of their own health.

Disclosure: Lautt is co-founder and Chief Scientific Officer of SciMar, a Canadian biotechnology company (SciMar.ca) that has developed clinical products based on the hepatalin paradigm.

Acknowledgments: The science described in this review directly related to hepatalin and MIS was a gradual accumulation (1991–2021) of pieces in the missing link contributed to by undergraduate, graduate, and postdoctoral students in Winnipeg, Canada (Lautt et al.) and Lisbon, Portugal (Macedo et al.), and visiting scientists from Japan, China, Ukraine, and Portugal, and several technologists trained by my long-term collaborator, Dallas Legare. The science is being brought to market by Mick Lautt and John West. Manuscript was prepared by Wayne Lautt, with assistance from Melanie Lautt and Karen Sanders.

REFERENCES

Afonso, R.A., Gaspar, J.M., Lamarao, I., Lautt, W.W., Macedo, M.P., 2016. Postprandial insulin action relies on meal composition and hepatic parasympathetics: dependency on glucose and amino acids: Meal, parasympathetics and insulin action. J. Nutr. Biochem. 27, 70–78. https://doi.org/10.1016/J.jnutbio.2015.08.023.

Afonso, R.A., Lautt, W.W., Ribeiro, R.T., Legare, D.J., Macedo, M.P., 2007. Insulin resistance in two animal models of obesity: a comparison of HISS-dependent and HISS-independent insulin action in high-fat diet-fed and Zucker rats. Proc. West. Pharmacol. Soc. 50, 110–114. PMID: 18605245.

Afonso, R.A., Lautt, W.W., Schafer, J., Legare, D.J., Oliveira, A.G., Macedo, M.P., 2010. High-fat diet results in postprandial insulin resistance that involves parasympathetic dysfunction. Br. J. Nutr. 104, 1450–1459. https://doi.org/10.1017/S0007114510002400.

Anderson, R.A., Evans, M.L., Ellis, G.R., Graham, J., Morris, K., Jackson, S.K., Lewis, M.J., Rees, A., Frenneaux, M.P., 2001. The relationship between post-prandial lipaemia, endothelial function and oxidative stress in healthy individuals and patients with type 2 diabetes. Atherosclerosis 154, 475–483. https://doi.org/10.1016/s0021-9150(00)00499-8.

Begum, K.S., Sachchithanantham, K., De Somsubhra, S., 2011. Maternal obesity and pregnancy outcome. Clin. Exp. Obstet. Gynecol. 38(1), 14–20. PMID: 21485717.

Benichou, T., Pereira, B., Mermillod, M., Tauveron, I., Pfabigan, D., Maqdasy, S., Dutheil, P., 2018. Heart rate variability in type 2 diabetes mellitus: a systemic review and meta-analysis. PLOS ONE 13(4), e0195166. https://doi.org/10.1371/journal.pone.0195166.

Biesalski, H.K., 2002. Free radical theory of aging. Curr. Opin. Clin. Metab. Care 5, 5–10. https://doi.org/10.1097/00075197-200201000-00002.

Bliss, M., 1984. The discovery of insulin. McClelland and Stewart Ltd., The University of Chicago Press.

Britton, A., Shipley, M., Malik, M., Hnatkova, K., Hemingway, H., Marmot, M., 2007. Changes in heart rate and heart rate variability over time in middle-aged men and women in the general population (from the Whitehall II Cohort Study). Am. J. Cardiol. 100, 524–527. https://doi.org/10.1016/j.amjcard.2007.03.056.

Brownlee, M., 2001. Biochemistry and molecular cell biology of diabetic complications. Nature 414, 813–819. htpps://doi.org/10.1038/414813a.

Ceriello, A., 2000. The post-prandial state and cardiovascular disease: relevance to diabetes mellitus. Diabetes Metab. Res. Rev. 16, 125–132. https://doi.org/10.1002/(sici)1520-7560(200003/04)16:2<125::aid-dmrr90>3.0.co;2-4.

Ceriello, A., Bortolotti, N., Motz, E., Pieri, C., Marra, M., Tonutti, L., Lizzio, S., Feletto, F., Catone, B., Taboga, C., 1999. Meal-induced oxidative stress and low-density lipoprotein (LDL) oxidation in diabetes: the possible role of hyperglycemia. Metabolism 48, 1503–1508. https://doi.org/10.1016/s0026-0495(99)90237-8. 10.1016/s0026-0495(99)90237-8.

Ceriello, A., Giugliano, D., Quatraro, A., Dello Russo, P., Torella, R., 1988. Blood glucose may condition factor VII levels in diabetic and normal subjects. Diabetologia 31, 889–891. https://doi.org/10.1007/BF00265372.

Ceriello, A., Taboga, C., Tonutti, L., Giacomello, R., Stel, L., Mortz, E., Pirisi, M., 1996. Post-meal coagulation activation in diabetes mellitus: the effect of Acarbose. Diabetologia 39, 469–473. https://doi.org/10.1007/BF00400679.

Chou, H.F., McGivern, R., Berman, N., Ipp, E., 1991. Oscillations of circulating plasma insulin concentrations in the rat. Life Sci. 48, 1463–1469. https://doi.org/10.1016/0024-3205(91)90183-c. 10.1016/0024-3205(91)90183-c.

Chowdhury, K.K., Legare, D.J., Lautt, W.W., 2011. Insulin sensitization by voluntary exercise in aging rats is mediated through hepatic insulin sensitizing substance (HISS). Exp. Gerontol. 46(1), 73–80. https://doi.org/10.1016/j.exger.2010.10.006. 10.1016/j.exger.2010.10.006.

Chowdhury, K.K., Legare, D.J., Lautt, W.W., 2013a. Exercise enhancement of hepatic insulin sensitizing substance-mediated glucose uptake in diet-induced pre-diabetic rats. Br. J. Nutr. 109(5), 844–852. https://doi.org/10.1017/S000711451200226710.1017/S0007114512002267.

Chowdhury, K.K., Legare, D.J., Lautt, W.W., 2013b. Review Article: Lifestyle impact on mealinduced insulin sensitization (MIS) in health and prediabetes: a focus on diet, antioxidants, and exercise interventions. Can. J. Physiol. Pharmacol. 91(2), 91–100. https://doi.org/10.1139/cjpp-2012-0228. 10.1139/cjpp-2012-0228.

Chowdhury, K.K., Legare, D.J., Lautt, W.W., 2013c. Interaction of antioxidants and exercise on insulin sensitivity in healthy and prediabetic rats. Can. J. Physiol. Pharmacol. 91(7), 570–577. https://doi.org/10.1139/cjpp-2012-0425. 10.1139/cjpp-2012-0425.

DECODE Study Group, on behalf of the European Diabetes Epidemiology Group, 1999. Glucose tolerance and mortality comparison of WHO and American Diabetes Association diagnostic criteria. Lancet 354, 617–621. PMID: 10466661.

De Vegt, F., Dekker, J.M., Ruhé, H.G., Stehouwer, C.D.A., Nijpels, G., Bouter, L.M., Heine, R.J., 1999. Hyperglycaemia is associated with all-cause and cardiovascular mortality in the Hoorn population: the Hoorn study. Diabetologia 42, 926–931. https://doi.org/10.1007/s001250051249. 10.1007/s001250051249.

Engelgau, M.M., Narayan, K.M.V., Herman, W.H., 2000. Screening for type 2 diabetes. Diabetes Care 23, 1563–1580. https://doi.org/10.2337/diacare.23.10.1563. 10.2337/diacare.23.10.1563.

Fernandes, A.B., Patarrao, R.S., Videira, P.A., Macedo, M.P., 2011. Understanding postprandial glucose clearance by peripheral organs: The role of the hepatic parasympathetic system. J. Neuroendocrinol. 23, 1288–1295. https://doi.org/10.1111/j.1365-2826.2011.02226.x. 10.1111/j.1365-2826.2011.02226.x.

Ferraro, Z.M., Barrowman, N., Prud'homme, D., Walker, M., Wen, S.W., Rodger, M., Adamo, K.B., 2012. Excessive gestational weight gain predicts large for gestational age neonates independent of maternal body mass index. J. Matern. Fetal. Neonatal. Med. 25(5), 538–542. https://doi.org/10.3109/14767058.2011.638953.

Fitzgerald, M.E., Tolley, E., Jackson, B., Zagvazdin, Y.S., Cuthbertson, S.L., Hodos, W., Reiner, A., 2005. Anatomical and functional evidence for progressive age-related decline in parasympathetic control of choroidal blood flow in pigeons. Exp. Eye Res. 81, 478–491. https://doi.org/10.1016/j.exer.2005.03.008. 10.1016/j.exer.2005.03.008.

Guarino, M.P., Correia, N.C., Lautt, W.W., Macedo, M.P., 2004. Insulin sensitivity is mediated by the activation of the Ach/NO/cGMP pathway in rat liver. Am. J. Physiol. Gastrointest. Liver Physiol. 287, G527–G532. https://doi.org/10.1152/ajpgi.00085.2004. 10.1152/ajpgi.00085.2004.

Guarino, M.P., Correia, N.C., Raposo, J.F., Macedo M.P., 2001. Nitric oxide synthase inhibition decreases output of hepatic insulin sensitizing substance (HISS), which is reversed by SIN-1 but not by nitroprusside. Proc West Pharmacol Soc 44, 25–26. PMID: 11793984.

Goodner, C.J., Walike, B.C., Koerker, D.J., Ensinck, J.W., Brown, A.C., Chideckel, E.W., Palmer, J., Kalnasy, L., 1977. Insulin, glucagon, and glucose exhibit synchronous, sustained oscillations in fasting monkeys. Science 195, 177–179. https://doi.org/10.1126/science.401543.

Greenway, C.V., Lautt, W.W., 1979. Effects of infusions of catecholamines, angiotensin, vasopressin and histamine on hepatic blood volume in the anaesthetized cat. Br. J. Pharmacol. 44, 177–185.

Grundy, S.M., Cleeman, J.I., Daniels, S.R., Donato, K.A., Eckel, R.H., Franklin, B.A., Gordon, D.J., Krauss, R.M., Savage, P.J., Smith Jr., S.C., Spertus, J.A., Costa, F., 2005. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. Circulation 112(17), 2735–2752. https://doi.org/10.1161/CIRCULATIONAHA.105.169404.

Hanefeld, M., Fischer, S., Julius, U., Schulze, J., Schwanebeck, U., Schmechel, H., Ziegelasch, J.J., Lindner, J., 1996. Risk factors for myocardial infarction and death in newly detected NIDDM: the Diabetes Intervention Study, 11-year follow-up. Diabetologia 39, 1577–1583. https://doi.org/10.1007/s001250050617.

Harman, D., 1956. Aging: A theory based on free radical and radiation chemistry. J. Gerontol. 11, 298–300. https://doi.org/10.1093/geronj/11.3.298.

Hogg, N., 2002. The biochemistry and physiology of S-nitrosothiols. Ann. Rev. Pharmacol. Toxicol. 42, 585–600. https://doi.org/10.1146/annurev.pharmtox.42.092501.104328.

Hoshi, R.A., Santos, I.S., Dantas, E.M., Andreao, R.V., Schmidt, M.I., Duncan, B.B., Mill, J.G., Lotufo, P.A., Bensenor, I., 2019. Decreased heart rate variability as a predictor for diabetes - A prospective study of the Brazilian longitudinal study of adult health. Diabetes Metab. Res. Rev. 35, e3175. https://doi.org/10.1002/dmrr.4175.

Hosking, D.J., Bennett, T., Hampton, J.R., 1978. Diabetic autonomic neuropathy. Diabetes 27, 1043–1054. https://doi.org/10.2337/diab.27.10.1043.

Ingall, T.J., McLeod, J.G., O'Brien, P.C., 1990. The effect of ageing on autonomic nervous system function. Aust. NZ J. Med. 20, 570–577. https://doi.org/10.1111/j.1445-5994.1990.tb01315.x.

Ingelsson, E., Sundström, J., Arnlöv, J., Zethelius, B., Lindet L., 2005. Insulin resistance and risk of congestive heart failure. JAMA 294, 334–341. https://doi.org/10.1001/jama.294.3.334.

International Diabetes Federation. IDF Diabetes Atlas, 9th edn. Brussels, Belgium: 2019. Available at: https://www.diabetesatlas.org

Kelly, S.J., Mahoney, J.C., Randich, A., West, J.R., 1991. Indices of stress in rats: effects of sex, perinatal alcohol and artificial rearing. Physiol. Behav. 49, 751–756. https://doi.org/10.1016/0031-9384(91)90314-e.

Ladyman, S.R., Augustine, R.A., Grattan, D.R., 2010. Hormone interactions regulating energy balance during pregnancy. J. Neuroendocrinol. 22(7), 805–817. https://doi.org/10.1111/j.1365-2826.2010.02017.x.

Lang, D.A., Matthews, D.R., Peto, J., Turner, R.C. 1979. Cyclic oscillations of basal plasma glucose and insulin concentrations in human beings. N. Engl. J. Med. 301, 1023–1027. https://doi.org/10.1056/NEJM197911083011903.

Latour, M.G., Chan, C.C. 2002. A rapid insulin sensitivity test (RIST) in the anesthetized mice. Diabetes, 51(Suppl 2): A422.

Latour, M.G., Lautt, W.W., 2002a. The hepatic vagus nerve in the control of insulin sensitivity in the rat. Autonomic Neurosci. 95, 125–130. https://doi.org/10.1016/s1566-0702(01)00390-3.

Latour, M.G., Lautt, W.W., 2002b. Insulin sensitivity regulated by feeding in the conscious unrestrained rat. Can. J. Physiol. Pharmacol. 80, 8–12. https://doi.org/10.1139/y01-094.

Lautt, W.W., 1980. Hepatic parasympathetic neuropathy as cause of maturity-onset diabetes? Gen. Pharmac. 11, 343–345. https://doi.org/10.1016/0306-3623(80)90096-8.

Lautt, W.W., 1998. State of the Art 1997: Hepatic parasympathetic nerves and glucose metabolism, in: Haussinger, D, Jungermann, K. (Eds.), Liver and Nervous System. Falk Symposium No. 103. Kluwer Academic Publishers, UK, pp. 1–14.

Lautt, W.W., 1999. The HISS story overview: A novel hepatic neurohumoral regulation of peripheral insulin sensitivity in health and diabetes. Can. J. Physiol. Pharmacol. 77(8), 553–562. PMID: 10543718.

Lautt, W.W., 2003. Practice and principles of pharmacodynamic determination of HISSdependent and HISS-independent insulin action: methods to quantitate mechanisms of insulin resistance. Med. Res. Rev. 23(1), 1–14. https://doi.org/10.1002/med.10022.

Lautt, W.W., 2007. Postprandial insulin resistance as an early predictive cardiovascular risk factor. Therap. Clin. Risk Managem. 3, 761–770. PMID: 18473001.

Lautt, W.W., Brown, L.C., Durham, J.S., 1980. Active and passive control of hepatic blood volume responses to hemorrhage at normal and raised hepatic venous pressure in cats. Can. J. Physiol. Pharmacol. 58, 1049–1057. https://doi.org/10.1139/y80-158.

Lautt, W.W, Cote, M.G., 1977. The effect of 6-hydroxydopamine-induced hepatic sympathectomy on the early hyperglycemic response to surgical trauma under anesthesia. J. Trauma 17, 270–274. https://doi.org/10.1097/00005373-197704000-00002.

Lautt, W.W., Dwan, P.D., Singh, R.R., 1982. Control of the hyperglycemic response to hemorrhage in cats. Can. J. Physiol. Pharmacol. 60, 1618–1623. https://doi.org/10.1139/y82-239.

Lautt, W.W., Legare, D.J., Martens, E.S., 1983. A possible role for somatostatin in depression of insulin and glucagon levels during hemorrhage. Can. J. Physiol. Pharmacol. 61, 237–240. https://doi.org/10.1139/y83-036.

Lautt, W.W., Legare, D.J., Reid, M.A.G., Sadri, P., Ting, J.W., Prieditis, H., 2005. Alcohol suppresses meal-induced insulin sensitization (MIS). Metab. Synd. Rel. Disord. 3, 51–59. https://doi.org/10.1089/met.2005.3.51.

Lautt, W.W., Macedo, M.P., Sadri, P., Takayama, S., Ramos, F.D., Legare, D.J., 2001. Hepatic parasympathetic (HISS) control of insulin sensitivity determined by feeding and fasting. Am. J. Physiol. Gastrointest. Liver Physiol. 281, G29–G36. https://doi.org/10.1152/ajpgi.2001.281.1.G29.

Lautt, W.W., Ming, Z., Legare, D.J., 2010. Attenuation of age- and sucrose-induced insulin resistance and syndrome X by a synergistic antioxidant cocktail: the AMIS syndrome and HISS hypothesis. Can. J. Physiol. Pharmacol. 88(3), 313–323. https://doi.org/10.1139/Y09-130.

Lautt, W.W., Ming, Z., Legare, D.J., Chowdhury, K.K., Hatch, G.M., Wang, H.H., 2015. Fatty Liver and Fatty Heart –Where do they stand in the AMIS syndrome? Healthcare 3, 666–682. https://doi.org/10/3390/healthcare3030666.

Lautt, W.W., Ming, Z., Macedo, M.P., Legare, D.J., 2008. HISS-dependent insulin resistance (HDIR) in aged rats is associated with adiposity, progresses to syndrome X, and is attenuated by a unique antioxidant cocktail. Exp. Gerontol. 43, 790–800. https://doi.org/10.1016/j.exger.2008.04.013.

Lautt, W.W., Schafer, J., Macedo, M.P., Legare, D.J., 2011. Bethanechol and n-acetylcysteine mimic feeding signals and reverse insulin resistance in fasted and sucrose-induced diabetic rats. Can. J. Physiol. Pharmacol. 89, 135–142. https://doi.org/10.1139/y11-001.

Lautt, W.W., Wang, H.H., 2014a. Obesity as an early symptom of the AMIS syndrome. J. Clin. Med. 3, 1178–1198. https://doi.org/10.3390/jcm3041178.

Lautt, W.W., Wang, H.H., 2014b. EP2803362A1. Hepatic insulin sensitizing substance and test meal for insulin sensitization. European Patent Application #14168369.8, published 19.11.2014, bulletin 2014/47.

Lautt, W.W., Wang, X., Sadri, P., Legare, D.J., Macedo, M.P., 1998. Rapid insulin sensitivity test (RIST). Can. J. Physiol. Pharmacol. 76, 1080–1086. https://doi.org/10.1139/cjpp-76-12-1080.

Lautt, W.W., Wong, C., 1978. Hepatic parasympathetic neural effect on glucose balance in the intact liver. Can. J. Physiol. Pharmacol. 56, 679–682. https://doi.org/10.1139/y78-163.

Lautt, W.W., Xie, H., 1998. Intraportal acetylcholine reverses insulin resistance caused by chronic bile duct ligation. Proc. West. Pharmacol. Soc. 41: 35-36. PMID: 9836233.

Levin, B.E., 2006. Metabolic imprinting: critical impact of the perinatal environment on the regulation of energy homeostasis. Philos. Trans. R Soc. Lond. B Biol. Sci. 361(1471), 1107–1121. https://doi.org/10.1098/rstb.2006.1851.

Lieber, C.S., 1999. Role of S-adenosyl-L-methionine in the treatment of liver diseases. Hepatology 30(6), 1155–1159. https://doi.org/10.1016/s0168-8278(99)80274-8.

Lovat, N.E.J., Legare, D.J., Gieni, R.S., Lautt, W.W. 2020. Gestational postprandial insulin sensitivity in the Sprague Dawley rat: the putative role of hepatic insulin sensitizing substance in glucose partitioning in pregnancy. Can. J. Physiol. Pharmacol. 98(8), 541–547. https://doi.org/10.1139/cjpp-2019-0575.

Lovat, N.E.J., Legare, D.J., Lautt W.W., 2021. An animal model of gestational obesity and prediabetes: HISS-dependent insulin resistance induced by a high sucrose diet in Sprague Dawley rats. Can. J. Physiol. Pharmacol. 99, 599–608. https://doi.org/10.1139/cjpp-2020-0340.

Macedo, M.P., Lima, I.S., Gaspar, J.M., Afonso, R.A., Patarrao, R.S., Kim, Y.B., Ribeiro, R.T., 2013. Risk of postprandial insulin resistance: The liver/vagus rapport. Rev. Endocr. Metab. Disord. 15, 67–77. https://doi.org/10.1007/s11154-013-9281-5.

Martins, F.O., Delgado, T.C., Viegas, J., Gaspar J.M., Scott, D.K., O'Doherty, R.M., Macedo, M.P., Jones, J.G., 2016. Mechanisms by which the thiazolidinedione troglitazone protects against sucrose-induced hepatic fat accumulation and hyperinsulinemia. Brit. J. Pharmacy 173, 267–278. https://doi.org/10.1111/bph.13362.

Merker, K., Stolzing, A., Grune, T., 2001. Proteolysis, caloric restriction and aging. Mech. Ageing Dev. 122, 595–615. https://doi.org/10.1016/s0047-6374(01)00219-6.

Mero, N., Syvanne, M., Taskinen, M-R., 1998. Postprandial lipid metabolism in diabetes. Atherosclerosis 141, S53–55. https://doi.org/10.1016/s0021-9150(98)00218-4.

Ming, Z., Fan, Y-J., Yang, X., Lautt, W.W., 2006. Synergistic protection by S-adenosyl-methionine with vitamins C and E on liver injury induced by thioacetamide in rats. Free Rad. Biol. Med. 40, 617–624. https://doi.org/10.1016/j.freeradbiomed.2005.09.034.

Ming, Z., Legare, D.J., Lautt, W.W., 2009. Obesity, syndrome X, and diabetes: the role of HISSdependent insulin resistance altered by sucrose, an antioxidant cocktail, and age. Can. J. Physiol. Pharmacol. 87, 873–882. https://doi.org/10.1139/Y09-079. Ming, Z., Legare, D.J., Lautt, W.W., 2011. Absence of meal-induced insulin sensitization (AMIS) in aged rats is associated with cardiac dysfunction that is protected by antioxidants J. Appl. Physiol. 111, 704–714. https://doi.org/10.1152/japplphysiol.00057.2011.

Ming, Z., Lautt, W.W., 2011. HISS, not insulin, causes vasodilation in response to administered insulin. J. Appl. Physiol. 110, 60–68. https://doi.org/10.1152/japplphysiol.00714.2010.

Minuk, G.Y., Meyers, A.F.A., Legare, D.J., Sadri, P., Lautt, W.W., 1998. Fetal exposure to alcohol results in adult insulin resistance in the rat. Proc. West. Pharmacol. Soc. 41, 39–40. PMID: 9836235.

Mondon, C.E., Burton, S.D., 1971. Factors modifying carbohydrate metabolism and effect of insulin in perfused rat liver. Am. J. Physiol. 220(3), 724–734. https://doi.org/10.1152/ajplegacy.1971.220.3.724.

Nelson, S.M., Matthews, P., Poston, L., 2010. Maternal metabolism and obesity: modifiable determinants of pregnancy outcome. Hum. Reprod. Update 16(3), 255–275. https://doi.org/10.1093/humupd/dmp050.

O'Brien, I.A.D., O'Hare, P., Corrall, R.J.M., 1986. Heart rate variability in healthy subjects: Effect of age and the derivation of normal ranges for tests of autonomic function. Br. Heart J. 55, 348–354.

Olson, G., Blackwell, S.C., 2011. Optimization of gestational weight gain in the obese gravida: a review. Obstet. Gynecol. Clin. North Am. 38(2), 397–407. https://doi.org/10.1016/j.ogc.2011.03.003.

Olson, C.M., Strawderman, M.S., Hinton, P.S., Pearson, T.A., 2003. Gestational weight gain and postpartum behaviors associated with weight change from early pregnancy to 1 y postpartum. Int. J. Obes. Relat. Metab. Disord. 27(1), 117–127. https://doi.org/10.1038/sj.ijo.0802156.

Patarrao, R.S., Lautt, W.W., Afonso, R.A., Riberio, R.T., Fernandes, A.B., Boavida, J.M., Macedo, M.P., 2012. Postprandial but not fasting insulin resistance is an early identifier of dysmetabolism in overweight subjects. Can. J. Physiol. Pharmacol. 90, 923–931. https://doi.org/10.1139/y2012-086.

Patarrao, R.S., Lautt, W.W., Afonso, R.A., Ribeiro, R.T., Guarino, M.P., Fernandes, A.B., Boavida, J.M., Macedo, M.P., 2008. Meal-induced insulin sensitization (MIS) and its parasympathetic regulation in humans. Can. J. Physiol. Pharmacol. 86, 880–888. https://doi.org/10.1139/Y08-080.

Patarrao, R.S., Lautt, W.W., Guarino, M.P., Afonso, R.A., Ribeiro, R.T., Fernandes, A.B., Boavida, J.M., Macedo, M.P., 2007. A new technique to assess insulin sensitivity in humans: the rapid insulin sensitivity test (RIST). Proc. West. Pharmacol. Soc. 50, 105–109. PMID: 18605244.

Patarrao, R.S., Lautt, W.W., Macedo, M.P., 2015. Acute Glucagon Induces Postprandial Peripheral Insulin Resistance. PLOS ONE 10(5), e0127221. https://doi.org/101371/journal.pone.0127221.

Peitl, B., Dobronte, R., Nemeth, J., Pankucsi, C., Sari, R., Varga, A., Szilvassy, Z., 2009. Mealinduced enhancement in insulin sensitivity is not triggered by hyperinsulinemia in rats. Metabolism 58, 328–332. https://doi.org/10.1016/j.metabol.2008.10.004.

Peitl, B., Nemeth, J., Pankucsi, C., Szilvassy, Z., 2006. Insulin sensitization induced by oral cicletanine in conscious rabbits. Naunyn. Schmiedebergs Arch. Pharmacol. 373, 429–439. https://doi.org/10.1007/s00210-006-0096-8.

Petersen, K.F., Dufour, S., Savage, D.B., Bilz, S., Solomon, G., Yonemitsu, S., Cline, G.W., Befroy, D., Zemany, L., Kahn, B.B., Papademetris, S., Rothman, D.L., Shulman, G.I., 2007. The role of skeletal muscle insulin resistance in the pathogenesis of the metabolic syndrome. Proc. Natl. Acad. Sci. USA 104, 12587–12594. https://doi.org/10.1073/pnas.0705408104.

Phillips, R.J., Powley, T.L., 2007. Innervation of the gastrointestinal tract: Patterns of aging. Auton. Neurosci. 136, 1–19. https://doi.org/10.1016/j.autneu.2007.04.005.

Reid, M.A., Latour, M.G., Legare, D.J., Rong, N., Lautt, W.W. 2002. Comparison of the rapid insulin sensitivity test (RIST), the insulin tolerance test (ITT), and the hyperinsulinemic euglycemic clamp (HIEC) to measure insulin action in rats. Can. J. Physiol. Pharmacol. 80, 811–818. https://doi.org/10.1139/y02-102.

Reid, M.A., Lautt, W.W., 2004. Pattern of insulin delivery affects hepatic insulin sensitizing substance (HISS) action and insulin resistance. Can. J. Physiol. Pharmacol. 82, 1068–1074. https://doi.org/10.1139/y04-111.

Ribeiro, R.T., Afonso, R.A., Guarino, M.P., Macedo, M.P., 2008. Loss of postprandial insulin sensitization during aging. J. Gerontol. A Biol. Sci. Med. Sci. 63, 560–565. https://doi.org/10.1093/gerona/63.6.560.

Ribeiro, R.T., Afonso, R.A., Macedo, M.P., 2002. The action of hepatic insulin-sensitizing substance: gender comparison in Wistar rats. Proc. West. Pharm. Soc. 45, 55–56. PMID: 12434527.

Ribeiro, R.T., Lautt, W.W., Legare, D.J., Macedo, M.P., 2005. Insulin resistance induced by sucrose-feeding is due to an impairment of the hepatic parasympathetic nerves. Diabetologia 48, 976–983. https://doi.org/10.1007/s00125-005-1714-6.

Sadri, P., Legare, D.J., Takayama, S., Lautt, W.W., 2003. Fetal ethanol exposure causes hepatic insulin sensitizing substance-dependent insulin resistance. Can. J. Diabetes 27(3), 239–247.

Sadri, P., Legare, D.J., Takayama, S., Lautt, W.W., 2005. Increased incidence of HISS-dependent insulin resistance in female rats prenatally exposed to ethanol. Can. J. Physiol. Pharmacol. 83, 383–387. https://doi.org/10.1139/y05-023.

Sadri, P., Reid, M.A.G., Afonso, R.A., Schafer, J., Legare, D.J., Macedo, M.P., Lautt, W.W., 2006. Meal-induced insulin sensitization in conscious and anesthetized rat models comparing liquid mixed meal with glucose and sucrose. Br. J. Nutr. 95, 288–295. https://doi.org/10.1079/bjn20051644.

Sanger, F., 1959. Chemistry of insulin; determination of the structure of insulin opens the way to greater understanding of life processes. Science 129, 1340–1344. https://doi.org/10.1126/science.129.3359.1340.

Selye, H., 1950. Stress and the general adaptation syndrome. Br. Med. J. 1, 1383–1392. https://doi.org/10.1136/bmj.1.4667.1383.

Schafer, J., Legare, D.J., Lautt, W.W., 2010. Acetylcholinesterase antagonist potentiated insulin action in fed but not fasted state. J. Pharmacol. Exp. Ther. 333, 621–628. https://doi.org/10.1124/jpet.109.164509.

Schneider, T., Hein, P., Michel-Reher, M.B., Michel, M.C., 2005. Effects of ageing on muscarinic receptor subtypes and function in rat urinary bladder. Naunyn Schmiedebergs Arch. Pharmacol. 372, 71–78. https://doi.org/10.1007/s00210-005-1084-0.

Scognamiglio, R., Negut, C., Vigili De Kreutzenberg, S., Tiengo, A., Avogaro, A., 2005. Postprandial myocardial perfusion in healthy subjects and in type 2 diabetic patients. Circulation 112, 179–184. https://doi.org/10.1161/CIRCULATIONAHA.104.495127.

Seredycz, L., Lautt, W.W., 2006. Hemorrhage results in hepatic insulin sensitizing substancedependent insulin resistance mediated by somatostatin in rats. Neuroendocrinology 84, 94– 102. https://doi.org/10.1159/000097484.

Seredycz, L, Ming, Z., Lautt, W.W., 2006. Acute hemorrhage causes hepatic insulin sensitizing substance (HISS)-dependent insulin resistance. Can. J. Physiol. Pharmacol. 84, 1145–1151. https://doi.org/10.1139/y06-064.

Shaw, J.E., Hodge, A.M., de Courten, M., Chitson, P., Zimmet, P.Z., 1999. Isolated post-challenge hyperglycemia confirmed as a risk factor for mortality. Diabetologia 42, 1050–1054. https://doi.org/10.1007/s001250051269.

Shichiri, M., Kishikawa, H., Ohkubo, Y., Wake, N., 2000. Long-term results of the Kumamoto Study on optimal diabetes control in type 2 diabetic patients. Diabetes Care 23(Suppl 2), B21–29. PMID: 10860187.

Simon, C., Brandenberger, G., 2002. Ultradian oscillations of insulin secretion in humans. Diabetes 51, S258–261. https://doi.org/10.2337/diabetes.51.2007.s258.

Simon, C., Follenius, M., Brandenberger, G., 1987. Postprandial oscillations of plasma glucose insulin and C-peptide in man. Diabetologia 30, 769–773. https://doi.org/10.1007/BF00275742.

Skrapari, I., Tentolouris, N., Perrea, D., Bakoyiannis, C., Papazafiropoulou, A., Katsilambros, N., 2007. Baroreflex sensitivity in obesity: Relationship with cardiac autonomic nervous system activity. Obesity 15, 1685–1693. https://doi.org/10.1038/oby.2007.201.

Stern, N., Izkhakov, Y., 2006. The metabolic syndrome revisited: "cardiometabolic risk" emerges as common ground between differing views of the ADA and AHA. J. Cardiometab. Syndr. 1(5), 362–363. https://doi.org/10.1111/j.1524-6175.2006.05999.x.

Sun, S., Ji, Y., Kersten, S., Qi, L., 2012. Mechanisms of inflammatory responses in obese adipose tissue. Annu. Rev. Nutr. 32(1): 261-286. https://doi.org/10.1146/annurev-nutr-071811-150623.

Svensson, M.K., Lindmark, S., Wiklund, U., Rask, P., Karlsson, M., Myrin, J., Kullberg, J., Johansson, L., Eriksson, J.W., 2016. Alterations in heart rate variability during everyday life are linked to insulin resistance. A role of dominating sympathetic over parasympathetic nerve activity? Cardiovasc. Diabetol. 15, 91. https://doi.org/10.1186/s12933-016-0411-8.

Taylor, A.N., Branch, B.J., Liu, S.H., Kokka, N., 1982. Long-term effects of fetal ethanol exposure on pituitary-adrenal responses to stress. Pharmacol. Biochem. Behav. 16, 585–589. https://doi.org/10.1016/0091-3057(82)90420-8.

Ting, J.W, Lautt, W.W., 2006. The effect of acute, chronic, and prenatal ethanol exposure on insulin sensitivity. Pharmacol. Therap. 111, 346–373. https://doi.org/10.1016/j.pharmthera.2005.10.004.

Vaccaro, O., Ruffa, G., Imperatore, G., Iovino, V., Rivellese, A.A., Riccardi, G., 1999. Risk of diabetes in the new diagnostic category of impaired fasting glucose. Diabetes Care 22, 1490–1493. https://doi.org/10.2337/diacare.22.9.1490.

Van Oostrom, A.J.H.H.M., Sijmonsma, T.P., Rabelink, T.J., Van Asbeck, B.S., Castro Cabezas, M., 2003. Postprandial leukocyte increase in healthy subjects. Metabolism 52, 199–202. https://doi.org/10.1053/meta.2003.50037.

Von Kanel, R., Nelesen, R.A., Ziegler, M.G., Mausbach, B.T., Mills, P.J., Dimsdale, J.E., 2007. Relation of autonomic activity to plasminogen activator inhibitor-1 plasma concentration and the role of body mass index. Blood Coagul. Fibrinolysis 18, 353–359. https://doi.org/10.1097/MBC.0b013e3281139c67. Weinberg, J., 1992. Prenatal ethanol exposure alters adrenocortical response to predictable and unpredictable stressors. Alcohol 697, 86–96. https://doi.org/10.1016/0741-8329(92)90043-a.

Xie, H, Lautt, W.W., 1995a. Induction of insulin resistance by cholinergic blockade with atropine in the cat. J. Auton. Pharmacol. 15, 361–369. https://doi.org/10.1111/j.1474-8673.1995.tb00402.x.

Xie, H., Lautt, W.W. 1995b. M1 muscarinic receptor blockade causes insulin resistance in the cat. Proc. West. Pharmacol. Soc. 38, 83–84. PMID: 7480031.

Xie, H., Lautt, W.W., 1996a. Insulin resistance of skeletal muscle produced by hepatic parasympathetic interruption. Am. J. Physiol. 270: E858-E863. https://doi.org/10.1152/ajpendo.1996.270.5.E858.

Xie, H., Lautt, W.W., 1996b. Insulin resistance caused by hepatic cholinergic interruption and reversed by acetylcholine administration. Am. J. Physiol. 271: E587–E592. https://doi.org/10.1152/ajpendo.1996.271.3.E587.

Xie, H., Tsybenko, V.A., Johnson, M.V., Lautt, W.W., 1993. Insulin resistance of glucose response produced by hepatic denervations. Can. J. Physiol. Pharmacol. 71(2), 175–178. https://doi.org/10.1139/y93-024.

Xie, H., Zhu, L., Zhang, Y.L., Legare, D.J., Lautt, W.W., 1996. Insulin sensitivity tested with a modified euglycemic technique in cats and rats. J. Pharmacol. Toxicol. Meth. 35, 77–82. https://doi.org/10.1016/1056-8719(96)00003-2.

Zhu, M.J., Ma, Y., Long, N.M., Du, M., Ford, S.P., 2010. Maternal obesity markedly increases placental fatty acid transporter expression and fetal blood triglycerides at midgestation in the ewe. Am. J. Physiol. Regul. Integr. Comp. Physiol. 299(5), R1224–R1231. https://doi.org/10.1152/ajpregu.00309.2010.