Online Supplement to “Statistical Learning of Neuronal Functional Connectivity”

Appendix A: List of notations and symbols

List of notations and symbols.

- C : total number of neurons in the ensemble.
- T : total length of the experiment trial.
- n : total number of time bins. We assume $n \rightarrow \infty$.
- p_g : the length of the g th group for the parameters.
- $\delta = T/n$: the length of each time bin.
- $\tau_k = (t_{k-1}, t_k]$: the start and end time points of the k th time bin.
- $N_c(\tau_k)$: count the number of spikes fired by neuron c during τ_k .
- $\tau_{0:k} = (0, t_k]$: the interval from the start of the experiment up to the time point t_k .
- $N_{1:C}(\tau_{0:k}) = \{N_c(\tau_k) : c = 1, \dots, C; k = 1, \dots, n\}$ if $k \geq 1$ or empty if $k = 0$: the spiking history of all neurons up to the time point t_k .
- $\lambda_c(\tau_k | \cdot) = E\{N_c(\tau_k) | \cdot\}$: the conditional intensity function of neuron c during τ_k .
- $\exp(\gamma_{c;0})$: the baseline firing rate of neuron c .
- $\boldsymbol{\gamma}^{(c)} = (\gamma_{c;1}, \dots, \gamma_{c;P})^T$: the vector of autoregressive parameters of neuron c at different lags.
- $\boldsymbol{\gamma}^{(c,i)}$: the vector of interactive parameters representing the influence of neuron i onto neuron c at different lags. For parametrization (2.1), $\boldsymbol{\gamma}^{(c,i)} = (\gamma_{c,i;1}, \dots, \gamma_{c,i;Q})^T$ as in (2.4); for parametrization (4.1), $\boldsymbol{\gamma}^{(c,i)} = \gamma_{c,i;1}$ as in (4.2).
- $\mathbf{X}(\tau_k)$: a M dimensional covariate vector measured during τ_k .
- $\boldsymbol{\beta}_c = (\beta_{c,1}, \dots, \beta_{c,M})^T$: the parameters that correspond to the covariates for neuron c .
- P : the history windows of autoregressive parameters.
- Q : the history windows of interactive parameters.
- M : the length of covariate vector $\mathbf{X}(\tau_k)$.
- $\tilde{\boldsymbol{\theta}}_c$: the vector form of all parameters for neuron c , i.e. $\tilde{\boldsymbol{\theta}}_c = (\gamma_{c;0}, \boldsymbol{\theta}_c^T)^T$.
- $\boldsymbol{\theta}_c$: the vector form of all parameters for neuron c excluding the intercept $\gamma_{c;0}$.

- $\tilde{\boldsymbol{\theta}}_c^*$: the true unknown parameter vector.
- $d = P + (C - 1)Q + M$: the total number of covariates
- $\ell_c(\tilde{\boldsymbol{\theta}}_c)$: negative log-likelihood of neuron c .
- $\mathcal{P}_{(\eta_c, \alpha_c)}(\boldsymbol{\theta}_c)$: structured penalty of $\boldsymbol{\theta}_c$.
- $L_c(\tilde{\boldsymbol{\theta}}_c)$: negative log-likelihood of neuron c with the penalty.
- (α_c, η_c) : tuning parameters of the penalty.
- $\|\cdot\|_2$: the L_2 norm.
- $\|\cdot\|_1$: the L_1 norm.

Additional notations and symbols. Furthermore, we define by $Y_{c,k}$ and $\tilde{\mathbf{X}}_{c,k}$ the response scalar and covariate vector of (2.5) at the k th time bin, $k = 1, \dots, n$. Let $\tilde{\mathcal{X}}_c = (\tilde{\mathbf{X}}_{c,1}, \dots, \tilde{\mathbf{X}}_{c,n})^T$ denote the whole design matrix. Adopting the BD framework in Zhang *et al.* (2010), the problem of (2.5) is equivalent to the penalized BD estimation with link function $F(\cdot) = \log(\cdot)$ and

$$Q(Y, \mu) = \mu - Y \log(\mu) - Y + Y \log(Y)$$

generated by $q(\mu) = \mu - a - \mu\{\log(\mu) - \log(a)\}$ where a is a finite positive constant such that $q(\mu)$ is well-defined.

Appendix B: Proofs of Results in Section 3

Conditions. We have the following conditions in which B_1 and B_2 are sufficiently large constants. Those are not the weakest possible, but serve to facilitate the technical derivations.

- A1. The parameter space Θ of $\tilde{\boldsymbol{\theta}}_c$ is compact in $\mathbb{R}^{d+1}$ and $\|\tilde{\boldsymbol{\theta}}_c^*\|_1 < B_1$;
- A2. For all bins, $N_c(\tau_k)$ are uniformly bounded by B_2 ;
- A3. The minimum eigenvalue of $\Sigma_c = n^{-1} \tilde{\mathcal{X}}_c \tilde{\mathcal{X}}_c^T$ are uniformly bounded away from 0, i.e. $\lambda_{\min}(\Sigma_c) > \delta$ for some positive constant $\delta > 0$;

Condition A1 restrict the magnitude of parameters from being too large or too small respectively. Condition A2 reflects the biological limitation to the number of spikes within a

short time interval. The boundedness assumption of the response variable would also facilitate the technical derivations. Condition A3 is commonly used for the covariance matrix of predictor variables and it ensures the convexity of likelihood function and identifiability of the problem.

Proof of Theorem 1. Since $d = P + (C - 1)Q + M$ and P, Q, M are all fixed constants, d and C have the same diverging rate. Let $r_n = \sqrt{d/n}$ and $\tilde{\mathbf{u}} = (u_0, u_1, \dots, u_d)^T \in \mathbb{R}^{d+1}$. Similar to Zhang *et al.* (2010), it suffices to show that for any given $\epsilon > 0$, there is a large constant U_ϵ such that, for large n ,

$$\mathbb{P}\left\{\inf_{\|\tilde{\mathbf{u}}\|_2=U_\epsilon} L_c(\tilde{\boldsymbol{\theta}}_c^* + r_n \tilde{\mathbf{u}}) > L_c(\tilde{\boldsymbol{\theta}}_c^*)\right\} \geq 1 - \epsilon. \quad (\text{A.1})$$

Now we separate the loss function and penalty term by

$$\begin{aligned} L_c(\tilde{\boldsymbol{\theta}}_c^* + r_n \tilde{\mathbf{u}}) - L_c(\tilde{\boldsymbol{\theta}}_c^*) \\ &= \{\ell_c(\tilde{\boldsymbol{\theta}}_c^* + r_n \tilde{\mathbf{u}}) - \ell_c(\tilde{\boldsymbol{\theta}}_c^*)\} + \{\mathcal{P}_{(\eta_c, \alpha_c)}(\tilde{\boldsymbol{\theta}}_c^* + r_n \tilde{\mathbf{u}}) - \mathcal{P}_{(\eta_c, \alpha_c)}(\tilde{\boldsymbol{\theta}}_c^*)\} \\ &\equiv I_1 + I_2. \end{aligned}$$

Further by Taylor expansion of I_1 ,

$$\begin{aligned} I_1 &= \frac{r_n}{n} \sum_{k=1}^n \{\exp(\tilde{\mathbf{X}}_{c,k}^T \tilde{\boldsymbol{\theta}}_c^*) - Y_{c,k}\} \tilde{\mathbf{X}}_{c,k}^T \tilde{\mathbf{u}} + \frac{r_n^2}{2n} \sum_{k=1}^n \exp(\tilde{\mathbf{X}}_{c,k}^T \tilde{\boldsymbol{\theta}}_c^*) (\tilde{\mathbf{X}}_{c,k}^T \tilde{\mathbf{u}})^2 \\ &\quad + \frac{r_n^3}{6n} \sum_{k=1}^n \exp(\tilde{\mathbf{X}}_{c,k}^T \tilde{\boldsymbol{\theta}}_c') (\tilde{\mathbf{X}}_{c,k}^T \tilde{\mathbf{u}})^3 = I_{1,1} + I_{1,2} + I_{1,3} \end{aligned}$$

where $\tilde{\boldsymbol{\theta}}_c'$ is located between $\tilde{\boldsymbol{\theta}}_c^*$ and $\tilde{\boldsymbol{\theta}}_c^* + r_n \tilde{\mathbf{u}}$. Then those terms can be bounded as

$$|I_{1,1}| \leq \left\| \frac{r_n}{n} \sum_{k=1}^n \{\exp(\tilde{\mathbf{X}}_{c,k}^T \tilde{\boldsymbol{\theta}}_c^*) - Y_{c,k}\} \tilde{\mathbf{X}}_{c,k} \right\|_2 \|\tilde{\mathbf{u}}\|_2 = O_P(r_n \sqrt{d/n}) U_\epsilon; \quad (\text{A.2})$$

$$I_{1,2} = 2^{-1} r_n^2 \tilde{\mathbf{u}}^T (\mathcal{X}_c^T \Omega \mathcal{X}_c / n) \tilde{\mathbf{u}} \geq c_1 r_n^2 U_\epsilon^2; \quad (\text{A.3})$$

$$|I_{1,3}| \leq O_P(r_n^3 d^{3/2}) U_\epsilon^3, \quad (\text{A.4})$$

where $\Omega = \text{diag}\{\exp(\tilde{\mathbf{X}}_{c,1}^T \tilde{\boldsymbol{\theta}}_c^*), \dots, \exp(\tilde{\mathbf{X}}_{c,n}^T \tilde{\boldsymbol{\theta}}_c^*)\}$ and c_1 is some positive constant. Here the data are not i.i.d., so we need to apply the martingale version of central limit theorem (Theorem 7.4 in Durrett, 2004) to obtain the rate of $I_{1,1}$ in (A.2), in which each element of $\sum_{k=1}^n \{\exp(\tilde{\mathbf{X}}_{c,k}^T \tilde{\boldsymbol{\theta}}_c^*) - Y_{c,k}\} \tilde{\mathbf{X}}_{c,k}^T$ is a martingale with bounded increments. (A.3) and (A.4) follow conditions A1-A3.

For the penalty term,

$$\begin{aligned}
I_2 \geq & -(1 - \alpha_c)\eta_c \left\{ \sqrt{P} \|r_n \mathbf{u}_{\gamma^{(c)}}\|_2 + \sqrt{Q} \sum_{i \in \{1, \dots, C\} \setminus c} \|r_n \mathbf{u}_{\gamma^{(c,i)}}\|_2 \right\} \\
& - \alpha_c \eta_c \left\{ \|r_n \mathbf{u}_{\gamma^{(c)}}\|_1 + \sum_{i \in \{1, \dots, C\} \setminus c} \|r_n \mathbf{u}_{\gamma^{(c,i)}}\|_1 \right\} - \eta_c \|r_n \mathbf{u}_{\beta_c}\|_1
\end{aligned}$$

where $\mathbf{u}_{\gamma^{(c)}}$, $\mathbf{u}_{\gamma^{(c,i)}}$ and $\mathbf{u}_{\beta_c}$ are components of $\tilde{\mathbf{u}}$ corresponding to $\gamma^{(c)}$, $\gamma^{(c,i)}$ and $\mathbf{u}_{\beta_c}$ respectively. Then,

$$\begin{aligned}
|I_2| & \leq \eta_c r_n \left\{ \sqrt{P} \|\mathbf{u}_{\gamma^{(c)}}\|_2 + \sqrt{Q} \sum_{i \in \{1, \dots, C\} \setminus c} \|\mathbf{u}_{\gamma^{(c,i)}}\|_2 + \sqrt{M} \|\mathbf{u}_{\beta_c}\|_2 \right\} \\
& \leq \eta_c r_n \sqrt{\max(P, Q, M)} \sqrt{C+1} \|\tilde{\mathbf{u}}\|_2 = O(\eta_c r_n \sqrt{d}) U_\epsilon.
\end{aligned}$$

Since $d^4/n = O(C^4/n) = o(1)$, we can choose some large C_ϵ such that $I_{1,1}$, $I_{1,3}$ and I_2 are all dominated by $I_{1,2}$, which is positive. This implies (A.1). ■

Appendix C: Figures and Tables in the paper

Section 4.1. Figures 1–2; Tables 1–2.

Figure 1: **(Simulation of simple network)** A simulated network with 10 neurons. The line types of arrows indicate two different types of interactive parameters in Figure 2, where the thick Red line represents type-A parameter (more excitatory) and the thin Blue line represents type-B parameter (more inhibitory).

Figure 2: **(Simulation of simple network)** True values (using $\circ$ connected with dashed lines) of parameters for the simulation. The horizontal dotted line indicates 0. Left panel: autoregressive parameters $\{\gamma_{c;p}\}_{p=1}^P$; middle panel: interactive type-A parameters $\{\gamma_{c,i;q}\}_{q=1}^Q$; right panel: interactive type-B parameters $\{\gamma_{c,i;q}\}_{q=1}^Q$.

Table 1: **(Simulation of simple network)** Comparison of estimated network by regularized methods with varying strength of parameters and length of spike train. Results are averaged over 100 replications, with standard deviations indicated in parentheses.

Length (bins)	Relative Strength	Method	Correct-All	detected-A	detected-B	Correct-NC
10000	100%	L_1 -short-10-3	1.58 (1.4)	1.34 (1.1)	0.24 (0.5)	79.11 (1.2)
		L_1 -10-10	0.61 (0.8)	0.59 (0.8)	0.02 (0.1)	79.72 (0.7)
		TR-5-5	3.01 (1.4)	2.47 (1.2)	0.54 (0.6)	76.72 (2.1)
		TR-10-10	1.73 (1.1)	1.54 (1.1)	0.19 (0.4)	77.90 (1.5)
		SGL-5-5	4.98 (1.6)	4.21 (1.4)	0.77 (0.7)	75.64 (3.1)
		SGL-10-10	4.73 (1.9)	3.57 (1.5)	1.16 (0.8)	67.02 (8.2)
		SGL-15-15	4.62 (1.8)	3.49 (1.4)	1.13 (0.8)	58.95 (9.7)
		True network	10.00 (0.0)	7.00 (0.0)	3.00 (0.0)	80.00 (0.0)
	125%	L_1 -short-10-3	4.02 (1.6)	3.40 (1.3)	0.62 (0.7)	78.23 (1.7)
		L_1 -10-10	1.70 (1.4)	1.66 (1.3)	0.04 (0.2)	79.68 (0.5)
		TR-5-5	4.46 (1.4)	3.79 (1.2)	0.67 (0.8)	77.30 (1.8)
		TR-10-10	2.78 (1.5)	2.58 (1.3)	0.20 (0.4)	78.00 (1.7)
		SGL-5-5	7.54 (1.4)	6.13 (0.9)	1.41 (1.0)	73.43 (4.0)
		SGL-10-10	7.54 (1.5)	5.69 (1.1)	1.85 (0.7)	62.93 (8.2)
		SGL-15-15	7.23 (1.6)	5.39 (1.1)	1.84 (0.8)	50.74 (11.1)
		True network	10.00 (0.0)	7.00 (0.0)	3.00 (0.0)	80.00 (0.0)

Table 2: *The caption is identical to that for Table 1, except that 15000 bins are used.*

Length (bins)	Relative Strength	Method	Correct-All	detected-A	detected-B	Correct-NC
15000	100%	L_1 -short-10-3	6.27 (1.7)	4.86 (1.3)	1.41 (0.8)	74.42 (3.3)
		L_1 -10-10	2.70 (1.7)	2.42 (1.4)	0.28 (0.5)	78.84 (1.2)
		TR-5-5	3.72 (1.6)	3.22 (1.4)	0.50 (0.6)	77.31 (2.2)
		TR-10-10	2.37 (1.2)	2.08 (1.1)	0.29 (0.5)	78.59 (1.2)
		SGL-5-5	7.33 (1.5)	5.70 (1.0)	1.63 (0.9)	74.62 (3.2)
		SGL-10-10	7.72 (1.6)	5.73 (1.1)	1.99 (0.9)	64.49 (7.4)
		SGL-15-15	7.42 (1.6)	5.46 (1.1)	1.96 (0.8)	51.80 (9.4)
		True network	10.00 (0.0)	7.00 (0.0)	3.00 (0.0)	80.00 (0.0)
	125%	L_1 -short-10-3	8.26 (1.4)	6.25 (0.8)	2.01 (0.9)	73.61 (2.9)
		L_1 -10-10	4.89 (1.7)	4.37 (1.4)	0.52 (0.6)	78.27 (1.5)
		TR-5-5	6.05 (1.7)	4.93 (1.4)	1.12 (0.8)	77.22 (2.0)
		TR-10-10	4.10 (1.4)	3.70 (1.3)	0.40 (0.6)	78.37 (1.6)
		SGL-5-5	9.13 (0.9)	6.82 (0.4)	2.31 (0.8)	72.26 (4.0)
		SGL-10-10	9.39 (0.9)	6.79 (0.4)	2.60 (0.6)	61.96 (8.5)
		SGL-15-15	9.24 (0.8)	6.65 (0.6)	2.59 (0.5)	47.15 (10.2)
		True network	10.00 (0.0)	7.00 (0.0)	3.00 (0.0)	80.00 (0.0)

Section 4.2. Figures 3, 4, 5 and 6; Table 3.

Figure 3: **(Simulation of complex network)** A simulated network with 60 neurons. The line types of arrows indicate two different types of interactive parameters in Figure 2, where the thick Red line represents type-A parameter and the thin Blue line represents type-B parameter.

Figure 4: **(Simulation of complex network)** The connectivity matrix of the simulated network. The color of the cell at the i th row and the j th column distinguishes how neuron j 's spiking history would affect the firing rate of neuron i . The background Green color means no connection; the Red color represents type-A parameter and the Blue color represents type-B parameter.

Figure 5: **(Simulation of complex network)** Boxplots comparing 5 methods. Left panel: number of “detected-A” connections; middle panel: number of “detected-B” connections; right panel: number of false detections. Here $n = 25000$ and the relative strength is 125%.

Figure 6: **(Simulation of complex network)** The estimated parameters of all type-A and type-B connections in a single simulation run. Left panels: plots of type-A parameters; right panels: plots of type-B parameters. Top panels: plots of the estimates obtained by L1-10-10; bottom panels: plots of the estimates obtained by SGL-10-10. Thick $\circ$ connected with dashed lines: true values; thin dashed lines: estimated values. Here $n = 25000$ and the relative strength is 125%.

Table 3: **(Simulation of complex network)** Comparison of estimated network by regularized methods with varying strength of parameters and length of spike train. Results are averaged over 100 replications, with standard deviations indicated in parentheses.

Length (bins)	Relative Strength	Method	Correct-All	detected-A	detected-B	Correct-NC
20000	100%	L_1 -short-10-3	23.65 (3.4)	17.19 (2.7)	6.46 (2.0)	3457.07 (7.7)
		L_1 -10-10	15.94 (3.2)	13.04 (2.7)	2.90 (1.6)	3464.85 (5.0)
		SGL-5-5	30.42 (3.2)	21.84 (2.3)	8.58 (2.5)	3460.30 (4.7)
		SGL-10-10	31.78 (3.9)	19.57 (2.5)	12.21 (3.0)	3466.68 (3.5)
		SGL-15-15	21.97 (3.7)	14.78 (2.6)	7.19 (2.5)	3473.01 (2.7)
		True network	60.00 (0.0)	30.00 (0.0)	30.00 (0.0)	3480.00 (0.0)
	125%	L_1 -short-10-3	35.23 (3.5)	24.77 (2.0)	10.46 (2.5)	3450.98 (7.8)
		L_1 -10-10	26.86 (3.5)	21.60 (2.6)	5.26 (2.0)	3463.54 (4.4)
		SGL-5-5	46.76 (3.0)	28.83 (1.1)	17.93 (2.7)	3454.38 (5.3)
		SGL-10-10	50.63 (3.0)	28.15 (1.3)	22.48 (2.6)	3459.48 (4.7)
		SGL-15-15	42.37 (3.4)	25.73 (1.6)	16.64 (2.8)	3466.64 (3.7)
		True network	60.00 (0.0)	30.00 (0.0)	30.00 (0.0)	3480.00 (0.0)
25000	100%	L_1 -short-10-3	41.70 (3.2)	25.92 (1.8)	15.78 (2.6)	3426.09 (9.9)
		L_1 -10-10	27.75 (3.6)	19.77 (2.5)	7.98 (2.3)	3454.48 (6.4)
		SGL-5-5	39.92 (3.0)	25.83 (1.9)	14.09 (2.6)	3456.61 (5.0)
		SGL-10-10	42.18 (3.8)	24.27 (2.3)	17.91 (2.9)	3463.31 (3.8)
		SGL-15-15	33.19 (3.7)	20.60 (2.8)	12.59 (2.5)	3469.86 (3.3)
		True network	60.00 (0.0)	30.00 (0.0)	30.00 (0.0)	3480.00 (0.0)
	125%	L_1 -short-10-3	50.69 (2.5)	29.35 (0.8)	21.34 (2.4)	3418.31 (9.5)
		L_1 -10-10	40.47 (3.3)	27.41 (1.6)	13.06 (2.7)	3453.88 (6.1)
		SGL-5-5	53.59 (2.4)	29.79 (0.5)	23.80 (2.3)	3450.05 (6.0)
		SGL-10-10	56.97 (1.8)	29.69 (0.6)	27.28 (1.5)	3456.29 (4.9)
		SGL-15-15	52.68 (2.4)	28.90 (0.9)	23.78 (2.3)	3463.03 (3.5)
		True network	60.00 (0.0)	30.00 (0.0)	30.00 (0.0)	3480.00 (0.0)

Section 4.3. Table 4.

Table 4: **(Simulation of real network)** Comparison of estimated network by regularized methods for the simulated network that mimics the real data, using $P = 30$ and $Q = 10$ in (2.1). Results are averaged over 100 replications, with standard deviations indicated in parentheses.

Method	Exper. 1 (with 73 neurons)		Exper. 4 (with 77 neurons)	
	Correct-All	Correct-NC	Correct-All	Correct-NC
L_1 -short-30-3	47.62 (1.5)	5048.70 (7.8)	53.95 (1.4)	5681.65 (7.9)
L_1 -30-10	45.47 (1.4)	5045.85 (7.7)	50.55 (1.4)	5689.68 (6.3)
SGL-30-10	48.44 (1.3)	5164.92 (5.5)	51.21 (1.0)	5745.54 (4.7)
True network	62.00 (0.0)	5194.00 (0.0)	71.00 (0.0)	5781.00 (0.0)

Section 5. Figures 7, 8, 9 and 10; Tables 5, 6, 7 and 8.

Figure 7: **(Neurophysiological data)** T-maze task: for each trial, the animal was placed in a start-box for a pre-determined time (a delay) and released with the removal of a starting gate. The rat would move to the junction of the “T” and choose the left or right arm. On a correct choice, the animal was rewarded and returned in the start-box. This figure is adapted from Devilbiss et al. (2012).

Figure 8: **(Neurophysiological data)** Estimates (using $\circ$ connected with dashed lines) of gain, i.e. $\exp(\gamma_{c;p})$, associated with autoregressive parameters by the SGL regularized method for selected neurons in Experiment 4.

Figure 9: **(Neurophysiological data)** The estimated network structures by the SGL regularized method. The thick Red color presents the excitatory connection and thin Blue color presents the inhibitory connection. Left panel: Experiment 1; right panel: Experiment 4.

Table 5: **(Neurophysiological data)** Summary statistics of the T-maze task dataset.

No.	Subject	Total length (sec)	Total number of neurons			Average firing rate (Hz)
			WS-type	NS-type	Total	
1	TM19	1449.2	53	4	73	0.567
2	TM19	1150.8	60	5	83	0.408
3	TM44	2243.7	44	6	59	0.772
4	TM44	2150.3	62	3	77	0.364
5	TM44	2191.1	57	3	71	0.192
6	TM44	2011.4	53	3	63	0.241
7	TM57	930.3	30	8	39	0.375
8	TM58	765.2	23	0	26	0.342

Figure 10: (Neurophysiological data) The estimated network matrices by the SGL regularized method. The Red color indicates the excitatory connection between the pair and the Blue color indicates the inhibitory connection, while the Green color means no functional connection. The exact color is determined by the strength of the connection. There is no estimate of interactive parameters for the blocks on the black diagonal line.

Table 6: **(Neurophysiological data)** *Description of the covariates used in the analysis.*

Column #	Name	Description
1	Delay	Stay in the resting area until the gate is released.
2	Run	Run through the tunnel from the gate to the branch point.
3	Decision	At the branch area of the T-maze and make a turn.
4	Choice	Run to the left or right end.
5	Reward	Eat reward in the correct trial.
6	Pick-up	Be picked up and put back to the resting area.
7	Left-trial	Whether made the left turn in the trial.
8	Correct-trial	Whether made the correct choice in the trial. (different from the previous one.)

Table 7: **(Neurophysiological data)** *Summary of estimated networks by regularized methods.*

Experiment	Possible pairs	L_1 -short-30-3		L_1 -30-10		SGL-30-10	
		Excit.	Inhib.	Excit.	Inhib.	Excit.	Inhib.
1	5256	48	40	17	14	43	19
2	6806	20	24	14	9	15	18
3	3422	116	53	49	13	63	16
4	5852	99	54	21	12	52	19
5	4970	21	9	3	0	9	4
6	3906	28	7	16	2	14	4
7	1482	55	6	13	2	23	0
8	650	0	0	0	0	0	0

Table 8: **(Neurophysiological data)** *Summary of estimated parameters by the SGL regularized method for 8 state indicator covariates described in Table 6.*

Experiment	Number of neurons with positive and negative estimated parameters															
	Delay		Run		Decision		Choice		Reward		Pick-up		Left		Correct	
	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-
1	6	5	2	0	11	3	0	0	0	0	3	4	4	3	2	6
2	9	7	6	1	1	1	2	0	2	1	3	2	0	2	0	1
3	8	29	6	1	6	3	2	0	0	1	7	0	2	8	1	3
4	8	26	7	0	6	0	7	0	0	0	5	7	8	3	6	3
5	5	13	2	1	1	0	2	0	0	2	6	1	1	3	1	3
6	11	14	2	0	2	0	2	0	4	1	3	4	1	4	0	1
7	10	7	16	1	6	0	1	1	1	6	1	10	4	6	0	6
8	0	0	0	0	0	0	0	0	1	0	4	0	1	0	1	0