

Supplementary Material

Planktonic foraminiferal assemblages reflect warming during two recent mid-latitude marine heatwaves

M. Kelsey Lane,* Jennifer S. Fehrenbacher, Jennifer L. Fisher, Melanie R. Fewings, Byron C. Crump, Craig M. Risien, Grace M. L. Meyer, Faith Schell

* **Correspondence:** *M. Kelsey Lane, lanemary@oregonstate.edu


1 Supplementary Data

Dataset S1 (separate file). Planktonic foraminifera count data. Station ID is in the format #####NH###, which indicates Month (##), Day (##), Year (##), NH Line Station (NH###).

Dataset S2 (separate file). Environmental data. Station ID is in the format #####NH###, which indicates Month (##), Day (##), Year (##), NH

2 Supplementary Figures and Tables

2.1 Supplementary Figure



Supplementary Figure 1 (Fig. S1). Temperature (°C) vertical profiles in the upper 100 m of the water column for each station sampled each year.

2.2 Supplementary Table

Table S1. **Environmental covariation.**

Covariation between environmental variables in a Pearson correlation matrix.

Parameters	SST	SSS	Ave. Temp	Ave. Sal	Log (Ext. Chl-a)	Dist. Offshore	BEUTI
SST							
SSS	-0.01						
Ave. Temp	0.66*	0.14					
Ave. Sal	-0.76*	0.25	-0.78*				
Log (Ext. Chl-a)	-0.58	-0.35	-0.46	0.45			
Dist. Offshore	0.47	0.06	0.68*	-0.57	-0.45		
BEUTI	-0.50	-0.57	-0.39	0.17	0.45	-0.08	

*Correlation >0.6