

## Supporting Information

# Chlorinated Iridoid Glucosides from *Veronica longifolia* and Their Antioxidant Activity

Søren R. Jensen,<sup>1,2</sup> Charlotte H. Gotfredsen,<sup>†</sup> U. Sebnem Harput,<sup>3</sup> and Iclal Saracoglu<sup>‡</sup>

*Department of Chemistry, The Technical University of Denmark, DK-2800, Lyngby,  
Denmark, and*

*Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, TR-06100,  
Ankara, Turkey*

---

<sup>1</sup> Technical University of Denmark.

<sup>2</sup> To whom correspondence should be addressed. Tel.: +45 45252103. Fax: +45 45933968.

E-mail: [srj@kemi.dtu.dk](mailto:srj@kemi.dtu.dk).

<sup>3</sup> Hacettepe University.

## List of Supporting Information

		<b>Page</b>
Figure S1	<sup>1</sup> H NMR (CD <sub>3</sub> OD, 500 MHz) of longifolioside A	3
Figure S2	<sup>13</sup> C NMR (CD <sub>3</sub> OD, 75 MHz) of longifolioside A	4
Figure S3	<sup>1</sup> H- <sup>1</sup> H NOESY NMR (CD <sub>3</sub> OD, 500 MHz) of longifolioside A	5
Figure S4	<sup>1</sup> H NMR (CD <sub>3</sub> OD, 500 MHz) of longifolioside B	6
Figure S5	<sup>13</sup> C NMR (CD <sub>3</sub> OD, 75 MHz) of longifolioside B	7




Figure S1




Figure S2




Figure S3




Figure S4




Figure S5