

Supporting Information



Sequential Synthesis of (*E*)- α,β -Unsaturated Primary Amides with Complete Stereoselectivity

José M. Concellón,* Humberto Rodríguez-Solla,* Carmen Concellón, Carmen Simal, and Noemí Alvaredo




Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, Julián Clavería, 8, 33071 Oviedo, Spain.
jmcg@uniovi.es

2-Chloro-3-hydroxydecanamide (3a).....	S2
2-Chloro-3-cyclohexyl-3-hydroxypropanamide (3b).....	S3
2-Chloro-3-hydroxy-5-phenylpentanamide (3c).....	S4
(<i>E</i>)-2-Decenamide (4a).....	S5
(<i>E</i>)-4-Methyl-2-hexenamide (4b).....	S6
(<i>E</i>)-3-Cyclohexylacrilamide (4c).....	S7
(<i>E</i>)-5-Phenyl-2-pentenamide (4d).....	S8
(<i>E</i>)-5,7,7-Trimethyl-2-octenamide (4e).....	S9
(2 <i>E</i> ,9 <i>Z</i>)-Dodeca-2,9-dienamide (4f).....	S10




2-Chloro-3-hydroxydecanamide (3a)





2-Chloro-3-cyclohexyl-3-hydroxypropanamide (3b)




2-Chloro-3-hydroxy-5-phenylpentanamide (3c)





(E)-2-Decenamide (4a)





(E)-4-Methyl-2-hexenamide (4b)





(E)-3-Cyclohexylacrylamide (4c)



(E)-5-Phenyl-2-pentenamide (4d)



(E)-5,7,7-Trimethyl-2-octenamide (4e)



(2E,9Z)-Dodeca-2,9-dienamide (4f)

