

Elucidating the encapsulation of siRNA in PEGylated cationic liposomes

Kevin Buyens, Joseph Demeester, Stefaan S. De Smedt* and Niek N. Sanders

[*] Corresponding Author: Prof. Stefaan C. De Smedt, Ghent Research Group on Nanomedicines, Laboratory of General Biochemistry and Physical Pharmacy, Faculty of Pharmaceutical Sciences, Ghent University, Harelbekestraat 72, 9000 Gent (Belgium)

E-mail: Stefaan.Desmedt@Ugent.be

Niek N. Sanders, Laboratory of Gene Therapy, Department of Nutrition, Genetics and Ethology, Faculty of Veterinary Medicine, Ghent University, Heidestraat 19, 9820 Merelbeke, Belgium

Supporting Information

To demonstrate that the total particle number in the size distribution plot (Figure 2 in the manuscript) is indeed 100%, a cumulative size distribution plot is depicted in Figure S1. It might seem that the total particle number in the size distribution plot is lower than 100%. However, the reason why it might seem less is that the size intervals used to plot this graph are quite narrow. Therefore, the number of particles within each interval is relatively low.




Figure S1. Cumulative size distribution of *CLASSIC* (black) and *HYDRA* (gray) siPLexes in absence (solid lines) and presence (dashed lines) of serum (FBS). SiPLexes were incubated for 1 hour at 37°C.