Supporting Information

Selective Antiproliferative Activity of Hydroxynaphthyl-β-D-xylosides

Mårten Jacobsson,[a] Ulf Ellervik,[a*] Mattias Belting,[b] Katrin Mani[c*]

[a)] *Organic Chemistry, Lund University, P.O. Box 124, SE-221 00 Lund, Sweden.*
[b)] *Department of Clinical Sciences, Section of Oncology, Lund University, Barngatan 2:1, SE-221 85, Lund, Sweden and* [c)] *Department of Experimental Medical Science, Division of Neuroscience, Lund University, Biomedical Center C13, SE-221 84, Lund, Sweden.*

[*] Department of Experimental Medical Science, Division of Neuroscience, Lund University, Biomedical Center C13, SE-221 84, Lund, Sweden, Tel. +46-46-2224047, Fax. +46-46-2223128, e-mail. katrin.mani@medkem.lu.se

Table of Contents

Results from crystal violet method S2-S7

^{1}H-NMR- and UV-spectrum for 4-(1,2-dihydroxynaphthyl)-β-D-xylopyranoside S8-S9



Figure 1: Inhibitory effect of compounds **1a**-**10a** (aglycons) on HFL-1 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 2: Inhibitory effect of compounds **1b**-**10b,c** (xylosides) on HFL-1 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 3: Inhibitory effect of compounds **1a**-**10a** (aglycons) on T24 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 4: Inhibitory effect of compounds **1b**-**10b,c** (xylosides) on T24 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 5: Inhibitory effect of compounds **1b**-**10b,c** (xylosides) on SV40 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 6: Inhibitory effect of compounds **1b**-**10b,c** (xylosides) on SV40 cells at varying concentrations as scored by the crystal violet method after 96 h exposure.



Figure 7: ^{1}H-NMR spectra for 4-(1,2-dihydroxynaphthyl)-β-D-xylopyranoside.



Figure 8: UV absorbance spectrum (200-500 nm) for 4-(1,2-dihydroxynaphthyl)-β-D-xylopyranoside.