

---

**Supporting Information for ic035132i to Inorg. Chem.**

**Nickel Complexes with New Bidentate P,N Phosphinito-Oxazoline and –  
Pyridine Ligands; Application for the Catalytic Oligomerization of  
Ethylene**

**Fredy Speiser, and Pierre Braunstein<sup>\*,†</sup>**

*Laboratoire de Chimie de Coordination (UMR 7513 CNRS), Université Louis Pasteur,  
4 rue Blaise Pascal, F-67070 Strasbourg Cédex, France*

**Lucien Saussine**


*Institut Français du Pétrole, Direction Catalyse et Séparation, CEDI René Navarre, BP 3,  
F-69390 Vernaison, France*

**Richard Welter**


*Laboratoire DECMET, UMR CNRS 7513, Université Louis Pasteur,  
4 rue Blaise Pascal, F-67070 Strasbourg Cedex, France.*

**FIGURES.** ORTEP Views of the molecular structures of complexes **14-17**. Thermal ellipsoids enclose 50% of the electron density


<sup>†</sup> For correspondence: E-mail: [braunst@chimie.u-strasbg.fr](mailto:braunst@chimie.u-strasbg.fr). Fax: +33-390 241 322.




**Figure 1.** ORTEP view of the molecular structure of **14** with the complete numbering scheme. Thermal ellipsoids enclose 50% of the electron density.



**Figure 2.** ORTEP view of the molecular structure of **15** with the complete numbering scheme. Thermal ellipsoids enclose 50% of the electron density.



**Figure 3.** ORTEP view of the molecular structure of **16** with the complete numbering scheme. Thermal ellipsoids enclose 50% of the electron density.



**Figure 4.** ORTEP view of the molecular structure of **17** with the complete numbering scheme. Thermal ellipsoids enclose 50% of the electron density.