

General experimental

Solvents and reagents. Unless otherwise stated, all reactions were carried out under an atmosphere of nitrogen in heat- or oven-dried glassware and using anhydrous solvent. Solvents and reagents were purchased from commercial suppliers and used as received. Reactions were monitored by thin layer chromatography (TLC) or liquid chromatography-mass spectroscopy (LCMS). Heating was conducted using hotplates with DrySyn adaptors.

Chromatography. Thin layer chromatography (TLC) was carried out using plastic-backed 50 precoated silica plates (particle size 0.2 mm). Spots were visualised by ultraviolet (UV) light (λ_{max} = 254 nm or 365 nm) and then stained with potassium permanganate solution followed by gentle heating. Flash column chromatography was carried out using the Teledyne ISCO CombiFlash® Rf+ apparatus with RediSep® or GraceResolv™ silica cartridges.

Liquid chromatography mass spectrometry. LCMS analysis carried out using system 1 described below unless otherwise stated as using system 2.

System 1

LCMS analysis was carried out on an H₂O_s Acquity UPLC instrument equipped with a BEH column (50 mm x 2.1 mm, 1.7 μ m packing diameter) and H₂O_s micromass ZQ MS using alternate-scan positive and negative electrospray. Analytes were detected as a summed UV wavelength of 210 – 350 nm. Three liquid phase methods were used:

Method A - Formic – 40 °C, 1 mL/min flow rate. Gradient elution with the mobile phases as (A) H₂O containing 0.1% volume/volume (v/v) formic acid and (B) acetonitrile containing 0.1% (v/v) formic acid. Gradient conditions were initially 1% B, increasing linearly to 97% B over 1.5 min, remaining at 97% B for 0.4 min then increasing to 100% B over 0.1 min.

Method B - High pH – 40 °C, 1 mL/min flow rate. Gradient elution with the mobile phases as (A) 10 mM aqueous ammonium bicarbonate solution, adjusted to pH 10 with 0.88 M aqueous ammonia and (B) acetonitrile. Gradient conditions were initially 1% B, increasing linearly to 97% B over 1.5 min, remaining at 97% B for 0.4 min then increasing to 100% B over 0.1 min.

Method C - TFA – 40 °C, 1 mL/min flow rate. Gradient elution with the mobile phases as (A) H₂O containing 0.1% volume/volume (v/v) TFA and (B) acetonitrile containing 0.1% (v/v) TFA. Gradient conditions were initially 1% B, increasing linearly to 97% B over 1.5 min, remaining at 97% B for 0.4 min then increasing to 100% B over 0.1 min.

System 2

LCMS analysis was carried out on an Agilent 1290 Infinity equipped with an acquity BEH column (50 mm x 2.1 mm, 1.7 µm packing diameter). Analytes were detected as a summed UV wavelength of 190 – 400 nm. Liquid phase method used:

Formic – 35 °C, 0.6 mL/min flow rate. Gradient elution with the mobile phases as (A) H₂O containing 0.1% volume/volume (v/v) formic acid and (B) acetonitrile containing 0.1% (v/v) formic acid. Gradient conditions were initially 3% B for 0.4 min, increasing linearly to 98% B over 2.8 min, remaining at 98% B for 0.6 min then decreasing to 3% B over 0.4 min then remaining at 3% B for 0.3 min. .

Nuclear magnetic resonance (NMR) spectroscopy. Proton (¹H), carbon (¹³C), and fluorine (¹⁹F) spectra were recorded in deuterated solvents at ambient temperature (unless otherwise stated) using standard pulse methods on any of the following spectrometers and signal frequencies: Bruker AV-400 (¹H = 400 MHz, ¹³C = 101 MHz, ¹⁹F = 376 MHz) and Bruker AV-600 (¹H = 600 MHz, ¹³C = 150 MHz). Chemical shifts are reported in ppm and are referenced to tetramethylsilane (TMS) or the following solvent peaks: CDCl₃ (¹H = 7.27 ppm, ¹³C = 77.0 ppm), DMSO-*d*₆ (¹H = 2.50 ppm, ¹³C = 39.5 ppm), and MeOH-*d*₄ (¹H = 3.31 ppm, ¹³C = 49.15 ppm). Peak assignments were made on the basis of chemical shifts, integrations, and coupling constants, using COSY, DEPT, and HSQC where appropriate. Coupling constants are quoted to the nearest 0.1 Hz and multiplicities are described as singlet (s), doublet (d), triplet (t), quartet (q), quintet (quin), sextet (sxt), septet (sept), br. (broad) and multiplet (m).

Infrared (IR) spectroscopy. Infrared spectra were recorded using a Perkin Elmer Spectrum 1 machine. Absorption maxima (ν_{\max}) are reported in wavenumbers (cm⁻¹) and are described as strong (s), medium (m), weak (w) and broad (br).

High-resolution mass spectrometry (HRMS). High-resolution mass spectra were recorded on a Micromass Q-ToF Ultima hybrid quadrupole time-of-flight mass spectrometer, with

analytes separated on an Agilent 1100 Liquid Chromatography equipped with a Phenomenex Luna C18 (2) reversed phase column (100 mm x 2.1 mm, 3 μ m packing diameter). LC conditions were 0.5 mL/min flow rate, 35 °C, injection volume 2–5 μ L. Gradient elution with (A) H₂O containing 0.1% (v/v) formic acid and (B) acetonitrile containing 0.1% (v/v) formic acid. Gradient conditions were initially 5% B, increasing linearly to 100% B over 6 min, remaining at 100% B for 2.5 min then decreasing linearly to 5% B over 1 min followed by an equilibration period of 2.5 min prior to the next injection. Mass to charge ratios (m/z) are reported in Daltons.

Melting points. Melting points were recorded on a Stuart SMP10 melting point apparatus.

Mass directed auto preparation (MDAP). Mass-directed automated purification was carried out using an H₂O_s ZQ MS using alternate-scan positive and negative electrospray and a summed UV wavelength of 210–350 nm. Two liquid phase methods were used:

Formic – Sunfire C18 column (100 mm x 19 mm, 5 μ m packing diameter, 20 mL/min flow rate) or Sunfire C18 column (150 mm x 30 mm, 5 μ m packing diameter, 40 mL/min flow rate). Gradient elution at ambient temperature with the mobile phases as (A) H₂O containing 0.1% volume/volume (v/v) formic acid and (B) acetonitrile containing 0.1% (v/v) formic acid.

High pH – Xbridge C18 column (100 mm x 19 mm, 5 μ m packing diameter, 20 mL/min flow rate) or Xbridge C18 column (150 mm x 30 mm, 5 μ m packing diameter, 40 mL/min flow rate). Gradient elution at ambient temperature with the mobile phases as (A) 10 mM aqueous ammonium bicarbonate solution, adjusted to pH 10 with 0.88 M aqueous ammonia and (B) acetonitrile.

TFA – Sunfire C18 column (100 mm x 19 mm, 5 μ m packing diameter, 20 mL/min flow rate) or Sunfire C18 column (150 mm x 30 mm, 5 μ m packing diameter, 40 mL/min flow rate). Gradient elution at ambient temperature with the mobile phases as (A) H₂O containing 0.1% volume/volume (v/v) TFA and (B) acetonitrile containing 0.1% (v/v) TFA.

The elution gradients used were at a flow rate of 40 mL/min over 10 or 20 min:

Gradient A	5-30% B
Gradient B	15-55% B


Gradient C	30-85% B
Gradient D	50-99% B
Gradient E	80-99% B

X-ray Crystallography.

- PDB file for compound **5** bound to LpPLA₂ available for immediate release upon publication.

Experimental

3-Hydroxy-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid – 7



To 1-bromo-4-(trifluoromethyl)benzene (197 g, 876 mmol) in THF (1000 mL) was added n-butyllithium (2.5M in hexanes) (500 mL, 1250 mmol) at -78°C . The reaction was stirred for 2 h before 3-oxocyclobutanecarboxylic acid (50 g, 438 mmol) in THF (250 mL) was added dropwise at -78°C . The reaction was allowed to warm to ambient temperature and was stirred for 16 h. The reaction mixture was quenched with saturated NH_4Cl solution (200 mL) and basified with 2M NaOH solution (500 mL) and washed with ethyl acetate (700 mL). The layers were separated and the aqueous layer was acidified with 2M HCl solution (1000 mL). The aqueous was extracted with ethyl acetate (1000 mL) which was then dried over Na_2SO_4 and evaporated under reduced pressure to afford 3-hydroxy-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid (100 g, 339 mmol, 77%) as a pale brown solid.


M.p. $113 - 115^{\circ}\text{C}$; LCMS (System 2, ESI) $R_t = 2.21$ $[\text{M}-\text{H}]^- = 259.1$, 88% purity; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) $\delta = 12.45 - 11.90$ (br. s., 1H), 7.77 – 7.68 (m, 4H), 2.85 – 2.74 (m, 1H), 2.67 – 2.53 (m, 4H) 1.92 (s, 1H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) $\delta = 175.7$, 151.6, 127.3 (q, $^2J_{\text{C-F}} = 31.5$ Hz), 125.7, 124.9 (q, $^3J_{\text{C-F}} = 3.6$ Hz), 124.3 (q, $^1J_{\text{C-F}} = 271.4$ Hz), 70.8, 40.9, 28.5; ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) $\delta = -61.3$ (s, 3F); $\nu_{\text{max}}/\text{cm}^{-1}$ (thin film) 3208 (br), 2957, 1701, 1683.

3-Chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid 8

To 3-hydroxy-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid (16.18 g, 62.2 mmol) in toluene (300 mL) was added conc. HCl (50 mL, 617 mmol) at ambient temperature and the reaction was sonicated for 7 h. The layers were separated and the aqueous layer was extracted with toluene (2×50 mL). The organics were combined and washed with water (2×50 mL) and brine (2×50 mL). The organics were filtered through a hydrophobic frit and evaporated under reduced pressure to give 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid (13.1 g, 46.9 mmol, 75%) as a white solid. The isomers can be separated *via* the


following procedure. The mixture was triturated with hexane and filtered to collect the solid washing with hexane. The solid was dried to afford *cis*-3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid as an off-white solid. The filtrate was evaporated under reduced pressure to afford *trans*-3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid as an off-white solid.

***cis*-3-Chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid – 8**




M.p. 172 – 174 °C; LCMS (Method B, UV, ESI) $R_t = 0.72$ $[M-H]^-$ = No mass ion peak, 100% purity; 1H NMR (400 MHz, DMSO- d_6) $\delta = 12.50$ (br. s., 1H), 7.88 – 7.74 (m, 4H), 3.34 – 3.21 (m, 2H), 3.08 – 2.97 (m, 2H), 2.92 – 2.80 (m, 1H); ^{13}C NMR (101 MHz, DMSO- d_6) $\delta = 174.5, 147.8, 128.5$ (q, $^2J_{C-F} = 31.5$ Hz), 127.0, 125.2 (q, $^3J_{C-F} = 3.6$ Hz), 124.0 (q, $^1J_{C-F} = 272.2$ Hz), 65.0, 41.9, 31.1; ^{19}F NMR (376 MHz, $CDCl_3$) $\delta = -62.8$ (s, 3F); ν_{max}/cm^{-1} (thin film) 2989, 1687; HRMS: Calculated for $C_{12}H_9ClF_3O_2$ 277.0249 Found $[M-H]^-$: 277.0251 (0.8 ppm).

***trans*-3-Chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid – 8**



M.p. 114 – 116 °C; LCMS (Method B, UV, ESI) $R_t = 0.71$ $[M-H]^- = 277.4$, 81% purity; 1H NMR (400 MHz, $CDCl_3$) $\delta = 7.66$ (d, $J = 8.3$ Hz, 2H), 7.49 (d, $J = 8.1$ Hz, 2H), 3.78 (quin, $J = 8.8$ Hz, 1H), 3.15 – 3.02 (m, 4H); ^{13}C NMR (101 MHz, $CDCl_3$) $\delta = 179.6, 148.3, 130.3$ (q, $^2J_{C-F} = 32.8$ Hz), 125.7 (q, $^3J_{C-F} = 4.0$ Hz), 125.5, 123.8 (q, $^1J_{C-F} = 272.3$ Hz), 66.8, 41.5, 32.5; ^{19}F NMR (376 MHz, $CDCl_3$) $\delta = -63.2$ (s, 3F); ν_{max}/cm^{-1} (thin film) 2947, 1705; HRMS: Calculated for $C_{12}H_{11}ClF_3O_2$ 279.0394 Found $[M+H]^+$: 279.0401 (2.3 ppm).

Methyl 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylate – 9




To a solution of 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylic acid (10 g, 35.9 mmol) and HCl (4M in 1,4-dioxane) (45 mL, 180 mmol) in 1,4-dioxane (100 mL) stirred under nitrogen at ambient temperature was added methanol (100 mL, 2472 mmol). The reaction mixture was stirred at ambient temperature for 3 h. Saturated aq. NaHCO₃ (20 mL) and ethyl acetate (20 mL) were added and the layers were separated. The aqueous layer was extracted with ethyl acetate (3 × 20 mL) and the combined organics were evaporated under reduced pressure to give methyl 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylate (10.5 g, 35.9 mmol, quant.) as a pale yellow liquid.

Analogously, the isomers of **9** could be produced using isomerically pure starting materials.


Data provided of individual isomers:

***cis*-Methyl 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylate – 9**




M.p. 52 – 54 °C; LCMS (Method B, UV, ESI) R_t = 1.29 [M-H]⁻ = No mass ion peak, 100% purity; ¹H NMR (400 MHz, CDCl₃) δ = 7.71 – 7.63 (m, 4H), 3.77 (s, 3H), 3.28 – 3.16 (m, 4H), 2.95 – 2.84 (m, 1H); ¹³C NMR (101 MHz, CDCl₃) δ = 173.6, 147.3, 130.3 (q, ²J_{C-F} = 33.1 Hz), 126.3, 125.7 (q, ³J_{C-F} = 3.6 Hz), 123.8 (q, ¹J_{C-F} = 272.2 Hz), 63.4, 52.2, 42.4, 31.8; ¹⁹F NMR (376 MHz, CDCl₃) δ = -62.7 (s, 3F); ν_{max}/cm⁻¹ (thin film) 2957, 1739.

***trans*-Methyl 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylate – 9**



LCMS (Method B, UV, ESI) $R_t = 1.31$ $[M+H]^+ =$ No mass ion peak, 80% purity; 1H NMR (400 MHz, $CDCl_3$) $\delta = 7.66$ (d, $J = 8.3$ Hz, 2H), 7.50 (d, $J = 8.3$ Hz, 2H), 3.80 – 3.67 (m, 4H), 3.14 – 2.98 (m, 4H); ^{13}C NMR (101 MHz, $CDCl_3$) $\delta = 174.1$, 148.5, 130.2 (q, $^2J_{C-F} = 32.8$ Hz), 125.7 (q, $^3J_{C-F} = 4.0$ Hz), 125.5, 123.8 (q, $^1J_{C-F} = 272.3$ Hz), 67.1, 52.0, 41.6, 32.6; ^{19}F NMR (376 MHz, $CDCl_3$) $\delta = -63.2$ (s, 3F); ν_{max}/cm^{-1} (thin film) 2955, 1733; HRMS: Calculated for $C_{13}H_{13}ClF_3O_2$ 293.0551 Found $[M+H]^+$: 293.0551 (0.1 ppm).


Methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.0]butane-1-carboxylate – 10



To solid sodium hydride (60% dispersion in oil) (8.2 g, 205 mmol) in THF (250 mL) stirred under nitrogen at ambient temperature was added a solution of *cis*-methyl 3-chloro-3-(4-(trifluoromethyl)phenyl)cyclobutanecarboxylate (50 g, 171 mmol) in THF (250 mL) portionwise. The reaction mixture was stirred at ambient temperature for 3 h. To the reaction mixture was added $NH_4Cl_{(aq)}$ (100 mL) and ethyl acetate (350 mL) then water (100 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (3 x 50 mL). The organics were combined and dried through a hydrophobic frit. The solvent was evaporated under reduced pressure to give methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.0]butane-1-carboxylate (43.1 g, 168 mmol, 98%) as a yellow solid.

M.p. 102 – 104, LCMS (Method B, UV, ESI) $R_t = 1.21$ $[M+H]^+ =$ No mass ion peak, 100% purity; 1H NMR (400 MHz, $CDCl_3$) $\delta = 7.56$ (d, $J = 8.3$ Hz, 2H), 7.39 (d, $J = 8.1$ Hz, 2H), 3.51 (s, 3H), 2.96 (t, $J = 1.1$ Hz, 2H), 1.67 (t, $J = 1.1$ Hz, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) $\delta = 169.4$, 138.3, 129.0 (q, $^2J_{C-F} = 33.1$ Hz), 126.1, 125.4 (q, $^3J_{C-F} = 43.3$ Hz), 124.1 (q, $^1J_{C-F} = 272.1$ Hz), 52.0, 36.0, 31.6, 24.4; ^{19}F NMR (376 MHz, $CDCl_3$) $\delta = -62.5$ (s, 3F); ν_{max}/cm^{-1} (thin film) 2956, 1708; HRMS: Calculated for $C_{13}H_{12}F_3O_2$ 257.0784 Found $[M+H]^+$: 257.0785 (0.5 ppm).


Methyl 2,2-dichloro-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate – 11



Methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.0]butane-1-carboxylate (5 g, 19.51 mmol) was dissolved in diethylene glycol dimethyl ether (10 mL) and tetrachloroethylene (PCE) (30 mL) and the reaction mixture was heated to 120 °C. Solid sodium 2,2,2-trichloroacetate (18.09 g, 98 mmol) was added portionwise over 30 min and the reaction was stirred at 140 °C for an additional 30 min. Water (100 mL) and CH₂Cl₂ (100 mL) were added and the layers were separated. The organic layer was washed with brine (10 × 50 mL) and was then filtered through a hydrophobic frit. Added CH₂Cl₂ (100 mL) and this was washed with brine (5 × 50 mL) and then 5% LiCl_(aq) (5 × 50 mL) to remove diethylene glycol dimethyl ether. The organic layer was filtered through a hydrophobic frit and the solvent was removed under reduced pressure to give the crude product. This was purified using column chromatography eluting with 0-20 cyclohexane:TBME. Collected fractions containing product were evaporated under reduced pressure to give methyl 2,2-dichloro-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate (2.5 g, 7.37 mmol, 38%) as a yellow oil.

LCMS (Method B, UV, ESI) $R_t = 1.38$ [M-H]⁻ = No mass ion peak, 81% purity; ¹H NMR (600 MHz, CDCl₃) $\delta = 7.66$ (d, $J = 8.1$ Hz, 2H), 7.51 (d, $J = 7.9$ Hz, 2H), 3.85 (s, 3H), 3.14 (d, $J = 1.3$ Hz, 2H), 2.47 (d, $J = 1.3$ Hz, 2H); ¹³C NMR (150 MHz, CDCl₃) $\delta = 165.4$, 135.9 (q, $^4J_{C-F} = 1.4$ Hz), 130.9 (q, $^2J_{C-F} = 32.6$ Hz), 127.6, 125.5 (q, $^3J_{C-F} = 3.8$ Hz), 123.9 (q, $^1J_{C-F} = 272.3$ Hz), 94.0, 57.0, 52.5, 52.4, 49.4; ¹⁹F NMR (376 MHz, CDCl₃) $\delta = -63.2$ (s, 3F); $\nu_{\text{max}}/\text{cm}^{-1}$ (thin film) 2957, 1737; HRMS: Calculated for C₁₄H₁₃Cl₂F₃O₂ 339.0161 Found [M+H]⁺: 339.0169 (2.5 ppm).


Methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate – 12



To azobis(cyclohexanecarbonitrile) (ACHN) (94 mg, 0.39 mmol) was added methyl 2,2-dichloro-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate (2.3 g, 6.78 mmol) in toluene (50 mL). Then tris(trimethylsilyl)silane (TTMSS) (11 mL, 38.9 mmol) was added. The reaction was heated under reflux for 3 h. ACHN (1.9 g, 7.77 mmol) was added portionwise over 4 days with continued heating under reflux. The solvent was removed under reduced pressure and the crude product was purified using column chromatography eluting with 0-20% cyclohexane:TBME. Fractions containing product were impure and these were purified using column chromatography eluting with 0-10% cyclohexane:TBME. Fractions containing product were evaporated under reduced pressure to give methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate (1.36 g, 5.03 mmol, 74%) as a white solid.

M.p. 54 – 56 °C; LCMS (Method B, UV, ESI) $R_t = 1.31$ $[M+H]^+ =$ No mass ion peak, 100% purity; ^1H NMR (400 MHz, CDCl_3) $\delta = 7.59$ (d, $J = 8.1$ Hz, 2H), 7.34 (d, $J = 7.8$ Hz, 2H), 3.75 (s, 3H), 2.38 (s, 6H); ^{13}C NMR (101 MHz, CDCl_3) $\delta = 170.3$, 143.5 (q, $^4J_{\text{C-F}} = 1.5$ Hz), 129.2 (q, $^2J_{\text{C-F}} = 32.3$ Hz), 126.5, 125.2 (q, $^3J_{\text{C-F}} = 3.7$ Hz), 124.2 (q, $^1J_{\text{C-F}} = 272.2$ Hz), 53.4, 51.7, 41.5, 37.1; ^{19}F NMR (376 MHz, CDCl_3) $\delta = -62.5$ (s, 3F); $\nu_{\text{max}}/\text{cm}^{-1}$ (thin film) 2982, 1737; HRMS: Calculated for $\text{C}_{14}\text{H}_{14}\text{F}_3\text{O}_2$ 271.0940 Found $[M+H]^+$: 271.0944 (1.4 ppm).


3-(4-(Trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid – 13



To methyl 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylate (1.3 g, 4.81 mmol) in 1,4-dioxane (10 mL) was added lithium hydroxide (9.62 mL, 9.62 mmol). This was stirred at ambient temperature for 3h. To the reaction was added 2M HCl_(aq) (5 mL) and ethyl acetate (30 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (3 x 10 mL). The organic layers were combined and evaporated under reduced pressure to give 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid (1.17 g, 4.57 mmol, 95%) as a white solid.

M.p. 233 – 235 °C; LCMS (Method B, UV, ESI) $R_t = 0.73$ [M-H]⁻ = 255.6, 98% purity; ¹H NMR (400 MHz, DMSO-d₆) $\delta = 12.60 - 12.27$ (m, 1H), 7.68 (d, $J = 8.1$ Hz, 2H), 7.47 (d, $J = 7.8$ Hz, 2H), 2.28 (s, 6H); ¹³C NMR (101 MHz, DMSO-d₆) $\delta = 170.9, 144.2, 127.5$ (q, $^2J_{C-F} = 32.8$ Hz), 126.9, 125.1 (q, $^3J_{C-F} = 3.7$ Hz), 124.3 (q, $^1J_{C-F} = 272.2$ Hz), 52.6, 40.6, 36.8; ¹⁹F NMR (376 MHz, DMSO-d₆) $\delta = -60.9$ (s, 3F); $\nu_{\max}/\text{cm}^{-1}$ (thin film) 2939, 1697; HRMS: Calculated for C₁₃H₁₂F₃O₂ 257.0784 Found [M+H]⁺: 257.0789 (2.0 ppm).


***N*-(2-(Diethylamino)ethyl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide – 15**



T3P (50% in ethyl acetate) (2.8 mL, 4.70 mmol), 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid (800 mg, 3.12 mmol) and triethylamine (0.9 mL, 6.46 mmol) was stirred in ethyl acetate (10 mL) for 30 min. Then *N*1,*N*1-diethylethane-1,2-diamine (0.66 mL, 4.70 mmol) was added and the reaction mixture was stirred for 16 h. Water (15 mL) and CH₂Cl₂ (20 mL) were added and the layers separated. The aqueous was extracted with CH₂Cl₂ (3 x 5 mL) and the organics were combined and evaporated under reduced pressure to give *N*-(2-(diethylamino)ethyl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide (1.10 g, 3.10 mmol, 99%) as a orange solid.

M.p. 81 – 83 °C; LCMS (Method B, UV, ESI) *R*_t = 1.17 [M+H]⁺ = 355.5, 94% purity; ¹H NMR (400 MHz, CDCl₃) δ = 7.58 (d, *J* = 7.8 Hz, 2H), 7.34 (d, *J* = 8.1 Hz, 2H), 6.48 – 6.36 (m, 1H), 3.39 – 3.27 (m, 2H), 2.61 – 2.52 (m, 6H), 2.32 (s, 6H), 1.05 (t, *J* = 7.1 Hz, 6H); ¹³C NMR (101 MHz, CDCl₃) δ = 169.7, 143.7, 129.2 (q, ²*J*_{C-F} = 33.0 Hz), 126.5, 125.2 (q, ³*J*_{C-F} = 3.7 Hz), 124.1 (q, ¹*J*_{C-F} = 272.2 Hz), 52.8, 51.3, 47.0, 40.5, 38.8, 36.7, 12.1; ¹⁹F NMR (376 MHz, CDCl₃) δ = -62.5 (s, 3F); ν_{max}/cm⁻¹ (thin film) 3320 (br) 2975, 1642; HRMS: Calculated for C₁₉H₂₆F₃N₂O 355.1992 Found [M+H]⁺: 355.2000 (2.2 ppm).


N1,N1-Diethyl-N2-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)ethane-1,2-diamine – 16



To *N*-(2-(diethylamino)ethyl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide (978 mg, 2.76 mmol) in THF (30 mL) was added LiAlH₄ (2.3 M in 2-MeTHF) (2.4 mL, 5.52 mmol) at ambient temperature and the reaction mixture was stirred for 16 h. The reaction was quenched by the addition of wet THF (20 mL) initially dropwise then portionwise. Water (20 mL) was then added portionwise, followed by 2M HCl (50 mL) and ethyl acetate (50 mL). The layers were separated and the organic layer was extracted with 2M HCl (3 x 10 mL). The aqueous layers were combined and basified with 2M NaOH (100 mL) then extracted with CH₂Cl₂ (10 x 50 mL). The organics were combined and filtered through a hydrophobic frit and evaporated under reduced pressure to give the crude *N1,N1*-diethyl-*N2*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)ethane-1,2-diamine (527 mg, 1.55 mmol, 56%). LCMS showed 87% product and this was taken on as is to the next step without further purification.

LCMS (Method B, UV, ESI) $R_t = 1.29$ $[M+H]^+ = 340.6$, 87% purity.

***N*-(2-(Diethylamino)ethyl)-2-(2-((4-fluorobenzyl)thio)-4-oxo-4,5,6,7-tetrahydro-1*H*-cyclopenta[d]pyrimidin-1-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide, Formic acid salt – 5**



2-(2-((4-Fluorobenzyl)thio)-4-oxo-4,5,6,7-tetrahydro-1*H*-cyclopenta[d]pyrimidin-1-yl)acetic acid (634 mg, 1.90 mmol), triethylamine (0.350 mL, 2.509 mmol) and T3P (50% in ethyl acetate) (1.1 mL, 1.85 mmol) were stirred in CH₂Cl₂ (5 mL) for 30 min. Then *N*1,*N*1-diethyl-*N*2-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)ethane-1,2-diamine (427 mg, 1.25 mmol) in CH₂Cl₂ (5 mL) was added and this was stirred for 16 h. To the reaction mixture was added 2M NaOH_(aq) (20 mL), water (20 mL) and CH₂Cl₂ (40 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (10 x 15 mL). The organics were combined and evaporated under reduced pressure. The crude product was purified by HpH MDAP, Method D to give product contaminated with formic acid and HpH modifier. The product was purified using HpH MDAP (Method D) and the solvent was removed under reduced pressure to give *N*-(2-(diethylamino)ethyl)-2-(2-((4-fluorobenzyl)thio)-4-oxo-4,5,6,7-tetrahydro-1*H*-cyclopenta[d]pyrimidin-1-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide, Formic acid salt (526 mg, 0.75 mmol, 60%) as a yellow solid.


M.p. 88 – 90 °C; LCMS (Method B, UV, ESI) $R_t = 1.44$ $[M+H]^+ = 657.7$, 100% purity; ¹H NMR (400 MHz, DMSO-*d*₆, 373 K) δ = 8.14 (s, 1H), 7.64 (d, $J = 8.1$ Hz, 2H), 7.47 – 7.41 (m, 2H), 7.41 – 7.36 (m, 2H), 7.03 (br. s., 2H), 5.00 – 4.80 (m, 2H), 4.46 (s, 2H), 3.57 (br. s., 2H), 3.40 (t, $J = 6.5$ Hz, 2H), 2.79 (t, $J = 7.6$ Hz, 2H), 2.67 – 2.52 (m, 8H), 2.14 – 1.95 (m, 8H), 0.98 (t, $J = 7.1$ Hz, 6H); ¹³C NMR (101 MHz, CDCl₃) δ = 167.6, 167.1, 164.9, 163.5, 160.7, 156.0, 155.6, 144.2, 143.2, 131.6, 131.1, 130.1, 126.3, 125.3, 125.2, 121.4, 121.2, 115.7, 115.4, 115.4, 52.4, 52.2, 51.4, 49.7, 49.6, 48.8, 47.8, 47.4, 47.1, 46.4, 43.8, 42.3, 41.8, 37.2, 37.0, 36.3, 36.3, 32.0, 31.9, 28.3, 20.8, 20.7, 11.6, 9.8; ¹⁹F NMR (376 MHz, CDCl₃) δ =

-62.5 (s, 3F), -114.2 (s, 1F); $\nu_{\text{max}}/\text{cm}^{-1}$ (thin film) 2969, 1618; HRMS: Calculated for $\text{C}_{35}\text{H}_{41}\text{F}_4\text{N}_4\text{O}_2\text{S}$ 657.2881 Found $[\text{M}+\text{H}]^+$: 657.2883 (0.4 ppm).

N-(2-(diethylamino)ethyl)-2-(2-((4-fluorobenzyl)thio)-4-oxo-4,5,6,7-tetrahydro-1*H*-cyclopenta[d]pyrimidin-1-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide, Formic acid salt (100 mg,) was dissolved in MeOH (2 mL) then passed through an SAX cartridge eluting with MeOH (3 CVs). This collected fractions were evaporated under reduced pressure to give *N*-(2-(diethylamino)ethyl)-2-(2-((4-fluorobenzyl)thio)-4-oxo-4,5,6,7-tetrahydro-1*H*-cyclopenta[d]pyrimidin-1-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide (87 mg, 0.132 mmol) as a yellow solid.

LCMS (Method B, UV, ESI) $R_t = 1.45$ $[\text{M}+\text{H}]^+ = 657.3$, 100% purity; ^1H NMR (400 MHz, CDCl_3) $\delta = 7.58$ (dd, $J = 3.5, 7.9$ Hz, 2H), 7.40 – 7.29 (m, 2H), 7.25 (t, $J = 8.1$ Hz, 2H), 7.02 – 6.94 (m, 1H), 6.89 – 6.81 (m, 1H), 4.92 – 4.64 (m, 2H), 4.53 (d, $J = 4.2$ Hz, 2H), 3.63 – 3.52 (m, 3H), 3.40 – 3.31 (m, 1H), 2.92 – 2.75 (m, 6H), 2.62 – 2.52 (m, 3H), 2.17 – 1.94 (m, 9H), 1.19 – 1.06 (m, 3H), 1.02 (t, $J = 7.2$ Hz, 3H).

***N*-(1-(2-Methoxyethyl)piperidin-4-yl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide – 19**




T3P (50% in DMF) (0.68 mL, 1.17 mmol), 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid (200 mg, 0.781 mmol) and Et_3N (0.22 mL, 1.58 mmol) were stirred in CH_2Cl_2 (1 mL) for 20 min. Then 1-(2-methoxyethyl)piperidin-4-amine (0.194 mL, 1.17 mmol) was added and the reaction mixture was stirred for 3 h. Water (2 mL) and CH_2Cl_2 (5 mL) were added and the layers separated. The organic layer was washed with 5% aq. LiCl (2 x 2 mL). The combined aqueous layers were extracted with DCM (2 x 3 mL). The organics were combined and filtered through a

hydrophobic frit. The solvent was removed under reduced pressure to give *N*-(1-(2-methoxyethyl)piperidin-4-yl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide (228 mg, 0.58 mmol, 74% yield) as an off white solid.

M.p. 185 – 187 °C; LCMS (Method B, UV, ESI) $R_t = 1.07$ $[M+H]^+ = 397.2$, 100% purity; 1H NMR (400 MHz, DMSO- d_6) $\delta = 7.68$ (d, $J = 8.1$ Hz, 2H), 7.62 (d, $J = 7.8$ Hz, 1H), 7.45 (d, $J = 8.1$ Hz, 2H), 3.58 - 3.47 (m, 1H), 3.41 (t, $J = 6.0$ Hz, 2H), 3.23 (s, 3H), 2.84 (m, 2H), 2.45 (t, $J = 6.0$ Hz, 2H), 2.22 (s, 6H), 2.04 – 1.94 (m, 2H), 1.70 – 1.61 (m, 2H), 1.53 – 1.40 (m, 2H); ^{13}C NMR (101 MHz, DMSO- d_6) $\delta = 168.2$, 144.7, 127.3 (q, $^2J_{C-F} = 31.5$ Hz), 126.8, 125.1 (q, $^3J_{C-F} = 3.8$ Hz), 124.3 (q, $^1J_{C-F} = 271.4$ Hz), 70.1, 57.9, 57.0, 52.8, 52.3, 46.2, 38.3, 31.5 (1C concealed); ^{19}F NMR (376 MHz, $CDCl_3$) $\delta = -62.5$ (s, 3F); ν_{max}/cm^{-1} (thin film) 3345, 2942, 1637; HRMS: Calculated for $C_{21}H_{28}F_3N_2O_2$ 397.2097 Found $[M+H]^+$: 397.2096 (–0.4 ppm).

1-(2-Methoxyethyl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)piperidin-4-amine, formic acid salt – 20




To $[Ir(COE)_2Cl]_2$ (48.4 mg, 0.054 mmol) was added diethylsilane (0.210 mL, 1.62 mmol) and CH_2Cl_2 (0.5 mL). To this was added *N*-(1-(2-methoxyethyl)piperidin-4-yl)-3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxamide (107 mg, 0.27 mmol) in CH_2Cl_2 (1.5 mL). The reaction mixture was stirred for 20 h at ambient temperature. To the reaction mixture was added 4M HCl in 1,4-dioxane (3 mL), however no solid precipitation. 2M NaOH(aq) was added until alkaline pH achieved. CH_2Cl_2 (20 mL) was added and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (3 x 10 mL). The organic layers were combined and evaporated under reduced pressure to give a crude product. The crude product LCMS showed only desired product. NMR showed diethylsilane peaks. To the crude product was added CH_2Cl_2 (10 mL) and 2M HCl (10 mL) and the layers were

separated. The organic layer was extracted with 2M HCl (3 x 5 mL). The aqueous layers were combined and 2M NaOH_(aq) was added until alkaline pH. CH₂Cl₂ (30 mL) was added and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (6 x 20 mL) and the organics were combined and evaporated under reduced pressure to give a residue which was purified using HpH MDAP (Method C) to give 1-(2-methoxyethyl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)piperidin-4-amine, formic acid salt (61 mg, 0.16 mmol, 59%) as a yellow solid.

M.p. 118 – 120 °C; LCMS (Method B, UV, ESI) $R_t = 1.33$ $[M+H]^+ = 383.2$, 100% purity; ¹H NMR (400 MHz, CDCl₃) $\delta = 8.46$ (s, 1H), 7.54 (d, $J = 8.1$ Hz, 2H), 7.31 – 7.27 (m, 2H), 5.23 (br. s., 2H), 3.61 (t, $J = 5.1$ Hz, 2H), 3.35 (s, 3H), 3.26 – 3.19 (m, 2H), 3.05 – 2.96 (m, 3H), 2.80 (t, $J = 5.3$ Hz, 2H), 2.47 – 2.37 (m, 2H), 2.17 – 2.10 (m, 8H), 1.95 – 1.82 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) $\delta = 167.5$, 144.0, 128.9 (q, $^2J_{C-F} = 32.3$ Hz), 126.4, 125.1 (q, $^3J_{C-F} = 3.6$ Hz), 69.2, 58.8, 56.8, 53.6, 52.2, 51.6, 45.6, 42.0, 36.4, 28.3 (1C concealed); ¹⁹F NMR (376 MHz, CDCl₃) $\delta = -62.4$ (s, 3F); ν_{max}/cm^{-1} (thin film) 2782 (br), 1619, 1560; HRMS: Calculated for C₂₁H₃₀F₃N₂O 383.2305 Found $[M+H]^+$: 383.2307 (0.5 ppm).

2-(2-((2,3-Difluorobenzyl)thio)-4-oxoquinolin-1(4*H*)-yl)-*N*-(1-(2-methoxyethyl)piperidin-4-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide – 22



2-(2-((2,3-Difluorobenzyl)thio)-4-oxoquinolin-1(4*H*)-yl)acetic acid (28.3 mg, 0.078 mmol), Et₃N (0.015 mL, 0.11 mmol) and T3P (50% in DMF) (0.046 mL, 0.078 mmol) in CH₂Cl₂ (0.5 mL) was stirred for 15 min at ambient temperature. To the reaction mixture was added 1-(2-methoxyethyl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)piperidin-4-amine (20 mg, 0.052 mmol) in CH₂Cl₂ (0.500 mL) at ambient

temperature stirred for 64 h. Water (1 mL) was added and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (3 x 1 mL). The organic layers were combined and evaporated under reduced pressure give a crude product. This was purified by HpH MDAP (Method D) to give 2-(2-((2,3-difluorobenzyl)thio)-4-oxoquinolin-1(4*H*)-yl)-*N*-(1-(2-methoxyethyl)piperidin-4-yl)-*N*-((3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)methyl)acetamide (20 mg, 0.028 mmol, 53% yield) as a yellow solid.

M.p. 57 – 59 °C; LCMS (Method B, UV, ESI) $R_t = 1.36$ [M+H]⁺ = 726.2, 100% purity; ¹H NMR (400 MHz, DMSO-d₆, 373 K) δ = 8.18 (dd, J = 1.5, 8.1 Hz, 1H), 7.71 – 7.59 (m, 3H), 7.47 – 7.32 (m, 4H), 7.32 – 7.21 (m, 2H), 7.17 – 7.07 (m, 1H), 6.35 (s, 1H), 5.41 (br. s., 2H), 4.43 (s, 2H), 3.83 – 3.72 (m, 1H), 3.57 (br. s., 2H), 3.48 (t, J = 5.7 Hz, 2H), 3.28 (s, 3H), 3.00 (m, 2H), 2.56 (t, J = 5.6 Hz, 2H), 2.27 – 1.68 (m, 12H); ¹³C NMR (101 MHz, CDCl₃) δ = 176.5, 176.4, 166.5, 165.2, 152.1, 152.0, 142.3, 132.6, 132.4, 127.0, 126.9, 126.4, 126.3, 125.9, 125.3, 125.1, 124.5, 123.9, 117.5, 117.4, 115.3, 115.1, 113.3, 113.1, 77.2, 69.9, 69.1, 58.9, 58.9, 57.5, 56.9, 54.8, 53.2, 53.0, 52.9, 52.8, 52.2, 50.1, 49.9, 44.7, 43.4, 41.7, 41.0, 38.1, 37.8, 33.1, 33.0, 30.8, 28.3; ¹⁹F NMR (376 MHz, DMSO-d₆) δ = –60.79 (s, 3F), –138.67 (m, 1F), –142.15 (m, 1F); $\nu_{\text{max}}/\text{cm}^{-1}$ (thin film) 3402 (br), 2964, 1652, 1617, 1593; HRMS: Calculated for C₃₉H₄₁F₅N₃O₃S 726.2783 Found [M+H]⁺: 726.2796 (1.8 ppm).

X-ray crystallography

Materials and methods

Lp-PLA2 (residues 46-428 with a C-terminal His6 tag) was expressed, purified and crystallised essentially as previously described¹ though the purified protein was concentrated to 15mg/ml for crystallisation. To obtain the protein complex structure with **analogue-5**, a crystal was soaked in 30% PEG3350, 0.1 M HEPES pH 7.3, 0.2 M NaCl, 0.04% pluronic F68, 10 mM CHAPS and 2mM compound dissolved in DMSO for ~1 day.

The crystal was cryoprotected by briefly soaking in well buffer with 10% glycerol. A diffraction dataset was acquired at 100 K at the ESRF (station ID29) using a Pilatus 6M detector, and processed by the synchrotron autoprocessing suite (XDS_PARALLELPROC)² using XDS³. Data were merged using AIMLESS⁴ within the CCP4 programming suite⁵.

The crystal structure was solved by Fourier Synthesis using REFMAC⁶ and a previously determined in-house Lp-PLA2 crystal structure as the starting model. Refinement was carried out with REFMAC and model building using COOT⁷. The final R_{factor} (and R_{free}) achieved were 16.2% (and 19.2%) and the structure was deposited into the PDB as entry 5LP1.

References

- ¹ Woolford, A. J.-A. *et al.*, 2016. *J. Med. Chem.* **59**, 5356–5367.
- ² Monaco S *et al.*, 2013. *J Appl Cryst.* **46**, 804-810.
- ³ Kabsch W. 2010. *Acta Cryst.* **D66**, 125-132.
- ⁴ Evans, P.R. and Murshudov, G.N. 2013. *Acta Cryst.* **D69**, 1204-1214.
- ⁵ Winn, M.D. *et al.*, 2010. *Acta Cryst.* **D67**, 235-242.
- ⁶ Murshudov G.N. *et al.*, 1997. *Acta Cryst.* **D53**, 240-255.
- ⁷ Emsley P and Cowtan K. 2004. *Acta Cryst.* **D60**, 2126-2132.

Table of Statistics

Lp-PLA2	Analogue 5
PARAMETER ^a	
DATA COLLECTION	
Space Group	C2
Cell Dimensions	
a,b,c (Å)	100.28,91.56,51.65
α, β, γ (°)	90.00,111.88, 90.00
Resolution (Å)	1.91 (1.98)
R _{merge} ^b	0.094 (0.624)
Average I/σI	7.8 (1.7)
Completeness (%)	98.7 (92.5)
Redundancy	3.4 (3.2)
No. Reflections	112961 (9618)
No. Unique Reflections	33162 (3045)
REFINEMENT	
Resolution (Å)	20.00-1.91
R _{work} /R _{free}	0.162/0.192
No. Reflections	31474
No. atoms	
Protein (chain A)	3040
Cmpd (L)	46
Water (W)	332
Cl (C)	2
B-factors [Å ²]	
Protein (chain A)	35.3
Cmpd (L)	37.0
Water (W)	47.4
Cl (C)	36.5
R.m.s deviations	
Bond lengths (Å)	0.007
Bond angles (°)	1.386

^a Data for the highest resolution shell are given in parentheses.

^b R_{merge} = Σ|I_j - <I_j>|/Σ<I_j>.