

Table S4. Patients taking pain and psychoactive drugs

	Baseline			3 Months			6 Months			12 Months		
	CM n=47	UC n=48	<i>P</i>	CM n=45	UC n=46	<i>P</i>	CM n=45	UC n=45	<i>P</i>	CM n=45	UC n=42	<i>P</i>
	n (%)	n (%)		n (%)	n (%)		n (%)	n (%)		n (%)	n (%)	
Opiate	1 (2)	4 (8)	.36	0	3 (7)	.24	0	2 (4)	.49	0	1 (2)	.48
Hypnotic	1 (2)	2 (4)	1	3 (7)	1 (2)	.36	3 (7)	1 (2)	.61	2 (4)	3 (7)	.67
Antidepressants	8 (17)	5 (10)	.38	9 (20)	8 (17)	.79	12 (27)	6 (13)	.19	8 (18)	8 (19)	1
Pregabalin	2 (4)	2 (4)	1	2 (4)	3 (7)	1	2 (4)	3 (7)	1	1 (2)	2 (5)	.61
Benzodiazepine	3 (6)	2 (4)	.68	3 (7)	4 (9)	1	1 (2)	4 (9)	.36	0	1 (2)	.48

CM: Case Management; UC: Usual Care; IQR: interquartile range